

Thue-Morse sequences of squares in compact groups

Johannes Morgenbesser

joint work with Michael Drmota

Institut für Diskrete Mathematik und Geometrie
Technische Universität Wien



Institut de Mathématiques de Luminy
Université de la Méditerranée, Aix Marseille II



Workshop on digital expansions, dynamics and tilings
Aussois, April 4-11, 2010

Summary

- ★ Thue-Morse sequence
- ★ Generalized Thue-Morse sequence and main result
- ★ Sketch of the proof
- ★ Applications
 - ★ La somme des chiffres des carrés
 - ★ Automatic sequences

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

0

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

01

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

0110

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

01101001

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

0110100110010110

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

01101001100101101001011001101001

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101 ...

$$t_0 = 0, \quad t_{2^n+k} = 1 - t_k \quad (0 \leq k < 2^n)$$

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101 ...

$$t_0 = 0, \quad t_{2^n+k} = 1 - t_k \quad (0 \leq k < 2^n)$$

$$t_n = s_2(n) \bmod 2$$

$$n = \sum_{i=0}^{\ell-1} \varepsilon_i(n) q^i \quad \varepsilon_i(n) \in \{0, 1, \dots, q-1\}, \quad s_q(n) = \sum_{i=0}^{\ell-1} \varepsilon_i(n)$$

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

$$\#\{0 \leq n < N : t_n = 0\} \approx \frac{N}{2}$$

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

$$\# \{0 \leq n < N : t_n = 0\} \approx \frac{N}{2}$$

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

$$\#\{0 \leq n < N : t_n = 0\} \approx \frac{N}{2}$$

$$\#\{0 \leq n < N : t_{3n} = 0\} \approx \frac{N}{2}$$

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

0 0 0 0 0 0 0 0 1 0 0 0 0 0 0 0 1 ...

$$\#\{0 \leq n < N : t_n = 0\} \approx \frac{N}{2}$$

$$\#\{0 \leq n < N : t_{3n} = 0\} \approx \frac{N}{2}$$

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

$$\# \{0 \leq p < N, p \text{ prime} : t_p = 0\}$$

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

Mauduit and Rivat (2009):

$$\# \{0 \leq p < N, p \text{ prime} : t_p = 0\} \approx \frac{\pi(N)}{2}$$

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

$$\# \{0 \leq n < N : t_{n^2} = 0\}$$

★ Thue-Morse sequence

Thue-Morse sequence $(t_n)_{n \geq 0}$:

011010011001011010010110011010011001011001101...

Mauduit and Rivat (2005+):

$$\#\{0 \leq n < N : t_{n^2} = 0\} \approx \frac{N}{2}$$

★ Generalized Thue-Morse sequences and main results

- H ... compact (Hausdorff) group
- $q \geq 2$ and $g_0, g_1, \dots, g_{q-1} \in H$ with $g_0 = e$ (identity element)
- $G \leq H$... closure of the subgroup generated by g_0, g_1, \dots, g_{q-1}

★ Generalized Thue-Morse sequences and main results

- H ... compact (Hausdorff) group
- $q \geq 2$ and $g_0, g_1, \dots, g_{q-1} \in H$ with $g_0 = e$ (identity element)
- $G \leq H$... closure of the subgroup generated by g_0, g_1, \dots, g_{q-1}

$$n = \sum_{i=0}^{\ell-1} \varepsilon_i(n) q^i$$

★ Generalized Thue-Morse sequences and main results

- H ... compact (Hausdorff) group
- $q \geq 2$ and $g_0, g_1, \dots, g_{q-1} \in H$ with $g_0 = e$ (identity element)
- $G \leq H$... closure of the subgroup generated by g_0, g_1, \dots, g_{q-1}

$$n = \sum_{i=0}^{\ell-1} \varepsilon_i(n) q^i$$

Generalized Thue-Morse sequence:

$$T(n) := g_{\varepsilon_0(n)} g_{\varepsilon_1(n)} \cdots g_{\varepsilon_{\ell-1}(n)}$$

★ Generalized Thue-Morse sequences and main results

- H ... compact (Hausdorff) group
- $q \geq 2$ and $g_0, g_1, \dots, g_{q-1} \in H$ with $g_0 = e$ (identity element)
- $G \leq H$... closure of the subgroup generated by g_0, g_1, \dots, g_{q-1}

$$n = \sum_{i=0}^{\ell-1} \varepsilon_i(n) q^i$$

Generalized Thue-Morse sequence:

$$T(n) := g_{\varepsilon_0(n)} g_{\varepsilon_1(n)} \cdots g_{\varepsilon_{\ell-1}(n)}$$

$$H = \langle \mathbb{Z}/2\mathbb{Z}, + \rangle, q = 2, g_0 = 0, g_1 = 1 : T(n) = s_2(n) \bmod 2 = t_n$$

Theorem (Drmota and Morgenbesser, 2010)

There exists a positive integer $m = m(q, g_0, \dots, g_{q-1})$ such that the following holds: Set

$$d\nu = \sum_{v=0}^m \mathbf{1}_{g_v U} \cdot Q(v, m) d\mu,$$

where

- $\mu \dots$ Haar measure on G ,
- $U = \text{cl}(\{T(mn) : n \geq 0\}) \dots$ normal subgroup of G of index m ,
- $Q(v, m) = \#\{0 \leq n < m : n^2 \equiv v \pmod{m}\}$.

Then $(T(n^2))_{n \geq 0}$ is ν -uniformly distributed in G , that is,

$$\frac{1}{N} \sum_{n=0}^{N-1} \delta_{T(n^2)} \rightarrow \nu.$$

Theorem (Drmota and Morgenbesser, 2010)

There exists a positive integer $m = m(q, g_0, \dots, g_{q-1})$ such that the following holds: Set

$$d\nu = \sum_{v=0}^m \mathbf{1}_{g_v U} \cdot Q(v, m) d\mu,$$

where

- $\mu \dots$ Haar measure on G ,
- $U = \text{cl}(\{T(mn) : n \geq 0\}) \dots$ normal subgroup of G of index m ,
- $Q(v, m) = \#\{0 \leq n < m : n^2 \equiv v \pmod{m}\}$.

Then $(T(n^2))_{n \geq 0}$ is ν -uniformly distributed in G , that is,

$$\frac{1}{N} \sum_{n=0}^{N-1} f(T(n^2)) \rightarrow \int_G f d\nu.$$

for all continuous functions $f : G \rightarrow \mathbb{C}$.

A **unitary group representation** is a continuous homomorphism $D : G \rightarrow U_n$ for some $n \geq 1$.

$U_n \dots$ group of unitary $n \times n$ matrices over \mathbb{C}

D is irreducible if there is no proper subspace W of \mathbb{C}^n with $D(x)W \subseteq W$ for all $x \in G$

A **unitary group representation** is a continuous homomorphism $D : G \rightarrow U_n$ for some $n \geq 1$.

$U_n \dots$ group of unitary $n \times n$ matrices over \mathbb{C}

D is irreducible if there is no proper subspace W of \mathbb{C}^n with $D(x)W \subseteq W$ for all $x \in G$

Lemma

Let G be a compact group and ν a regular normed Borel measure on G . Then a sequence $(x_n)_{n \geq 0}$ is ν -uniformly distributed in G iff

$$\frac{1}{N} \sum_{n=0}^{N-1} D(x_n) \rightarrow \int_G D \, d\nu$$

holds for all irreducible unitary representations D of G .

Remarks:

- The *characteristic integer* m is the largest integer such that $m \mid q - 1$ and such that there exists a representation D of G with

$$D_1(g_u) = e^{-2\pi i \frac{u}{m}} \quad \text{for all } u \in \{0, 1, \dots, q - 1\}.$$

- $(T(n^2))_{n \geq 0}$ is uniformly distributed in G (i.e., $\nu = \mu$) iff $m \leq 2$.
- If G is connected, then $T(n^2)$ is uniformly distributed in G .

★ Sketch of the proof

$$\frac{1}{N} \sum_{0 \leq n < N} D(T(n^2))$$

★ Sketch of the proof

$$\frac{1}{N} \sum_{0 \leq n < N} D(T(n^2))$$

- D_0, \dots, D_{m-1} :

$$D_k(g_u) = e^{-2\pi i \frac{k}{m} u} \quad \text{for all } 0 \leq u < q \text{ and } 0 \leq k < m$$

- all other irreducible unitary representations...

Van der Corput type inequality:

$$\left\| \sum_{0 \leq n < N} D(T(n^2)) \right\|_{\mathbb{F}} \leq \left(\frac{dN}{R} \sum_{|r| < R} \left(1 - \frac{|r|}{R} \right) \left\| \sum_{\substack{0 \leq n \leq B \\ 0 \leq n+r \leq B}} D(T(n+r)^2) D(T(n^2))^H \right\|_{\mathbb{F}} \right)^{1/2} + \frac{f}{2} R$$

Van der Corput type inequality:

$$\left\| \sum_{0 \leq n < N} D(T(n^2)) \right\|_{\mathbb{F}} \leq \left(\frac{dN}{R} \sum_{|r| < R} \left(1 - \frac{|r|}{R} \right) \left\| \sum_{\substack{0 \leq n \leq B \\ 0 \leq n+r \leq B}} D(T(n+r)^2) D(T(n^2))^H \right\|_{\mathbb{F}} \right)^{1/2} + \frac{f}{2} R$$

$$T(n) = g_{\varepsilon_0(n)} g_{\varepsilon_1(n)} \cdots g_{\varepsilon_{\lambda-1}(n)} g_{\varepsilon_{\lambda}(n)} \cdots g_{\varepsilon_{\ell-1}(n)}$$

Van der Corput type inequality:

$$\left\| \sum_{0 \leq n < N} D(T(n^2)) \right\|_{\mathbb{F}} \leq \left(\frac{dN}{R} \sum_{|r| < R} \left(1 - \frac{|r|}{R} \right) \left\| \sum_{\substack{0 \leq n \leq B \\ 0 \leq n+r \leq B}} D(T(n+r)^2) D(T(n^2))^H \right\|_{\mathbb{F}} \right)^{1/2} + \frac{f}{2} R$$

$$T_{\lambda}(n) = g_{\varepsilon_0(n)} g_{\varepsilon_1(n)} \cdots g_{\varepsilon_{\lambda-1}(n)}$$

Van der Corput type inequality:

$$\left\| \sum_{0 \leq n < N} D(T(n^2)) \right\|_{\mathbb{F}} \leq \left(\frac{dN}{R} \sum_{|r| < R} \left(1 - \frac{|r|}{R} \right) \left\| \sum_{\substack{0 \leq n \leq B \\ 0 \leq n+r \leq B}} D(T(n+r)^2) D(T(n^2))^H \right\|_{\mathbb{F}} \right)^{1/2} + \frac{f}{2} R$$

$$T_{\lambda}(n) = g_{\varepsilon_0(n)} g_{\varepsilon_1(n)} \cdots g_{\varepsilon_{\lambda-1}(n)}$$

$$(n+r)^2 = (\varepsilon_{\ell-1} \varepsilon_{\ell-2} \cdots \varepsilon_{\lambda} \cdots)_q, \quad n^2 = (\varepsilon_{\ell-1} \varepsilon_{\ell-2} \cdots \varepsilon_{\lambda} \cdots)_q$$

Van der Corput type inequality:

$$\left\| \sum_{0 \leq n < N} D(T(n^2)) \right\|_{\mathbb{F}} \leq \left(\frac{dN}{R} \sum_{|r| < R} \left(1 - \frac{|r|}{R} \right) \left\| \sum_{\substack{0 \leq n \leq B \\ 0 \leq n+r \leq B}} D(T(n+r)^2) D(T(n^2))^H \right\|_{\mathbb{F}} \right)^{1/2} + \frac{f}{2} R$$

$$T_{\lambda}(n) = g_{\varepsilon_0(n)} g_{\varepsilon_1(n)} \cdots g_{\varepsilon_{\lambda-1}(n)}$$

$$(n+r)^2 = (\varepsilon_{\ell-1} \varepsilon_{\ell-2} \cdots \varepsilon_{\lambda} \cdots)_q, \quad n^2 = (\varepsilon_{\ell-1} \varepsilon_{\ell-2} \cdots \varepsilon_{\lambda} \cdots)_q$$

$$D(T((n+r)^2)) D(T(n^2))^H$$

$$= D(T_{\lambda}((n+r)^2)) D(g_{\varepsilon_{\lambda}}) \cdots D(g_{\varepsilon_{\ell-1}}) D(g_{\varepsilon_{\ell-1}})^H \cdots D(g_{\varepsilon_{\lambda}})^H D(T_{\lambda}(n^2))^H$$

Van der Corput type inequality:

$$\left\| \sum_{0 \leq n < N} D(T(n^2)) \right\|_{\mathbb{F}} \leq \left(\frac{dN}{R} \sum_{|r| < R} \left(1 - \frac{|r|}{R} \right) \left\| \sum_{\substack{0 \leq n \leq B \\ 0 \leq n+r \leq B}} D(T(n+r)^2) D(T(n^2))^H \right\|_{\mathbb{F}} \right)^{1/2} + \frac{f}{2} R$$

$$T_{\lambda}(n) = g_{\varepsilon_0(n)} g_{\varepsilon_1(n)} \cdots g_{\varepsilon_{\lambda-1}(n)}$$

$$(n+r)^2 = (\varepsilon_{\ell-1} \varepsilon_{\ell-2} \cdots \varepsilon_{\lambda} \cdots)_q, \quad n^2 = (\varepsilon_{\ell-1} \varepsilon_{\ell-2} \cdots \varepsilon_{\lambda} \cdots)_q$$

$$D(T((n+r)^2)) D(T(n^2))^H$$

$$= D(T_{\lambda}((n+r)^2)) D(g_{\varepsilon_{\lambda}}) \cdots \overbrace{D(g_{\varepsilon_{\ell-1}}) D(g_{\varepsilon_{\ell-1}})^H}^{I_n} \cdots D(g_{\varepsilon_{\lambda}})^H D(T_{\lambda}(n^2))^H$$

$$= D(T_{\lambda}((n+r)^2)) D(T_{\lambda}(n^2))^H$$

$$T_\lambda(n) = g_{\varepsilon_0}(n)g_{\varepsilon_1}(n) \cdots g_{\varepsilon_{\mu-1}}(n)g_{\varepsilon_\mu}(n) \cdots g_{\varepsilon_{\lambda-1}}(n)$$

Fourier terms:

$$F_\lambda(h) = \frac{1}{q^\lambda} \sum_{0 \leq u < q^\lambda} e^{-2\pi i \frac{hu}{q^\lambda}} D(T_\lambda(u))$$

$$T_{\mu,\lambda}(n) = g_{\varepsilon_\mu(n)} \cdots g_{\varepsilon_{\lambda-1}(n)}$$

Fourier terms:

$$F_\lambda(h) = \frac{1}{q^\lambda} \sum_{0 \leq u < q^\lambda} e^{-2\pi i \frac{hu}{q^\lambda}} D(T_\lambda(u))$$

$$T_{\mu,\lambda}(n) = g_{\varepsilon_\mu(n)} \cdots g_{\varepsilon_{\lambda-1}(n)}$$

Fourier terms:

$$F_\lambda(h) = \frac{1}{q^\lambda} \sum_{0 \leq u < q^\lambda} e^{-2\pi i \frac{hu}{q^\lambda}} D(T_\lambda(u))$$

$$F_{\mu,\lambda}(h) := \frac{1}{q^\lambda} \sum_{0 \leq u < q^\lambda} e^{-2\pi i \frac{hu}{q^\lambda}} D(T_{\mu,\lambda}(u))$$

A subtle Fourier analysis of the double truncated sum (following the ideas of Mauduit and Rivat) leads to the following expression:

A subtle Fourier analysis of the double truncated sum (following the ideas of Mauduit and Rivat) leads to the following expression:

$$\frac{2}{\pi} \log \left(\frac{4e^{\pi/2} q^\lambda}{\pi} \right) q^{\lambda/2} \max_{0 \leq \ell < q^\lambda} \sum_{d|q^\lambda} d^{1/2} \sum_{\substack{0 \leq h_1, h_2, h_3, h_4 < q^\lambda \\ (h_1 + h_2 + h_3 + h_4, q^\lambda) = d \\ d|2r(h_1 + h_2) + 2sq^\mu(h_2 + h_3) + \ell}} \|F_{\mu, \lambda}(h_1)\|_{\mathbb{F}} \|F_{\mu, \lambda}(h_2)\|_{\mathbb{F}} \|F_{\mu, \lambda}(h_3)\|_{\mathbb{F}} \|F_{\mu, \lambda}(h_4)\|_{\mathbb{F}}$$

A subtle Fourier analysis of the double truncated sum (following the ideas of Mauduit and Rivat) leads to the following expression:

$$\frac{2}{\pi} \log \left(\frac{4e^{\pi/2} q^\lambda}{\pi} \right) q^{\lambda/2} \max_{0 \leq \ell < q^\lambda} \sum_{d|q^\lambda} d^{1/2} \sum_{\substack{0 \leq h_1, h_2, h_3, h_4 < q^\lambda \\ (h_1 + h_2 + h_3 + h_4, q^\lambda) = d \\ d|2r(h_1 + h_2) + 2sq^\mu(h_2 + h_3) + \ell}} \|F_{\mu, \lambda}(h_1)\|_{\mathbb{F}} \|F_{\mu, \lambda}(h_2)\|_{\mathbb{F}} \|F_{\mu, \lambda}(h_3)\|_{\mathbb{F}} \|F_{\mu, \lambda}(h_4)\|_{\mathbb{F}}$$

The analogue of this expression appears in Mauduit and Rivat's work and can be estimated as in their case.

★ Applications

★ La somme des chiffres des carrés:

Theorem (Mauduit and Rivat, 2009)

Let $q, r \geq 2$ and set $m = \gcd(q - 1, r)$. Then we have

- (i) $\lim_{x \rightarrow \infty} \frac{1}{x} \# \{n \leq x : s_q(n^2) \equiv a \pmod r\} = \frac{1}{r} Q(a, m),$
- (ii) $(\alpha s_q(n^2))_{n \in \mathbb{N}}$ is uniformly distributed modulo 1 iff α is irrational.

★ Applications

★ La somme des chiffres des carrés:

Theorem (Mauduit and Rivat, 2009)

Let $q, r \geq 2$ and set $m = \gcd(q - 1, r)$. Then we have

- (i) $\lim_{x \rightarrow \infty} \frac{1}{x} \# \{n \leq x : s_q(n^2) \equiv a \pmod r\} = \frac{1}{r} Q(a, m),$
- (ii) $(\alpha s_q(n^2))_{n \in \mathbb{N}}$ is uniformly distributed modulo 1 iff α is irrational.

Proof:

(i) ▶ $H = \mathbb{Z}/r\mathbb{Z}, \quad g_j = j \pmod r, 0 \leq j \leq q - 1.$

★ Applications

★ La somme des chiffres des carrés:

Theorem (Mauduit and Rivat, 2009)

Let $q, r \geq 2$ and set $m = \gcd(q - 1, r)$. Then we have

$$(i) \lim_{x \rightarrow \infty} \frac{1}{x} \# \{n \leq x : s_q(n^2) \equiv a \pmod{r}\} = \frac{1}{r} Q(a, m),$$

(ii) $(\alpha s_q(n^2))_{n \in \mathbb{N}}$ is uniformly distributed modulo 1 iff α is irrational.

Proof:

- (i) ▶ $H = \mathbb{Z}/r\mathbb{Z}$, $g_j = j \pmod{r}$, $0 \leq j \leq q - 1$.
- ▶ Then $G = H$, $T(n) = s_q(n) \pmod{r}$.

★ Applications

★ La somme des chiffres des carrés:

Theorem (Mauduit and Rivat, 2009)

Let $q, r \geq 2$ and set $m = \gcd(q - 1, r)$. Then we have

$$(i) \lim_{x \rightarrow \infty} \frac{1}{x} \# \{n \leq x : s_q(n^2) \equiv a \pmod{r}\} = \frac{1}{r} Q(a, m),$$

(ii) $(\alpha s_q(n^2))_{n \in \mathbb{N}}$ is uniformly distributed modulo 1 iff α is irrational.

Proof:

- (i)
 - ▶ $H = \mathbb{Z}/r\mathbb{Z}$, $g_j = j \pmod{r}$, $0 \leq j \leq q - 1$.
 - ▶ Then $G = H$, $T(n) = s_q(n) \pmod{r}$.
 - ▶ $\chi_k(u) = e^{2\pi i u k / r}$ $0 \leq k < r$... representations of G

★ Applications

★ La somme des chiffres des carrés:

Theorem (Mauduit and Rivat, 2009)

Let $q, r \geq 2$ and set $m = \gcd(q - 1, r)$. Then we have

$$(i) \lim_{x \rightarrow \infty} \frac{1}{x} \# \{n \leq x : s_q(n^2) \equiv a \pmod{r}\} = \frac{1}{r} Q(a, m),$$

(ii) $(\alpha s_q(n^2))_{n \in \mathbb{N}}$ is uniformly distributed modulo 1 iff α is irrational.

Proof:

- (i)
 - ▶ $H = \mathbb{Z}/r\mathbb{Z}$, $g_j = j \pmod{r}$, $0 \leq j \leq q - 1$.
 - ▶ Then $G = H$, $T(n) = s_q(n) \pmod{r}$.
 - ▶ $\chi_k(u) = e^{2\pi i u k / r}$ $0 \leq k < r$... representations of G
 - ▶ $m = \gcd(q - 1, r)$, $D_1(u) := \chi_{r/m}(u) = e^{2\pi i u / m}$.

★ Applications

★ La somme des chiffres des carrés:

Theorem (Mauduit and Rivat, 2009)

Let $q, r \geq 2$ and set $m = \gcd(q - 1, r)$. Then we have

$$(i) \lim_{x \rightarrow \infty} \frac{1}{x} \# \{n \leq x : s_q(n^2) \equiv a \pmod r\} = \frac{1}{r} Q(a, m),$$

(ii) $(\alpha s_q(n^2))_{n \in \mathbb{N}}$ is uniformly distributed modulo 1 iff α is irrational.

Proof:

- (i)
 - ▶ $H = \mathbb{Z}/r\mathbb{Z}$, $g_j = j \pmod r$, $0 \leq j \leq q - 1$.
 - ▶ Then $G = H$, $T(n) = s_q(n) \pmod r$.
 - ▶ $\chi_k(u) = e^{2\pi i u k / r}$ $0 \leq k < r$... representations of G
 - ▶ $m = \gcd(q - 1, r)$, $D_1(u) := \chi_{r/m}(u) = e^{2\pi i u / m}$. ■

★ Applications

★ La somme des chiffres des carrés:

Theorem (Mauduit and Rivat, 2009)

Let $q, r \geq 2$ and set $m = \gcd(q - 1, r)$. Then we have

- (i) $\lim_{x \rightarrow \infty} \frac{1}{x} \# \{n \leq x : s_q(n^2) \equiv a \pmod r\} = \frac{1}{r} Q(a, m)$,
- (ii) $(\alpha s_q(n^2))_{n \in \mathbb{N}}$ is uniformly distributed modulo 1 iff α is irrational.

Proof:

- (ii) ▶ $H = \mathbb{R}/\mathbb{Z}$, $g_j = \alpha j \pmod 1$, $0 \leq j \leq q - 1$.

★ Applications

★ La somme des chiffres des carrés:

Theorem (Mauduit and Rivat, 2009)

Let $q, r \geq 2$ and set $m = \gcd(q - 1, r)$. Then we have

- (i) $\lim_{x \rightarrow \infty} \frac{1}{x} \# \{n \leq x : s_q(n^2) \equiv a \pmod r\} = \frac{1}{r} Q(a, m),$
- (ii) $(\alpha s_q(n^2))_{n \in \mathbb{N}}$ is uniformly distributed modulo 1 iff α is irrational.

Proof:

- (ii)
 - ▶ $H = \mathbb{R}/\mathbb{Z}, \quad g_j = \alpha j \pmod 1, \quad 0 \leq j \leq q - 1.$
 - ▶ Then $G = H$ iff α irrational

★ Applications

★ La somme des chiffres des carrés:

Theorem (Mauduit and Rivat, 2009)

Let $q, r \geq 2$ and set $m = \gcd(q - 1, r)$. Then we have

- (i) $\lim_{x \rightarrow \infty} \frac{1}{x} \# \{n \leq x : s_q(n^2) \equiv a \pmod r\} = \frac{1}{r} Q(a, m)$,
- (ii) $(\alpha s_q(n^2))_{n \in \mathbb{N}}$ is uniformly distributed modulo 1 iff α is irrational.

Proof:

- (ii)
 - ▶ $H = \mathbb{R}/\mathbb{Z}$, $g_j = \alpha j \pmod 1$, $0 \leq j \leq q - 1$.
 - ▶ Then $G = H$ iff α irrational
 - ▶ $T(n) = \alpha s_q(n) \pmod 1$

★ Applications

★ La somme des chiffres des carrés:

Theorem (Mauduit and Rivat, 2009)

Let $q, r \geq 2$ and set $m = \gcd(q - 1, r)$. Then we have

$$(i) \lim_{x \rightarrow \infty} \frac{1}{x} \# \{n \leq x : s_q(n^2) \equiv a \pmod r\} = \frac{1}{r} Q(a, m),$$

(ii) $(\alpha s_q(n^2))_{n \in \mathbb{N}}$ is uniformly distributed modulo 1 iff α is irrational.

Proof:

- (ii)
 - ▶ $H = \mathbb{R}/\mathbb{Z}$, $g_j = \alpha j \pmod 1$, $0 \leq j \leq q - 1$.
 - ▶ Then $G = H$ iff α irrational
 - ▶ $T(n) = \alpha s_q(n) \pmod 1$




★ Applications

★ Automatic sequences:

Definition

A sequence $(u_n)_{n \geq 0}$ is called a q -automatic sequence, if u_n is the output of an automaton when the input is the q -ary expansion of n .




★ Applications

★ Automatic sequences:

Definition

A sequence $(u_n)_{n \geq 0}$ is called a q -automatic sequence, if u_n is the output of an automaton when the input is the q -ary expansion of n .




$$32 = (1012)_3$$

★ Applications

★ Automatic sequences:

Definition

A sequence $(u_n)_{n \geq 0}$ is called a q -automatic sequence, if u_n is the output of an automaton when the input is the q -ary expansion of n .




$$32 = (1012)_3$$

★ Applications

★ Automatic sequences:

Definition

A sequence $(u_n)_{n \geq 0}$ is called a q -automatic sequence, if u_n is the output of an automaton when the input is the q -ary expansion of n .




$$32 = (1012)_3$$

★ Applications

★ Automatic sequences:

Definition

A sequence $(u_n)_{n \geq 0}$ is called a q -automatic sequence, if u_n is the output of an automaton when the input is the q -ary expansion of n .




$$32 = (1012)_3$$

★ Applications

★ Automatic sequences:

Definition

A sequence $(u_n)_{n \geq 0}$ is called a q -automatic sequence, if u_n is the output of an automaton when the input is the q -ary expansion of n .




$$32 = (1012)_3 \quad u_{32} = a,$$

★ Applications

★ Automatic sequences:

Definition

A sequence $(u_n)_{n \geq 0}$ is called a q -automatic sequence, if u_n is the output of an automaton when the input is the q -ary expansion of n .



$$32 = (1012)_3 \quad u_{32} = a,$$


$$61 = (2021)_3$$

★ Applications

★ Automatic sequences:

Definition

A sequence $(u_n)_{n \geq 0}$ is called a q -automatic sequence, if u_n is the output of an automaton when the input is the q -ary expansion of n .



$$32 = (1012)_3 \quad u_{32} = a,$$


$$61 = (2021)_3$$

★ Applications

★ Automatic sequences:

Definition

A sequence $(u_n)_{n \geq 0}$ is called a q -automatic sequence, if u_n is the output of an automaton when the input is the q -ary expansion of n .



$$32 = (1012)_3 \quad u_{32} = a,$$


$$61 = (2021)_3$$

★ Applications

★ Automatic sequences:

Definition

A sequence $(u_n)_{n \geq 0}$ is called a q -automatic sequence, if u_n is the output of an automaton when the input is the q -ary expansion of n .



$$32 = (1012)_3 \quad u_{32} = a,$$


$$61 = (2021)_3$$

★ Applications

★ Automatic sequences:

Definition

A sequence $(u_n)_{n \geq 0}$ is called a q -automatic sequence, if u_n is the output of an automaton when the input is the q -ary expansion of n .



$$32 = (1012)_3 \quad u_{32} = a,$$


$$61 = (2021)_3 \quad u_{61} = b$$

★ Applications

★ Automatic sequences:

Definition


A sequence $(u_n)_{n \geq 0}$ is called a q -automatic sequence, if u_n is the output of an automaton when the input is the q -ary expansion of n .




$$32 = (1012)_3 \quad u_{32} = a,$$


$$61 = (2021)_3 \quad u_{61} = b$$

$(u_n)_{n \geq 0} : aaaaabaabaabaabaabbbaaabaabbbaaabaabbbaaaaaaba \dots$






$$M_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$



$$M_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$


$$M_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$



$$M_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_2 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$




$$M_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_2 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

$$32 = (1012)_3 :$$

$$\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$




$$M_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_2 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

$$32 = (1012)_3 :$$

$$M_1 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$




$$M_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_2 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

$$32 = (1012)_3 :$$

$$M_0 \circ M_1 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$




$$M_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_2 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

$$32 = (1012)_3 :$$

$$M_1 \circ M_0 \circ M_1 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$




$$M_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_2 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

$$32 = (1012)_3 : \quad M_2 \circ M_1 \circ M_0 \circ M_1 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$




$$M_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_2 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

$$S(n) := M_{\varepsilon_0(n)} M_{\varepsilon_1(n)} \cdots M_{\varepsilon_{\ell-1}(n)}$$

$$u_n = f(S(n) \mathbf{e}_1), \quad \mathbf{e}_j^T S(n) \mathbf{e}_1 = \begin{cases} 1, & \text{output state } s_j \\ 0, & \text{otherwise.} \end{cases}$$




$$M_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M_2 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$


Definition

A q -automatic sequence is called *invertible* if there exists an automaton such that all transition matrices are invertible and M_0 is the identity matrix.




$(u_n)_{n \geq 0} : aaaaabaabaabaabbbaaabaabbbaaabaabbbaaaaaaba\dots$

Frequency of a in $(u_n)_{n \geq 0} : \lim_{N \rightarrow \infty} \frac{1}{N} \#\{0 \leq n < N : u_n = a\}$




$(u_n)_{n \geq 0} : \text{aaaaabaabaabaabbaaabaabbaaabaabbaaaaaaba} \dots$

Frequency of a in $(u_{3n})_{n \geq 0} : \lim_{N \rightarrow \infty} \frac{1}{N} \#\{0 \leq n < N : u_{3n} = a\}$



$(u_n)_{n \geq 0} : \text{aaabaabaabaabbbaabaabbbaabaabbbaaaaaaba} \dots$


Frequency of a in $(u_{n^2})_{n \geq 0} : \lim_{N \rightarrow \infty} \frac{1}{N} \#\{0 \leq n < N : u_{n^2} = a\} = ?$



$(u_n)_{n \geq 0} : \text{aaabaabaabaabbbaabaaabbbaabaaabbbaaaaaaba} \dots$

Frequency of a in $(u_{n^2})_{n \geq 0} : \lim_{N \rightarrow \infty} \frac{1}{N} \#\{0 \leq n < N : u_{n^2} = a\} = ?$

$H \dots$ group of permutation matrices, $g_i = M_i$,




$(u_n)_{n \geq 0} : \text{aaabaabaabaabbbaabaaabbbaabaaabbbaaaaaaba} \dots$

Frequency of a in $(u_{n^2})_{n \geq 0} : \lim_{N \rightarrow \infty} \frac{1}{N} \#\{0 \leq n < N : u_{n^2} = a\} = ?$

$H \dots$ group of permutation matrices, $g_i = M_i$,

$$T(n) = S(n),$$




$(u_n)_{n \geq 0} : \text{aaabaabaabaabbbaabaaabbbaabaaabbbaaaaaaba} \dots$

Frequency of a in $(u_{n^2})_{n \geq 0} : \quad \lim_{N \rightarrow \infty} \frac{1}{N} \#\{0 \leq n < N : u_{n^2} = a\} = ?$

$H \dots$ group of permutation matrices, $g_i = M_i$,

$$T(n) = S(n), \quad \frac{1}{N} \#\{0 \leq n < N : u_{n^2} = s_j\} = \frac{1}{N} \sum_{n=0}^{N-1} \mathbf{e}_j^T T(n^2) \mathbf{e}_1$$



$(u_n)_{n \geq 0} : \text{aaabaabaabaabbbaabaaabbbaabaaabbbaaaaaaba} \dots$

Frequency of a in $(u_{n^2})_{n \geq 0}$: $\lim_{N \rightarrow \infty} \frac{1}{N} \#\{0 \leq n < N : u_{n^2} = a\} = ?$

$H \dots$ group of permutation matrices, $g_i = M_i$,

$$T(n) = S(n), \quad \frac{1}{N} \#\{0 \leq n < N : u_{n^2} = s_j\} = \frac{1}{N} \sum_{n=0}^{N-1} \mathbf{e}_j^T T(n^2) \mathbf{e}_1$$

Theorem (Drmota and Morgenbesser, 2010)


Let $q \geq 2$ and $(u_n)_{n \geq 0}$ be an invertible q -automatic sequence. Then the frequency of each letter of the subsequence $(u_{n^2})_{n \geq 0}$ exists.

Thank you!

- 1 LAUWERENS KUIPERS AND HARALD NIEDERREITER: *Uniform Distribution of Sequences*. Wiley-Interscience Publication, 1974
- 2 JEAN-PAUL ALLOUCHE AND JEFFREY SHALLIT: *Automatic sequences*. Cambridge University Press, 2003
- 3 ALEKSANDR O. GELFOND: *Sur les nombres qui ont des propriétés additives et multiplicatives données*. Acta Arithmetica, 1968
- 4 CHRISTIAN MAUDUIT AND JOËL RIVAT: *La somme des chiffres des carrés*. Acta Mathematica, 2009
- 5 MICHAEL DRMOTA AND JOHANNES F. MORGENBESSER: *Generalized Thue-Morse Sequence of Squares*. submitted

Examples of automatic sequences

Thue-Morse sequence:



Rudin-Shapiro sequence:

