

# **A cartesian bicategory of polynomial functors in homotopy type theory**

---

Samuel Mimram (joint with Eric Finster, Maxime Lucas and Thomas Seiller)

Journées SCALP – November 4, 2021

# In a nutshell

The situation:

- the category of polynomial functors is cartesian closed

Our contributions:

The situation:

- the category of polynomial functors is cartesian closed
- the 2-category of polynomial functors is not cartesian closed [Girard]

Our contributions:

The situation:

- the category of polynomial functors is cartesian closed
- the 2-category of polynomial functors is not cartesian closed [Girard]
- we should consider polynomial functors in groupoids [Kock]

Our contributions:

The situation:

- the category of polynomial functors is cartesian closed
- the 2-category of polynomial functors is not cartesian closed [Girard]
- we should consider polynomial functors in groupoids [Kock]

Our contributions:

- we have formalized polynomials in groupoids (or spaces) in HoTT/Agda

The situation:

- the category of polynomial functors is cartesian closed
- the 2-category of polynomial functors is not cartesian closed [Girard]
- we should consider polynomial functors in groupoids [Kock]

Our contributions:

- we have formalized polynomials in groupoids (or spaces) in HoTT/Agda
- we have shown that the resulting bicategory is cartesian closed

The situation:

- the category of polynomial functors is cartesian closed
- the 2-category of polynomial functors is not cartesian closed [Girard]
- we should consider polynomial functors in groupoids [Kock]

Our contributions:

- we have formalized polynomials in groupoids (or spaces) in HoTT/Agda
- we have shown that the resulting bicategory is cartesian closed
- we have provided a small axiomatization of the type  $\mathbb{B}$  of natural numbers and bijections

Part I

# Polynomial functors



## Categorifying polynomials

A **polynomial** is a sum of monomials

$$P(X) = \sum_{0 \leq i < k} X^{n_i}$$

(no coefficients, but repetitions allowed)

## Categorifying polynomials

A **polynomial** is a sum of monomials

$$P(X) = \sum_{0 \leq i < k} X^{n_i}$$

(no coefficients, but repetitions allowed)

We can **categorify** this notion: replace natural numbers by elements of a set.

$$P(X) = \sum_{b \in B} X^{E_b}$$


# Polynomial functors

This data can be encoded as a **polynomial**  $P$ , which is a diagram in **Set**:

$$E \xrightarrow{P} B$$

where

- $b \in B$  is a monomial
- $E_b = p^{-1}(b)$  is the set of instances of  $X$  in the monomial  $b$ .



# Polynomial functors

This data can be encoded as a **polynomial**  $P$ , which is a diagram in **Set**:

$$E \xrightarrow{P} B$$

where

- $b \in B$  is a monomial
- $E_b = p^{-1}(b)$  is the set of instances of  $X$  in the monomial  $b$ .


It induces a **polynomial functor**

$$[[P]] : \mathbf{Set} \rightarrow \mathbf{Set}$$

$$X \mapsto \sum_{b \in B} X^{E_b}$$

## Polynomial functors

For instance, consider the polynomial corresponding to the function




The associated polynomial functor is

$$[[P]](X) : \mathbf{Set} \rightarrow \mathbf{Set}$$

$$X \mapsto X \times X \sqcup X \times X \times X$$

## Polynomial functors

For instance, consider the polynomial corresponding to the function

$$\mathbb{N} \xrightarrow{p} 1$$


The associated polynomial functor is

$$\begin{aligned} \llbracket P \rrbracket(X) &: \mathbf{Set} \rightarrow \mathbf{Set} \\ X &\mapsto X \times X \times X \times \dots \end{aligned}$$

## Polynomial functors

For instance, consider the polynomial corresponding to the function

$$\mathbb{N} \xrightarrow{p} 1$$

⋮

●

●

●



The associated polynomial functor is

$$[[P]](X) : \mathbf{Set} \rightarrow \mathbf{Set}$$

$$X \mapsto X \times X \times X \times \dots$$

A polynomial is **finitary** when each monomial is a finite product.


## Polynomial functors: typed variant

We will more generally consider a “colored variant” of polynomials  $P$

$$I \xleftarrow{s} E \xrightarrow{P} B \xrightarrow{t} J$$

this means that

- each monomial  $b$  has a color  $t(b) \in J$ ,
- each occurrence of a variable  $e \in E$  has a color  $s(e) \in I$ .





## Polynomial functors: typed variant

We will more generally consider a “colored variant” of polynomials  $P$

$$I \xleftarrow{s} E \xrightarrow{p} B \xrightarrow{t} J$$

this means that

- each monomial  $b$  has a color  $t(b) \in J$ ,
- each occurrence of a variable  $e \in E$  has a color  $s(e) \in I$ .

It induces a **polynomial functor**

$$\begin{aligned} \llbracket P \rrbracket(X) &: \mathbf{Set}^I \rightarrow \mathbf{Set}^J \\ (X_i)_{i \in I} &\mapsto \left( \sum_{b \in t^{-1}(j)} \prod_{e \in p^{-1}(b)} X_{s(e)} \right)_{j \in J} \end{aligned}$$



## Polynomial vs polynomial functors

A *polynomial*  $P$

$$I \xleftarrow{s} E \xrightarrow{p} B \xrightarrow{t} J$$

induces a *polynomial functor*

$$[[P]] : \mathbf{Set}^I \rightarrow \mathbf{Set}^J$$

We have mentioned that composition is defined for polynomials. However, on polynomials, it is not strictly associative: we can build a *bicategory* **Poly** of sets and polynomial functors.


This suggests that 2-cells are an important part of the story!

# Morphisms between polynomials

A morphism between two polynomials is

$$\begin{array}{ccccccc}
 I & \xleftarrow{s} & E & \xrightarrow{p} & B & \xrightarrow{t} & J \\
 \parallel & & \varepsilon \downarrow & \lrcorner & \downarrow \beta & & \parallel \\
 I & \xleftarrow{s'} & E' & \xrightarrow{p'} & B' & \xrightarrow{t'} & J
 \end{array}$$

We send monomials to monomials, preserving typing and arities:



# Morphisms between polynomials

A morphism between two polynomials is

$$\begin{array}{ccccccc}
 I & \xleftarrow{s} & E & \xrightarrow{p} & B & \xrightarrow{t} & J \\
 \parallel & & \varepsilon \downarrow & \lrcorner & \downarrow \beta & & \parallel \\
 I & \xleftarrow{s'} & E' & \xrightarrow{p'} & B' & \xrightarrow{t'} & J
 \end{array}$$

We send monomials to monomials, preserving typing and arities:

$$\begin{array}{ccc}
 \begin{array}{c}
 i_1 \quad i_2 \quad \dots \quad i_{n-1} i_n \\
 | \quad | \quad \dots \quad | \quad | \\
 \hline
 b \\
 | \\
 j
 \end{array}
 & \mapsto &
 \begin{array}{c}
 i_1 \quad i_2 \quad \dots \quad i_{n-1} i_n \\
 | \quad | \quad \dots \quad | \quad | \\
 \hline
 \beta(b) \\
 | \\
 j
 \end{array}
 \end{array}$$

We can build a bicategory **Poly** of sets, polynomials and morphisms of polynomials.

# Morphisms between polynomials

A morphism between two polynomials is

$$\begin{array}{ccccccc}
 I & \xleftarrow{s} & E & \xrightarrow{p} & B & \xrightarrow{t} & J \\
 \parallel & & \varepsilon \downarrow & \lrcorner & \downarrow \beta & & \parallel \\
 I & \xleftarrow{s'} & E' & \xrightarrow{p'} & B' & \xrightarrow{t'} & J
 \end{array}$$

We send monomials to monomials, preserving typing and arities:



We can build a bicategory **Poly** of sets, polynomials and morphisms of polynomials.

In the following, we will restrict to the case where 2-cells are *equivalences*.

## Morphisms between polynomial functors

A morphism between polynomial functors

$$[[P]], [[Q]] : \mathbf{Set}^I \rightarrow \mathbf{Set}^J$$

is a “suitable” natural transformation, and we can build a 2-category **PolyFun**.

## Cartesian structure

The category **PolyFun** is cartesian. Namely, given two polynomial functors in **Poly**

$$P : I \rightarrow J \qquad Q : I \rightarrow K$$

i.e., in **Cat**,

$$\llbracket P \rrbracket : \mathbf{Set}^I \rightarrow \mathbf{Set}^J \qquad \llbracket Q \rrbracket : \mathbf{Set}^I \rightarrow \mathbf{Set}^K$$

we have, in **Cat**,

$$\langle P, Q \rangle : \mathbf{Set}^I \rightarrow \mathbf{Set}^J \times \mathbf{Set}^K \cong \mathbf{Set}^{J \sqcup K}$$

and the constructions preserve polynomiality: in **PolyFun**,

$$\langle P, Q \rangle : I \rightarrow (J \sqcup K)$$



## Closed structure

For the closed structure, we can hope for the same: given, in **PolyFun**,

$$P : I \sqcup J \rightarrow K$$

i.e., in **Cat**,

$$\llbracket P \rrbracket : \mathbf{Set}^{I \sqcup J} \rightarrow \mathbf{Set}^K$$

we have

$$\underline{\mathbf{Set}^{I \sqcup J} \rightarrow \mathbf{Set}^K}$$

## Closed structure

For the closed structure, we can hope for the same: given, in **PolyFun**,

$$P : I \sqcup J \rightarrow K$$

i.e., in **Cat**,

$$\llbracket P \rrbracket : \mathbf{Set}^{I \sqcup J} \rightarrow \mathbf{Set}^K$$

we have

$$\frac{\mathbf{Set}^{I \sqcup J} \rightarrow \mathbf{Set}^K}{\mathbf{Set}^I \times \mathbf{Set}^J \rightarrow \mathbf{Set}^K}$$

## Closed structure

For the closed structure, we can hope for the same: given, in **PolyFun**,

$$P : I \sqcup J \rightarrow K$$

i.e., in **Cat**,

$$\llbracket P \rrbracket : \mathbf{Set}^{I \sqcup J} \rightarrow \mathbf{Set}^K$$

we have

$$\frac{\frac{\mathbf{Set}^{I \sqcup J} \rightarrow \mathbf{Set}^K}{\mathbf{Set}^I \times \mathbf{Set}^J \rightarrow \mathbf{Set}^K}}{\mathbf{Set}^I \rightarrow (\mathbf{Set}^K)^{\mathbf{Set}^J}}$$

## Closed structure

For the closed structure, we can hope for the same: given, in **PolyFun**,

$$P : I \sqcup J \rightarrow K$$

i.e., in **Cat**,

$$\llbracket P \rrbracket : \mathbf{Set}^{I \sqcup J} \rightarrow \mathbf{Set}^K$$

we have

$$\frac{\frac{\frac{\mathbf{Set}^{I \sqcup J} \rightarrow \mathbf{Set}^K}{\mathbf{Set}^I \times \mathbf{Set}^J \rightarrow \mathbf{Set}^K}}{\mathbf{Set}^I \rightarrow (\mathbf{Set}^K)^{\mathbf{Set}^J}}}{\mathbf{Set}^I \rightarrow \mathbf{Set}^{\mathbf{Set}^J \times K}}$$

## Closed structure

For the closed structure, we can hope for the same: given, in **PolyFun**,

$$P : I \sqcup J \rightarrow K$$

i.e., in **Cat**,

$$\llbracket P \rrbracket : \mathbf{Set}^{I \sqcup J} \rightarrow \mathbf{Set}^K$$

we have

$$\frac{\frac{\mathbf{Set}^{I \sqcup J} \rightarrow \mathbf{Set}^K}{\mathbf{Set}^I \times \mathbf{Set}^J \rightarrow \mathbf{Set}^K}}{\mathbf{Set}^I \rightarrow (\mathbf{Set}^K)^{\mathbf{Set}^J}} \quad \frac{\mathbf{Set}^I \rightarrow (\mathbf{Set}^K)^{\mathbf{Set}^J}}{\mathbf{Set}^I \rightarrow \mathbf{Set}^{\mathbf{Set}^J \times K}}$$

which suggests defining the closure as

$$[J, K] = \mathbf{Set}^J \times K$$

## Closed structure

For the closed structure, we can hope for the same: given, in **PolyFun**,

$$P : I \sqcup J \rightarrow K$$

i.e., in **Cat**,

$$\llbracket P \rrbracket : \mathbf{Set}^{I \sqcup J} \rightarrow \mathbf{Set}^K$$

we have

$$\frac{\frac{\mathbf{Set}^{I \sqcup J} \rightarrow \mathbf{Set}^K}{\mathbf{Set}^I \times \mathbf{Set}^J \rightarrow \mathbf{Set}^K}}{\mathbf{Set}^I \rightarrow (\mathbf{Set}^K)^{\mathbf{Set}^J}}}{\mathbf{Set}^I \rightarrow \mathbf{Set}^{\mathbf{Set}^J \times K}}$$

which suggests defining the closure as

$$[J, K] = \mathbf{Set}^J \times K$$


for LL-people: this looks like  $!J \wp K$ .

## Closed structure

In terms of operations, the intuition behind the bijection

$$\mathbf{PolyFun}(I \sqcup J, K) \cong \mathbf{PolyFun}(I, \mathbf{Set}^J \times K)$$

is that we can formally transform operations as follows




## Closed structure

In terms of operations, the intuition behind the bijection

$$\mathbf{PolyFun}(I \sqcup J, K) \cong \mathbf{PolyFun}(I, \mathbf{Set}/J \times K)$$

is that we can formally transform operations as follows



via

$$\mathbf{Set}^J \simeq \mathbf{Set}/J$$



## Closed structure

There are two problems with our closure. The first one is that

$$[I, J] = \mathbf{Set}/I \times J$$

is too large to be an object of our category.

## Closed structure

There are two problems with our closure. The first one is that

$$[I, J] = \mathbf{Set}/I \times J$$

is too large to be an object of our category.

One can restrict to polynomial functors which are **finitary**: we can then take

$$[I, J] = \mathbf{Set}_{\text{fin}}/I \times J$$

## Closed structure

There are two problems with our closure. The first one is that

$$[I, J] = \mathbf{Set}/I \times J$$

is too large to be an object of our category.

One can restrict to polynomial functors which are **finitary**: we can then take

$$[I, J] = \mathbf{Set}_{\text{fin}}/I \times J$$

or rather

$$[I, J] = \mathbb{N}/I \times J$$

## Closed structure

There are two problems with our closure. The first one is that

$$[I, J] = \mathbf{Set}/I \times J$$

is too large to be an object of our category.

One can restrict to polynomial functors which are **finitary**: we can then take

$$[I, J] = \mathbf{Set}_{\text{fin}}/I \times J$$

or rather

$$[I, J] = \mathbb{N}/I \times J$$

Finitary polynomial functors are also known as **normal functors** [Girard].

### **Theorem**

*The category **PolyFun** is cartesian closed.*

## Theorem

*The category **PolyFun** is cartesian closed.*

## Remark (Girard)

The 2-category **PolyFun** is not cartesian closed.

## Failure of the cartesian closed structure

We would like to have an equivalence of categories

$$\mathbf{PolyFun}(I \sqcup J, K) \simeq \mathbf{PolyFun}(I, \mathbb{N}/J \times K)$$

## Failure of the cartesian closed structure

We would like to have an equivalence of categories

$$\mathbf{PolyFun}(I \sqcup J, K) \simeq \mathbf{PolyFun}(I, \mathbb{N}/J \times K)$$

but consider the polynomial functor

$$\llbracket P \rrbracket(X) = X^2 \quad : \quad \mathbf{Set}^{0 \sqcup 1} \rightarrow \mathbf{Set}^1$$



## Failure of the cartesian closed structure

We would like to have an equivalence of categories

$$\mathbf{PolyFun}(I \sqcup J, K) \simeq \mathbf{PolyFun}(I, \mathbb{N}/J \times K)$$

but consider the polynomial functor

$$\llbracket P \rrbracket(X) = X^2 \quad : \quad \mathbf{Set}^{0 \sqcup 1} \rightarrow \mathbf{Set}^1$$

which is induced by the polynomial

$$1 \longleftarrow 2 \longrightarrow 1 \longrightarrow 1$$

## Failure of the cartesian closed structure

We would like to have an equivalence of categories

$$\mathbf{PolyFun}(I \sqcup J, K) \simeq \mathbf{PolyFun}(I, \mathbb{N}/J \times K)$$

but consider the polynomial functor

$$\llbracket P \rrbracket(X) = X^2 \quad : \quad \mathbf{Set}^{0 \sqcup 1} \rightarrow \mathbf{Set}^1$$

which has two automorphisms

$$\begin{array}{ccccccc} 1 & \longleftarrow & 2 & \longrightarrow & 1 & \longrightarrow & 1 \\ \parallel & & \tau \downarrow \text{id}^{\perp} & & \downarrow & & \parallel \\ 1 & \longleftarrow & 2 & \longrightarrow & 1 & \longrightarrow & 1 \end{array}$$

## Failure of the cartesian closed structure

We would like to have an equivalence of categories

$$\mathbf{PolyFun}(I \sqcup J, K) \simeq \mathbf{PolyFun}(I, \mathbb{N}/J \times K)$$

but consider the polynomial functor

$$\llbracket P \rrbracket(X) = X^2 \quad : \quad \mathbf{Set}^{0 \sqcup 1} \rightarrow \mathbf{Set}^1$$

whose exponential transpose is

$$0 \longleftarrow 0 \longrightarrow 1 \xrightarrow{* \mapsto 2} \mathbb{N}$$

and has only one automorphism.

## Failure of the cartesian closed structure

We would like to have an equivalence of categories

$$\mathbf{PolyFun}(I \sqcup J, K) \simeq \mathbf{PolyFun}(I, \mathbb{N}/J \times K)$$

but consider the polynomial functor

$$\llbracket P \rrbracket(X) = X^2 \quad : \quad \mathbf{Set}^{0 \sqcup 1} \rightarrow \mathbf{Set}^1$$

whose exponential transpose is

$$0 \longleftarrow 0 \longrightarrow 1 \xrightarrow{* \mapsto 2} \mathbb{N}$$

and has only one automorphism.

The equivalence fails:

$$\mathbf{PolyFun}(0 \sqcup 1, 1) \not\simeq \mathbf{PolyFun}(0, \mathbb{N}/1 \times 1)$$

(two elements on the left, one on the right)

## Fixing the cartesian closed structure

The failure of the equivalence

$$\mathbf{PolyFun}(0 \sqcup 1, 1) \not\cong \mathbf{PolyFun}(0, \mathbb{N}/1 \times 1)$$

can be interpreted as being due to the fact that  $2 \in \mathbb{N}/1$  has no non-trivial isomorphism.

This suggests moving to **groupoids!**

## Fixing the cartesian closed structure

The failure of the equivalence

$$\mathbf{PolyFun}(0 \sqcup 1, 1) \not\cong \mathbf{PolyFun}(0, \mathbb{N}/1 \times 1)$$

can be interpreted as being due to the fact that  $2 \in \mathbb{N}/1$  has no non-trivial isomorphism.

This suggests moving to **groupoids!**

More precisely, we should replace  $\mathbb{N}$  by the groupoid  $\mathbb{B}$  of all symmetric groups.

## Polynomial functors in groupoids

The notion of polynomial functor generalizes in any locally cartesian closed category.

## Polynomial functors in groupoids

The notion of polynomial functor generalizes in any locally cartesian closed category.

...but the category **Gpd** is not cartesian closed!



## Polynomial functors in groupoids

The notion of polynomial functor generalizes in any locally cartesian closed category.

...but the category **Gpd** is not cartesian closed!

Kock has identified that if we perform all the usual constructions up to homotopy (slice, pullbacks, etc.), we recover a suitable setting to define polynomial functors.

## Polynomial functors in groupoids

The notion of polynomial functor generalizes in any locally cartesian closed category.

...but the category **Gpd** is not cartesian closed!

Kock has identified that if we perform all the usual constructions up to homotopy (slice, pullbacks, etc.), we recover a suitable setting to define polynomial functors.

This requires properly defining and using all the usual constructions in a suitable 2-categorical sense.

## Polynomial functors in groupoids

Given a polynomial  $P$

$$E \xrightarrow{P} B$$

the induced polynomial functor

$$[[P]] : \mathbf{Gpd} \rightarrow \mathbf{Gpd}$$

$$X \mapsto \int^{b \in B} E_b$$

where  $E_b$  is the *homotopy fiber* of  $p$  at  $b$  and

$$\int^{b \in E} E_b = \sum_{b \in \pi_0(B)} X_b / \text{Aut}(b)$$

where the quotient is to be taken 2-categorically / homotopically...

Part II

# Formalization in Agda

There is a framework in which everything is constructed *up to homotopy* for free:  
**homotopy type theory**.

In particular, there is a well-known notion of groupoid in this setting:  
a type with no non-trivial equalities between equalities.

Let's formally develop the theory of polynomials in this setting.

## Formalizing polynomials

A polynomial

$$I \xleftarrow{s} E \xrightarrow{p} B \xrightarrow{t} J$$

is a **container**:

```
record Poly (I J : Type) : Type1 where
  field
    Op : J → Type
    Pm : (i : I) → {j : J} → Op j → Type
```

We sometimes write

$$I \rightsquigarrow J = \text{Poly } I \ J$$

## Composing polynomials

The polynomial functor induced by a polynomial  $P$  is

$$\llbracket \_ \rrbracket : I \rightsquigarrow J \rightarrow (I \rightarrow \text{Type}) \rightarrow (J \rightarrow \text{Type})$$
$$\llbracket \_ \rrbracket P X j = \Sigma (Op P j) (\lambda c \rightarrow (i : I) \rightarrow (p : Pm P i c) \rightarrow (X i))$$

## Composing polynomials

The polynomial functor induced by a polynomial  $P$  is

$$\llbracket \_ \rrbracket : I \rightsquigarrow J \rightarrow (I \rightarrow \text{Type}) \rightarrow (J \rightarrow \text{Type})$$
$$\llbracket \_ \rrbracket P X j = \Sigma (0p P j) (\lambda c \rightarrow (i : I) \rightarrow (p : Pm P i c) \rightarrow (X i))$$

The composite of two polynomials is

$$\_ \cdot \_ : I \rightsquigarrow J \rightarrow J \rightsquigarrow K \rightarrow I \rightsquigarrow K$$
$$0p (P \cdot Q) = \llbracket Q \rrbracket (0p P)$$
$$Pm (\_ \cdot \_ P Q) i (c, a) = \Sigma J (\lambda j \rightarrow \Sigma (Pm Q j c) (\lambda p \rightarrow Pm P i (a j p)))$$



# A bicategory

## **Theorem**

*We can build a pre-bicategory of types, polynomials and their morphisms.*

Note: by univalence, we can use propositional equality for 2-cells, which simplifies the definition.

## A bicategory

### **Theorem**

*We can build a pre-bicategory of types, polynomials and their morphisms.*

### **Theorem**

*We can build a bicategory of groupoids, polynomials in groupoids and their morphisms.*

## A bicategory

### **Theorem**

*We can build a pre-bicategory of types, polynomials and their morphisms.*

### **Theorem**

*We can build a bicategory of groupoids, polynomials in groupoids and their morphisms.*

### **Theorem**

*This bicategory is cartesian with  $\sqcup$  as coproduct.*

## Defining the exponential

In order to define the 1-categorical closure, the plan was:

$$\mathbf{Set} \rightsquigarrow \mathbf{Set}_{\text{fin}} \rightsquigarrow \mathbb{N}$$

## Defining the exponential

In order to define the 1-categorical closure, the plan was:

$$\mathbf{Set} \rightsquigarrow \mathbf{Set}_{\text{fin}} \rightsquigarrow \mathbb{N}$$

For the 2-categorical closure the plan is

$$\mathbf{Gpd} \rightsquigarrow \mathbf{Gpd}_{\text{fin}} \rightsquigarrow \mathbb{B}$$

Here,  $\mathbb{B}$  is the groupoid with  $n \in \mathbb{N}$  as objects and  $\Sigma_n$  as automorphisms on  $n$ .

## Finite types

We write `Fin n` for the canonical finite type with `n` elements:  
its constructors are `0` to `n-1`.

We write `Fin n` for the canonical finite type with `n` elements:  
its constructors are 0 to `n-1`.

```
data Fin : ℕ → Set where
  zero : {n : ℕ}           → Fin (suc n)
  suc  : {n : ℕ} (i : Fin n) → Fin (suc n)
```

## Finite types

The predicate of being **finite** is

```
is-finite : Type → Type
```

```
is-finite A =  $\Sigma \mathbb{N} (\lambda n \rightarrow \| A \simeq \text{Fin } n \|)$ 
```



## Finite types

The predicate of being **finite** is

```
is-finite : Type → Type
```

```
is-finite A =  $\Sigma \mathbb{N} (\lambda n \rightarrow \| A \simeq \text{Fin } n \|)$ 
```

The type of finite types is

```
FinType : Type1
```

```
FinType =  $\Sigma \text{Type is-finite}$ 
```

## Finite types

The predicate of being **finite** is

`is-finite` : `Type` → `Type`

`is-finite` A =  $\Sigma \mathbb{N} (\lambda n \rightarrow \parallel A \simeq \text{Fin } n \parallel)$

The type of finite types is

`FinType` : `Type`<sub>1</sub>

`FinType` =  $\Sigma \text{Type } \text{is-finite}$

(note that this is a *large* type)

## Finitary polynomials

A polynomial is **finitary** when, for each operation, the total space of its parameters is finite:

```
is-finitary : (P : I  $\rightsquigarrow$  J)  $\rightarrow$  Type
```

```
is-finitary P = {j : J} (c : Op P j)  $\rightarrow$  is-finite ( $\Sigma$  I ( $\lambda$  i  $\rightarrow$  Pm P i c))
```

## A small model for finite types

The type of **integers** is

```
data  $\mathbb{N}$  : Type where  
  zero :  $\mathbb{N}$   
  suc  :  $\mathbb{N} \rightarrow \mathbb{N}$ 
```

## A small model for finite types

The type  $\mathbb{B}$  is

```
data  $\mathbb{B}$  : Type where
  obj      :  $\mathbb{N} \rightarrow \mathbb{B}$ 
  hom      : {n :  $\mathbb{N}$ } ( $\alpha$  : Fin n  $\simeq$  Fin n)  $\rightarrow$  obj n  $\equiv$  obj n
  id-coh   : (n :  $\mathbb{N}$ )  $\rightarrow$  hom {n = n}  $\simeq$ -refl  $\equiv$  refl
  comp-coh : {m n o :  $\mathbb{N}$ } ( $\alpha$  : Fin m  $\simeq$  Fin n) ( $\beta$  : Fin n  $\simeq$  Fin o)  $\rightarrow$ 
             hom ( $\simeq$ -trans  $\alpha$   $\beta$ )  $\equiv$  hom  $\alpha$   $\cdot$  hom  $\beta$ 
```

(this is a small higher inductive type!)

## A small model for finite types

The type  $\mathbb{B}$  is

```
data  $\mathbb{B}$  : Type where
  obj      :  $\mathbb{N} \rightarrow \mathbb{B}$ 
  hom      :  $\{n : \mathbb{N}\} (\alpha : \text{Fin } n \simeq \text{Fin } n) \rightarrow \text{obj } n \equiv \text{obj } n$ 
  id-coh   :  $(n : \mathbb{N}) \rightarrow \text{hom } \{n = n\} \simeq\text{-refl} \equiv \text{refl}$ 
  comp-coh :  $\{m \ n \ o : \mathbb{N}\} (\alpha : \text{Fin } m \simeq \text{Fin } n) (\beta : \text{Fin } n \simeq \text{Fin } o) \rightarrow$ 
              $\text{hom } (\simeq\text{-trans } \alpha \ \beta) \equiv \text{hom } \alpha \cdot \text{hom } \beta$ 
```

(this is a small higher inductive type!)

### Theorem

$\text{FinType} \simeq \mathbb{B}$ .

# The closure

We define

$$\text{Exp} : \text{Type} \rightarrow \text{Type}_1$$
$$\text{Exp } I = I \rightarrow \text{Type}$$

## **Theorem**

*Ignoring size issues, for polynomials we have*

$$(I \sqcup J) \rightsquigarrow K \simeq I \rightsquigarrow (\text{Exp } J \times K)$$

We define

$\text{Exp} : \text{Type} \rightarrow \text{Type}_1$

$\text{Exp } I = \Sigma (I \rightarrow \text{Type}) (\lambda F \rightarrow \text{is-finite } (\Sigma I F))$

## **Theorem**

*Ignoring size issues, for finitary polynomials we have*

$$(I \sqcup J) \rightsquigarrow K \simeq I \rightsquigarrow (\text{Exp } J \times K)$$



# The closure

We define

$\text{Exp} : \text{Type} \rightarrow \text{Type}_1$

$\text{Exp } I = \Sigma \text{ FinType } (\lambda N \rightarrow \text{fst } N \rightarrow I)$

## **Theorem**

*Ignoring size issues, for finitary polynomials we have*

$$(I \sqcup J) \rightsquigarrow K \simeq I \rightsquigarrow (\text{Exp } J \times K)$$

# The closure

We define

$\text{Exp} : \text{Type} \rightarrow \text{Type}$

$\text{Exp } I = \Sigma \mathbb{B} (\lambda b \rightarrow \mathbb{B}\text{-to-Fin } b \rightarrow I)$

## **Theorem**

*For finitary polynomials we have*

$$(I \sqcup J) \rightsquigarrow K \simeq I \rightsquigarrow (\text{Exp } J \times K)$$

# The exponential

Note that

$$\text{Exp} : \text{Type} \rightarrow \text{Type}$$
$$\text{Exp } I = \Sigma \mathbb{B} (\lambda b \rightarrow \mathbb{B}\text{-to-Fin } b \rightarrow I)$$

is the free pseudo-commutative monoid!

Questions?