

Coinductive control of inductive data types

Paige Randall North and Maximilien Péroux

Published in CALCO 2023
arXiv:2303.16793

19 February 2024

Outline

Overview and background

Endofunctors

Overview

Theorem (N.-Péroux)

The category of algebras over an *accessible, lax symmetric monoidal* endofunctor on a *locally presentable, symmetric monoidal closed* category is enriched over the category of coalgebras of the same endofunctor.

Overview

Theorem (N.-Péroux)

The category of algebras over an *accessible, lax symmetric monoidal* endofunctor on a *locally presentable, symmetric monoidal closed* category is enriched over the category of coalgebras of the same endofunctor.

Examples

There are many examples, including polynomial endofunctors with extra structure.

Overview

Theorem (N.-Péroux)

The category of algebras over an *accessible, lax symmetric monoidal* endofunctor on a *locally presentable, symmetric monoidal closed* category is enriched over the category of coalgebras of the same endofunctor.

Examples

There are many examples, including polynomial endofunctors with extra structure.

Gain

Get more control over algebras

- ▶ Get more “initial algebras” (e.g. generalized W-types)

Review of categorical W-types

Let \mathcal{C} be a locally presentable, symmetric monoidal closed category, i.e. \mathbf{Set} .

Natural numbers

The type of natural numbers \mathbb{N} is the initial algebra for the endofunctor $X \mapsto X + 1$.

This endofunctor fulfills our hypotheses.

Lists

The type of lists $\mathbb{List}(A)$ is the initial algebra for the endofunctor $X \mapsto X \times A + 1$.

When A is equipped with the structure of a commutative monoid, this fulfills our hypotheses.

Previous work on coalgebraic enrichment

Universal measuring coalgebra (Wraith, Sweedler 1968)

For k -algebras A and B , there is a k -coalgebra $\underline{\text{Alg}}(A, B)$

- ▶ which underlies an enrichment of k -algebras in k -coalgebras
- ▶ whose *set-like elements*¹ are in bijection with $\text{Alg}(A, B)$.

Taking $B := k$, one gets the *dual* $\underline{\text{Alg}}(A, k)$ of A .

Extensions

- ▶ Anel-Joyal 2013 (dg-algebras)
- ▶ Hyland-Franco-Vasilakopoulou 2017 (monoids)
- ▶ Vasilakopoulou 2019 (\mathcal{V} -categories)
- ▶ Péroux 2022 (∞ -algebras of an ∞ -operad)
- ▶ McDermott-Rivas-Uustalu 2022 (monads)

¹those $c \in \text{Alg}(A, B)$ s.t. $\Delta c = c \otimes c$ and $\epsilon(c) = 1_A$

Enriched categories

Definition

An *enrichment* of a category \mathcal{C} in a monoidal category \mathcal{V} consists of

- ▶ a functor $\underline{\mathcal{C}}(-, -) : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{V}$
- ▶ a morphism $\mathbb{1} \rightarrow \underline{\mathcal{C}}(A, A)$ for each $A \in \text{ob } \mathcal{C}$
- ▶ a morphism $\underline{\mathcal{C}}(A, B) \otimes \underline{\mathcal{C}}(B, C) \rightarrow \underline{\mathcal{C}}(A, C)$ for $A, B, C \in \text{ob } \mathcal{C}$
- ▶ an isomorphism $\mathcal{V}(\mathbb{1}, \underline{\mathcal{C}}(A, B)) \cong \mathcal{C}(A, B)$ for $A, B \in \text{ob } \mathcal{C}$.

such that ...

Remark

Monoidal *closed* means enriched in itself.

Measuring in general

Fix a *locally presentable, symmetric monoidal closed category* \mathcal{C} and an *accessible, lax symmetric monoidal endofunctor* F .

Measuring

For algebras (A, α) , (B, β) a *measure* $(A, \alpha) \rightarrow (B, \beta)$ is a coalgebra (C, χ) together with a morphism $\phi : C \rightarrow \underline{\mathcal{C}}(A, B)$ satisfying:

$$\begin{array}{ccccccc}
 & & & FC & \xrightarrow{F(\phi)} & F(\underline{\mathcal{C}}(A, B)) & \longrightarrow & \underline{\mathcal{C}}(FA, FB) \\
 & \nearrow \chi & & & & & & \downarrow \beta \\
 C & & & & & & & \\
 & \searrow \phi & & \underline{\mathcal{C}}(A, B) & \xrightarrow{\alpha} & \underline{\mathcal{C}}(FA, B) & &
 \end{array}$$

The *universal measure* $\underline{\text{Alg}}(A, B)$ is the terminal one.

Measuring in general

Fix a *locally presentable, symmetric monoidal closed category* \mathcal{C} and an *accessible, lax symmetric monoidal endofunctor* F .

Measuring

For algebras $(A, \alpha), (B, \beta)$ a *measure* $(A, \alpha) \rightarrow (B, \beta)$ is a coalgebra (C, χ) together with a morphism $\phi : C \rightarrow \underline{\mathcal{C}}(A, B)$ satisfying:

$$\begin{array}{ccccc}
 & & FC & \xrightarrow{F(\phi)} & F(\underline{\mathcal{C}}(A, B)) & \longrightarrow & \underline{\mathcal{C}}(FA, FB) \\
 & \nearrow \chi & & & & & \downarrow \beta \\
 C & & & & & & \\
 & \searrow \phi & \underline{\mathcal{C}}(A, B) & \xrightarrow{\alpha} & \underline{\mathcal{C}}(FA, B) & &
 \end{array}$$

The *universal measure* $\underline{\text{Alg}}(A, B)$ is the terminal one.

Theorem (N.-Péroux)

The universal measure $\underline{\text{Alg}}(A, B)$ always exists, and these are the hom-coalgebras of an enrichment of $\text{Alg}(F)$ in $\text{CoAlg}(F)$.

Measuring for the natural numbers

Consider the endofunctor $X \mapsto X + 1$ on Set .

- ▶ Algebras are sets A together with $A + 1 \rightarrow A$
 - ▶ Have $-_A : \mathbb{N} \rightarrow A$
- ▶ Coalgebras are sets C together with $A \rightarrow A + 1$
 - ▶ Have $\llbracket - \rrbracket : C \rightarrow \mathbb{N}^\infty$

Measuring

For algebras A, B , a *measure* $A \rightarrow B$ is a coalgebra C together with a function $C \rightarrow A \rightarrow B$ such that

- ▶ $f_c(0_A) = 0_B$ for all $c \in C$;
- ▶ $f_c(a + 1) = 0_B$ for all $\llbracket c \rrbracket = 0$ and for all $a \in A$;
- ▶ $f_c(a + 1) = f_{c-1}(a) + 1$ for $\llbracket c \rrbracket \geq 1$ and for all $a \in A$.

The *universal measure* $\underline{\text{Alg}}(A, B)$ is the terminal measure $A \rightarrow B$.

Measuring for the natural numbers

Consider the endofunctor $X \mapsto X + 1$ on Set .

- ▶ Algebras are sets A together with $A + 1 \rightarrow A$
 - ▶ Have $-_A : \mathbb{N} \rightarrow A$
- ▶ Coalgebras are sets C together with $A \rightarrow A + 1$
 - ▶ Have $\llbracket - \rrbracket : C \rightarrow \mathbb{N}^\infty$

Measuring

For algebras A, B , a *measure* $A \rightarrow B$ is a coalgebra C together with a function $C \rightarrow A \rightarrow B$ such that

- ▶ $f_c(0_A) = 0_B$ for all $c \in C$;
- ▶ $f_c(a + 1) = 0_B$ for all $\llbracket c \rrbracket = 0$ and for all $a \in A$;
- ▶ $f_c(a + 1) = f_{c-1}(a) + 1$ for $\llbracket c \rrbracket \geq 1$ and for all $a \in A$.

The *universal measure* $\underline{\text{Alg}}(A, B)$ is the terminal measure $A \rightarrow B$.

What is this?

Set-like elements in general

Definition

The *set-like elements* are

$$\mathbb{1} \rightarrow \underline{\text{Alg}}(A, B) \quad \text{in } \text{CoAlg}(F)$$

i.e., elements of $\text{Alg}(A, B)$.

Set-like elements in general

Definition

The *set-like elements* are

$$\mathbb{1} \rightarrow \underline{\text{Alg}}(A, B) \quad \text{in } \text{CoAlg}(F)$$

i.e., elements of $\text{Alg}(A, B)$.

That is

- ▶ The *points* of $\text{Alg}(A, B)$ are total algebra homomorphisms $A \rightarrow B$.

Set-like elements in general

Definition

The *set-like elements* are

$$\mathbb{1} \rightarrow \underline{\text{Alg}}(A, B) \quad \text{in } \text{CoAlg}(F)$$

i.e., elements of $\text{Alg}(A, B)$.

That is

- ▶ The *points* of $\text{Alg}(A, B)$ are total algebra homomorphisms $A \rightarrow B$.
- ▶ If we're considering $(\text{Set}, \times, *)$, the underlying set of $\mathbb{1}$ is $*$, so these are 'special' elements of the underlying set of $\text{Alg}(A, B)$.

Set-like elements for the natural numbers

Set-like elements

The *set-like elements* are

$$\mathbb{N} \rightarrow \underline{\text{Alg}}(A, B)$$

Set-like elements for the natural numbers

Set-like elements

The *set-like elements* are

$$\mathbb{I} \rightarrow \underline{\text{Alg}}(A, B)$$

where \mathbb{I} has underlying set $\{*\}$ such that $* - 1 = *$

Set-like elements for the natural numbers

Set-like elements

The *set-like elements* are

$$\mathbb{1} \rightarrow \underline{\text{Alg}}(A, B)$$

where $\mathbb{1}$ has underlying set $\{*\}$ such that $* - 1 = *$

so $\mathbb{1} \rightarrow \underline{\text{Alg}}(A, B)$ is an element $* \in \underline{\text{Alg}}(A, B)$ s.t. $* - 1 = *$

Set-like elements for the natural numbers

Set-like elements

The *set-like elements* are

$$\mathbb{1} \rightarrow \underline{\text{Alg}}(A, B)$$

where $\mathbb{1}$ has underlying set $\{*\}$ such that $* - 1 = *$

so $\mathbb{1} \rightarrow \underline{\text{Alg}}(A, B)$ is an element $* \in \underline{\text{Alg}}(A, B)$ s.t. $* - 1 = *$

so f_* is a total algebra homomorphism

Measuring

...

- ▶ $f_c(0_A) = 0_B$ for all $c \in C$;

...

- ▶ $f_c(a + 1) = f_{c-1}(a) + 1$ for $\llbracket c \rrbracket \geq 1$ and for all $a \in A$.

Set-like elements for the natural numbers

Set-like elements

The *set-like elements* are

$$\mathbb{1} \rightarrow \underline{\text{Alg}}(A, B)$$

where $\mathbb{1}$ has underlying set $\{*\}$ such that $* - 1 = *$

so $\mathbb{1} \rightarrow \underline{\text{Alg}}(A, B)$ is an element $* \in \underline{\text{Alg}}(A, B)$ s.t. $* - 1 = *$

so f_* is a total algebra homomorphism

Measuring

...

- ▶ $f_*(0_A) = 0_B;$

...

- ▶ $f_*(a + 1) = f_*(a) + 1$ for all $a \in A.$

Set-like elements for the natural numbers

Set-like elements

The *set-like elements* are

$$\mathbb{1} \rightarrow \underline{\text{Alg}}(A, B)$$

where $\mathbb{1}$ has underlying set $\{*\}$ such that $* - 1 = *$

so $\mathbb{1} \rightarrow \underline{\text{Alg}}(A, B)$ is an element $* \in \underline{\text{Alg}}(A, B)$ s.t. $* - 1 = *$

so f_* is a total algebra homomorphism

that is, an element of $\text{Alg}(A, B)$.

Measuring

...

- ▶ $f_*(0_A) = 0_B;$

...

- ▶ $f_*(a + 1) = f_*(a) + 1$ for all $a \in A$.

Set-like elements for the natural numbers

Set-like elements

The *set-like elements* are

$$\mathbb{1} \rightarrow \underline{\text{Alg}}(A, B)$$

where $\mathbb{1}$ has underlying set $\{*\}$ such that $* - 1 = *$

so $\mathbb{1} \rightarrow \underline{\text{Alg}}(A, B)$ is an element $* \in \underline{\text{Alg}}(A, B)$ s.t. $* - 1 = *$

so f_* is a total algebra homomorphism

that is, an element of $\text{Alg}(A, B)$.

Example

$$\text{Alg}(\mathbb{N}, A) \cong *$$

$$\underline{\text{Alg}}(\mathbb{N}, A) \cong \mathbb{N}^\infty$$

What are the non-set-like elements?

Example

$$\underline{\text{Alg}}(\mathbb{N}, A) \cong \mathbb{N}^\infty$$

What are the non-set-like elements?

Example

$$\underline{\text{Alg}}(\mathbb{N}, A) \cong \mathbb{N}^\infty$$

So denote the elements of $\underline{\text{Alg}}(\mathbb{N}, A)$ by

- ▶ f_0
- ▶ f_1
- ▶ \dots
- ▶ f_∞

Measuring

...

- ▶ $f_0(0) = 0_B$
- ▶ $f_0(a + 1) = 0_B$ for all $a \in A$

What are the non-set-like elements?

Example

$$\underline{\text{Alg}}(\mathbb{N}, A) \cong \mathbb{N}^\infty$$

So denote the elements of $\underline{\text{Alg}}(\mathbb{N}, A)$ by

- ▶ $f_0(n) = 0_A$
- ▶ f_1
- ▶ \dots
- ▶ f_∞

Measuring

...

- ▶ $f_0(0) = 0_B$
- ▶ $f_0(a + 1) = 0_B$ for all $a \in A$

What are the non-set-like elements?

Example

$$\underline{\text{Alg}}(\mathbb{N}, A) \cong \mathbb{N}^\infty$$

So denote the elements of $\underline{\text{Alg}}(\mathbb{N}, A)$ by

- ▶ $f_0(n) = 0_A$
- ▶ f_1
- ▶ \dots
- ▶ f_∞

Measuring

...

- ▶ $f_1(0_A) = 0_B$
- ▶ $f_1(a + 1) = f_0(a) + 1$ for all $a \in A$

What are the non-set-like elements?

Example

$$\underline{\text{Alg}}(\mathbb{N}, A) \cong \mathbb{N}^\infty$$

So denote the elements of $\underline{\text{Alg}}(\mathbb{N}, A)$ by

- ▶ $f_0(n) = 0_A$
- ▶ $f_1(0) = 0_A; f_1(sn) = 1_A$
- ...
- ▶ f_∞

Measuring

...

- ▶ $f_1(0_A) = 0_B$
- ▶ $f_1(a + 1) = f_0(a) + 1$ for all $a \in A$

What are the non-set-like elements?

Example

$$\underline{\text{Alg}}(\mathbb{N}, A) \cong \mathbb{N}^\infty$$

So denote the elements of $\underline{\text{Alg}}(\mathbb{N}, A)$ by

- ▶ $f_0(n) = 0_A$
- ▶ $f_1(0) = 0_A; f_1(sn) = 1_A$
- ...
- ▶ f_∞

Measuring

...

- ▶ $f_\infty(0) = 0_B$
- ▶ $f_\infty(a + 1) = f_\infty(a) + 1$

What are the non-set-like elements?

Example

$$\underline{\text{Alg}}(\mathbb{N}, A) \cong \mathbb{N}^\infty$$

So denote the elements of $\underline{\text{Alg}}(\mathbb{N}, A)$ by

- ▶ $f_0(n) = 0_A$
- ▶ $f_1(0) = 0_A; f_1(sn) = 1_A$
- ...
- ▶ $f_\infty(n) = n_A$

Measuring

...

- ▶ $f_\infty(0) = 0_B$
- ▶ $f_\infty(a + 1) = f_\infty(a) + 1$

What are the non-set-like elements?

Example

$$\underline{\text{Alg}}(\mathbb{N}, A) \cong \mathbb{N}^\infty$$

So denote the elements of $\underline{\text{Alg}}(\mathbb{N}, A)$ by

- ▶ $f_0(n) = 0_A$
- ▶ $f_1(0) = 0_A; f_1(sn) = 1_A$
- ...
- ▶ $f_\infty(n) = n_A$

Definition

So we call elements of the underlying of $\underline{\text{Alg}}(A, B)$ *n-partial algebra homomorphisms*.

What are the non-set-like elements?

- ▶ Let \mathfrak{n} denote the quotient of \mathbb{N} by $m = n$ for all $m \geq n$.
- ▶ Let \mathfrak{n}° denote the subobject of \mathbb{N}^∞ consisting of $\{0, \dots, n\}$.

Example

$$\text{Alg}(\mathfrak{n}, A) \cong \begin{cases} * & \text{if } n_A = m_A \text{ for all } m \geq n; \\ \emptyset & \text{otherwise.} \end{cases}$$

What are the non-set-like elements?

- ▶ Let \mathfrak{n} denote the quotient of \mathbb{N} by $m = n$ for all $m \geq n$.
- ▶ Let \mathfrak{n}° denote the subobject of \mathbb{N}^∞ consisting of $\{0, \dots, n\}$.

Example

$$\text{Alg}(\mathfrak{n}, A) \cong \begin{cases} * & \text{if } n_A = m_A \text{ for all } m \geq n; \\ \emptyset & \text{otherwise.} \end{cases}$$

$$\underline{\text{Alg}}(\mathfrak{n}, A) \cong \begin{cases} \mathbb{N}^\infty & \text{if } n_A = m_A \text{ for all } m \geq n; \\ \mathfrak{n}^\circ & \text{otherwise.} \end{cases}$$

- ▶ So there is at least always an n -partial homomorphism out of n (which is unique).

What can we do with this?

Generalize W-types, i.e., initial algebras.

C-initial objects

For a coalgebra C , a *C-initial algebra* is the terminal algebra A such that for all other algebras B there is a unique

$$C \rightarrow \underline{\text{Alg}}(A, B).$$

Initial object

An initial object in a category \mathcal{C} is the (terminal) object A such that for all other algebras B there is a unique

$$* \rightarrow \mathcal{C}(A, B).$$

C-initial objects for the natural numbers

Examples

For the natural-numbers endofunctor:

- ▶ \mathbb{N} is the $\mathbb{1}$ -initial algebra
- ▶ \mathbb{N} is the \mathbb{N}^{∞} -initial algebra

C-initial objects for the natural numbers

Examples

For the natural-numbers endofunctor:

- ▶ \mathbb{N} is the $\mathbb{1}$ -initial algebra
- ▶ \mathbb{N} is the \mathbb{N}^∞ -initial algebra

- ▶ $\mathbb{1}$ - (or \mathbb{N}^∞ -) initial means initial with respect to total algebra homomorphisms

Theorem

n is the n° -initial algebra

- ▶ n° -initial means initial with respect to partial algebra homomorphisms

Examples

(Endofunctors on a locally presentable symmetric monoidal category)

(id) The identity endofunctor

(A) The constant endofunctor at fixed commutative monoid A

(GF) The composition of two instances

($F \otimes G$) The tensor of two instances (\mathcal{C} closed)

($F + G$) The coproduct of an instance F and an ' F -module' G

(id^A) The exponential id^A at object A (\mathcal{C} cartesian closed)

(W -type) The polynomial endofunctor associated to a morphism $f : X \rightarrow Y$, given a commutative monoid structure on Y and an oplax symmetric monoidal structure on the preimage functor $f^{-1} : \mathcal{C} \rightarrow \text{Set}$ ($\mathcal{C} = \text{Set}$)

(d.e.s.) A discrete equational system (monoidal structure on \mathcal{C} is cocartesian, \mathcal{C} has binary products that preserve filtered colimits)

Summary

We have

- ▶ that algebras are enriched in coalgebras (under certain hypotheses)
- ▶ an interpretation as notion of partial algebra homomorphism (especially in the case N)
- ▶ many examples
- ▶ a more refined notion of initial algebra

Conclusion

- ▶ Summary
 - ▶ Work out more of the examples in detail
 - ▶ Understand C -initial algebras in more examples and in general
 - ▶ Understand if this extra structure is useful for programming languages
 - ▶ Understand if there is a connection with domain theory
- ▶ Future work
 - ▶ that algebras are enriched in coalgebras (under certain hypotheses)
 - ▶ an interpretation as notion of partial algebra homomorphism (especially in the case N)
 - ▶ many examples
 - ▶ a more refined notion of initial algebra

Thank you!