Interpolation as Cut-introduction

² On the Computational Content of Craig-Lyndon Interpolation

₃ Alexis Saurin ⊠

4 IRIF – CNRS, Université Paris Cité & INRIA Picube

⁵ — Abstract

Analyzing Maehara's method for proving Craig's interpolation theorem, we extract a "proof relevant" 6 interpolation theorem for first-order LL in the sense that if π is a cut-free sequent proof of $A \vdash B$, we can find a formula C in the common vocabulary of A and B and proofs π_1, π_2 of $A \vdash C$ and $C \vdash B$ respectively such that π_1 composed with π_2 cut-reduces to π . As a direct corollary, we get 9 similar proof relevant interpolation results for LJ and LK using linear translations. This refined 10 interpolation is then rephrased in terms of a cut-introduction process synthetizing the interpolant. 11 Finally, we analyze the computational content of interpolation by proving and interpolation 12 result for Curien and Herbelin's Duality of Computation. 13 14 The present document uses color: a color-blind-friendly and printable version is available at https:

15 //www.irif.fr/_media/users/saurin/pub/interpolation_as_cut_introduction-cb.pdf.

16 2012 ACM Subject Classification

Keywords and phrases Classical Logic, Interpolation, Cut Elimination, Linear Logic, Sequent
 calculus, System L

¹⁹ Digital Object Identifier 10.4230/LIPIcs...

²⁰ 1 "- Why Not a Proof-Relevant Interpolation Theorem? ²¹ - Introduce Cuts, Of Course!"

In the words of Solomon Feferman, "though deceptively simple and plausible on the face 22 of it, Craig's interpolation theorem (...) has proved to be a central logical property that 23 has been used to reveal the deep harmony between the syntax and semantics of first order 24 logic" [17]. Indeed, Craig's interpolation (which states that in the predicate calculus, if 25 $A \vdash B$, there exists a formula C built from the relation symbols occurring both in A and B 26 such that $A \vdash C$ and $C \vdash B$) and its developments suggest far deeper connections between 27 models and proofs than the simple correspondence between provability and validity given by 28 Gödel completeness theorem. This could be argued to be in line with and pursue structural 29 proof-theoretic proofs of Gödel completeness theorem such as Schütte proof [36] or the more 30 recent analysis by Basaldella and Terui of completeness in Ludics [3, 2]. 31

First of all, one should recall that while the original proof of interpolation by Craig [7, 8] 32 was proof-theoretic as well as Maehara's method [25] its most striking applications were model-33 theoretic results whether we consider standard model-theoretic results that are consequences 34 of interpolation, such as Beth definability theorem [4] or Robinson's consistency theorem [33] 35 or if we consider modern uses of interpolation in model-checking [21, 27]. Among the proof-36 theoretical methods for proving interpolation, the success of Maehara's method is probably 37 due to its wide applicability to a range of logics and proof-systems, from intuitionistic 38 logic [28, 36] to modal logics [18, 24, 1, 37] or in infinitary logics and abstract model 39 theory [17, 15]. 40

While in most proof theory textbooks [20, 36, 40, 41] Craig's interpolation theorem is presented as an application of cut-elimination, one shall see here that it also has in fact much to do with cut-introduction, giving a proof-relevant and computational content to Interpolation theorem. This opens the way to an analysis of interpolation in terms of the denotational semantics of proofs and programs.



© Author: Please provide a copyright holder; licensed under Creative Commons License CC-BY 4.0

Leibniz International Proceedings in Informatics

LIPICS Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

XX:2 Interpolation as cut-introduction

Contributions and organization of the paper. More precisely, after providing back-46 ground on linear logic sequent calculus in Section 2, we state in Section 3 the following result 47 for first-order linear logic (LL in the following): For any first-order LL formulas A, B, if π 48 proves $A \vdash B$, there exists a formula C in the common vocabulary (that refers to the subset 49 of the first-order language occurring in a formula, in terms of relation symbols, possibly 50 taking into account polarity constraints as in Lyndon's interpolation) of A and B and proofs 51 π_1, π_2 of $A \vdash C$ and $C \vdash B$ respectively such that $(Cut)(\pi_1, \pi_2)$, that is π_1 composed with π_2 , 52 is equivalent to π (by $=_{cut}$ we refer to the congruence generated by the cut-reduction rules): 53

$$\frac{A \stackrel{n_1}{\vdash} C \quad C \stackrel{n_2}{\vdash} B}{A \vdash B} \quad (\mathsf{Cut}) \ =_{\mathsf{cut}} A \stackrel{\pi}{\vdash} B \cdot$$

Interpolation can therefore be achieved while preserving the computational / denotational
content of proofs, while factoring the computation through an interfacing, interpolant type
made only of the base types used in both the input and output types.

This result is then extended to classical and intuitionistic logics simply by means of linear translations, which sheds an interesting light on the relationship between Lyndon and Craig's interpolation.

Then, by a further analysis of Maehara's method, we show in Section 4 that the interpolation process for a cut-free proof deriving $A \vdash B$ can be in fact decomposed in two phases: (i) an ascending phase which equips each sequent of π with a splitting is followed by (ii) a descending phase which solves the interpolation problem. This latter descending phase happens to be a cut-introduction phase which provides a proof of the proof-relevant interpolation result of the previous section. In particular, the resulting proof is, *by construction*, denotationally equal to π .

Finally, in Section 5, we consider the computational content of interpolation and address this question using the framework of the Duality of Computation introduced by Curien and Herbelin: we consider the linear system L, a term calculus with typing rules very close to that of the sequent calculus¹, and prove a computational interpolation result.

Related works. Surprisingly we could not find any occurrence in the literature analyzing
 Maehara's method in terms of cut-elimination (or rather, cut-introduction) even though *all ingredients* were there since Maehara's seminal work.

On the other hand, another early proof-theoretic proof of interpolation theorem was pro-74 posed by Prawitz for natural deduction [31]. Just like for Maehara's method the strengthened, 75 proof-relevant interpolation result was at hand in this work as well and Cubrić actually 76 showed this in the setting of the simply typed λ -calculus as well as a corresponding factoriza-77 tion result for bicartesian closed categories in the early 90s [9, 10]. Sadly, Cubrić's paper 78 as well as his PhD thesis supervised by Makkai, received too little attention and very few 79 following works refer to his results: we could only find less than 10 references to these works 80 among them only three truly consider the interpolation aspect $[16, 26, 23]^2$. We hope that 81 the present work can contribute to foster interest in Cubrić results. 82

¹ System L can be viewed as achieving a Curry-Howard correspondence with sequent calculus proofs – while the λ -calculus achieves a correspondence with natural deduction proofs

² Matthes [26] extends Čubrić' results to a natural deduction with general elimination rules and a corresponding term calculus while Kanazawa [23] considers interpolation in purely implicational fragments of intuitionistic logic and finds workarounds for the lack of interpolation in this setting. On the way, he considers various sequent calculi and proves a result which has some similarities with our result but is weaker both in terms of the logical language – which is restricted to implicative LJ– and of the characterization of the equivalence between the interpolating proofs and the original proof.

Another related work is that of Carbone [6] where she establishes a strengthened form 83 of Maehara's interpolation paying great attention to the ancestor relation (formulated in 84 terms of flow graphs in that work) which allows her to get bounds on the complexity of the 85 interpolant but did not lead her to a study of proof-relevant interpolation, invariance by 86 cut-elimination nor interpolation as cut-introduction. Only few works consider interpolation 87 in (fragments of) linear logic, starting with Roorda [34]. In the framework of the calculus of 88 structure, Straßburger proves a decomposition theorem for MELL [38] that is advocated to 89 correspond to an interpolation theorem and may have a more fine-grained proof-theoretical 90 content. More recently, several papers investigated and formalized interpolation theorems in 91 substructural logics, including exponential-free linear logic [5, 14, 32]. 92

2 Background on LL proof theory

⁹⁴ In the following, we provide the necessary background on first-order LL.

As usual, we assume a first-order language \mathcal{L} (without equality nor function symbols). We assume the set of atomic formulas to be equipped with an involution, inducing a partitioning between **positive atomic formulas**, written *a*, and **negative atomic formulas**, written a^{\perp} . We introduce a language of first-order LL formulas:

⁹⁹ ► Definition 1. The grammar of first-order LL formulas is defined inductively as:

100

Linear negation is defined as usual as an involution on LL formulas (changing a^{\perp} , \perp , \top , \otimes , \otimes , \forall , ? into a, 1, 0, \otimes , \oplus , \exists , ! respectively). A one-sided LL sequent is an ordered list of LL formulas usually written $\vdash \Gamma$ to emphasize the sequent judgments.

Definition 2 (LL sequent calculus). The inference rules of LL sequent calculus are given in Figure 1. The inferences of Figure 1 show a relation between conclusion and premises formulas, the ancestor relation (or sub-occurrence relation) that is extended by transitivity to formulas of non consecutive sequents of a derivation. An LL proof of a sequent $\vdash \Gamma$ is a finite tree rooted in $\vdash \Gamma$ generated by the rules of Figure 1.

Remark 3. The ancestor relation defined above is (implicitly) used in designing a cutreduction system and plays a crucial role in expressing the validity condition for nonwellfounded and circular proofs as well as in the extension of Maehara's method we will show next, in order to propagate sequent splittings from conclusions to premises.

To state Craig-Lyndon interpolation, we need to make clear the required notion to express the interpolation condition on the vocabulary:

▶ Definition 4 ((polarized) vocabulary of a formula). Given an LL formula F, Voc(F) is the set of all predicate symbols occurring in F together with their linear negations, while $Voc^+(F)$ (resp. $Voc^-(F)$) is the set of positive (resp. negative) predicate symbols occurring in F. This is naturally lifted to lists of formulas.

¹²⁰ In the present paper, we shall heavily rely on LL cut-reduction rules (and their symmetric, ¹²¹ cut-introduction rules). Even though those rules are standard, we recall them in Appendix A ¹²² for completeness.

XX:4 Interpolation as cut-introduction

Figure 1 (a) Propositional MALL Inferences; (b) LL Exponential Inferences; (c) First-order Inferences;

¹²³ **3** Proof-relevant interpolation theorem

The following proof-relevant interpolation theorem can be established by refining Maehara's proof in order to take seriously into account the relations, not only between the interpolation sequents, but also between the interpolating proofs. This is presented in detail in Appendix B.

Theorem 5. Let Γ, Δ be lists of LL formulas and $\pi \vdash \Gamma, \Delta$ cut-free. There exists a LL formula C such that $\operatorname{Voc}^+(C) \subseteq \operatorname{Voc}^-(\Gamma) \cap \operatorname{Voc}^+(\Delta)$ and $\operatorname{Voc}^-(C) \subseteq \operatorname{Voc}^+(\Gamma) \cap \operatorname{Voc}^-(\Delta)$ and two cut-free proofs π_1, π_2 of $\vdash \Gamma, C$ and $\vdash C^{\perp}, \Delta$ respectively such that

$$\frac{\vdash \overset{\pi_1}{\Gamma,C} \quad \vdash \overset{\pi_2}{C^{\perp}}, \Delta}{\vdash \Gamma,\Delta} \quad \underset{(\mathsf{Cut})}{\longrightarrow} \overset{\star}{\underset{\mathsf{cut}}} \quad \pi.$$

In the following sections, we will follow two approaches which we consider to be more instructive from the computer science logic viewpoint, as they reveal the computational content of proof-relevant interpolation. We will proceed by "introducing cuts" so that we preserve the denotational equivalence of the *interpolated proof* with its interpolation (proofs).

But first, let us extend the proof-relevant interpolation result to LK and LJ. These results could be obtained directly, by refining Maehara's method as done in Appendix B or by following the cut-introduction or computational approach of the following section. On the other hand, one can directly deduce this result by using the linear embeddings of LK and LJ [19, 13] as a corollary of the above theorem.

► Corollary 6. Let Γ, Γ', Δ, Δ' be lists of LK formulas (resp. of LJ formulas, with Δ being empty and Δ' of length at most 1) and π be a cut-free LK (resp. LJ) proof of Γ, Γ' ⊢ Δ, Δ'. There exists a LK (resp. LJ) formula C such that Voc⁺(C) ⊆ Voc⁺(Γ, Δ) ∩ Voc⁺(Γ', Δ') and Voc⁻(C) ⊆ Voc⁻(Γ, Δ) ∩ Voc⁻(Γ', Δ') and two cut-free LK (resp. LJ) proofs π₁, π₂ of Γ ⊢ C, Δ and Γ', C ⊢ Δ' respectively such that $\frac{\Gamma ⊢ C, Δ ~ \Gamma', C ⊢ Δ'}{\Gamma, \Gamma' ⊢ Δ, Δ'} \xrightarrow{(Cut)} →_{cut}^{*} π.$

¹⁴⁵ **Proof.** We sketch this here for LK and provide details in appendices:

First, it is a matter of a simple check that proof-relevant interpolation presented in the above theorem also holds for the two-sided LL sequent calculus.

Second, a cut-free LK proof π (it would be similar for LJ) can be decorated with exponential modalities and inferences in order to turn it into a cut-free LL proof π' .

After interpolating this proof (obtaining I', π'_1 and π'_2), one can erase the linear information of the interpolants and the two interpolating proofs (that is, taking the classical *skeleton* of the proofs) and get back a pair of LK (resp. LJ) proofs π_1, π_2 together with a formula I in LK (resp. LJ) that lives in the appropriate vocabulary.

The properties of the linear embeddings ensure that the skeleton of a cut-free proof obtained from $Cut(\pi'_1, \pi'_2)$ can be obtained by eliminating cuts from $Cut(\pi_1, \pi_2)$.

¹⁵⁶ **4** Interpolation as cut-introduction

¹⁵⁷ We will now show how the proof-relevant interpolation theorem stated in the previous section ¹⁵⁸ can be established by exploiting the dynamics of cut-introduction that is, more precisely,

¹⁵⁹ how the synthesis of the interpolant is in fact a cut-introduction process.

¹⁶⁰ For this, we need to analyze and refine Maehara's proof method.

¹⁶¹ 4.1 Refining Maehara's method

The usual statement and proof method for interpolation, made more informative in the previous section (and Appendix B), actually obfuscate the fact that the interpolating formula and proofs are synthetized using a cut-introduction mechanism: this is due to the structure of the inductive reasoning used to establish the interpolation theorem under Maehara's method. Indeed, the inductive structure of the reasoning amounts to inspecting the structure of the sequent proofs until one reaches the base cases (the aximo and unit cases), after which the call to the induction hypotheses synthetize the interpolant.

Analyzing what happens in constructing the interpolating formula and proofs is made clearer by structuring the process in two phases, a bottom-up phase and a top-down phase³:

Ascending phase. This first phase consists in traversing the initial proof π bottom-up, from root (conclusion) to leaves (axioms), and building, for each visited sequent Γ , a splitting (Γ', Γ'') inherited from the splitting of the conclusion of the proof by the ancestor relation. In this way, each node of the proof is ultimately decorated with some additional information on how to split the sequent labelling the node. We will use the red-blue colors throughout the paper to represent such splittings, and from now on, enrich sequents with such splitting information.

Ultimately, for each logical axiom rule $\vdash A^{\perp}, A$, we are in one of the following situations: (i) $(\{A^{\perp}, A\}, \emptyset)$; (ii) $(\{A^{\perp}\}, \{A\})$; (iii) $(\{A\}, \{A^{\perp}\})$; (iv) $(\emptyset, \{A^{\perp}, A\})$ which are summed up with the color code as: $\vdash A^{\perp}, A$; $\vdash A^{\perp}, A$; $\vdash A^{\perp}, A$; $\vdash A^{\perp}, A$. (and similarly for each axiom corresponding to some unit, \top or 1.) This corresponds to the various base cases of the inductive proof of the previous section.

¹⁸³ Once every axiom has been reached, we switch to the descending phase, traversing again

the proof, top to bottom, in an asynchronous manner.

³ Note that the "top-down" and "bottom-up" terminologies refer to the usual tree presentation of sequent proofs, where the root of below, rather than to the usual view of terms: we move upward in the proof while inspecting subproofs.

¹⁸⁵ **Descending phase.** Equipped with the sequents splitting information one shall now apply ¹⁸⁶ cut-introduction rules to axioms, progressively moving the cuts down and merging them ¹⁸⁷ in such a way, ultimately, to reach the root sequent of the original proof. We call *active* a ¹⁸⁸ sequent such that all its premises are concluded with cut inferences. (Initially, since π is

¹⁸⁹ cut-free, only the conclusions of logical axioms or 0-ary unit rules are *active sequents*.)

¹⁹⁰ We apply cut-introductions to active sequents, maintaining the following two invariants:

¹⁹¹ = when a sequent is active with splitting (Γ', Γ'') , the cut formulas of its premises are ¹⁹² interpolants for the premise sequents wrt. their splitting (Note that this condition ¹⁹³ trivially satisfied initially since the active axioms have no premise).

¹⁹⁴ = when an inference (r) has conclusion S which is active, we apply a (sequence of) cut-¹⁹⁵ introduction step(s) on this inference, in such a way that (i) S becomes the conclusion ¹⁹⁶ of the introduced cut and (ii) the premises of this cut correspond to the splitting ¹⁹⁷ associated with sequent S.

¹⁹⁸ The descending phase therefore terminates when the cut reaches the root.

¹⁹⁸ Sequencing these two phases, $\vdash \Gamma, \Delta$ is to be interpolated as some $\frac{\vdash \Gamma, I}{\vdash \Gamma, \Delta} = \frac{\pi^R}{\vdash \Gamma, \Delta}$ (Cut).

Ultimately, one therefore builds a cut formula I and two cut-free proofs π_1 and π_2 such that C is an interpolant of the conclusion sequent with respect to the original splitting and such that $(C_{ut})(\pi_1, \pi_2) \longrightarrow_{cut}^* \pi$, a condition which is satisfied by construction.

To state precisely the properties of the two phases, we introduce the following notions:

A decorated proof is an LL proof such that each sequent is equipped with a splitting.
 A coherent decorated proof is such that for each node, the splitting of the conclusion and of its premises is coherent with respect to the ancestor relation: a formula belonging to the left (resp. right) component of the splitting has all its ancestors belonging to the left (resp. right) component of the splitting. We shall now consider only such coherent

proofs. (Of course, the notion of coherent decorations can be refined to two-sided calculi;
 see Appendix D.)

211 With the above notions, the following lemma is clear by induction on the proof:

▶ Lemma 7. For any LL proof and any splitting of its conclusion sequent, the ascending phase terminates with a coherent decorated proof.

4.2 PRIS: Proof-Relevant Interpolation Situation

We shall now focus on the top-down phase and, in order to make formal the discussion above, we shall introduce a useful class of coherent decorated proofs, called PRIS for proof-relevant interpolation situation; they are essentially partially solved interpolation problems:

Definition 8 (Proof-relevant Interpolation Situation). A PRIS for (Γ, Δ) is the data of:

- **the goal**, that is a cut-free LK proof π of conclusion $\vdash \Gamma, \Delta$ and with $n \ge 0$ open premises ($\vdash \Gamma_i, \Delta_i$)_{1 \le i \le n} such that for each $1 \le i \le n$ the formulas in Γ_i (resp. Δ_i) are ancestors of formulas in Γ (resp. of Δ);
- **the partial interpolants,** that is for each $1 \le i \le n$, a formula I_i st. $\operatorname{Voc}^+(I_i) \subseteq \operatorname{Voc}^-(\Gamma_i) \cap \operatorname{Voc}^+(\Delta_i)$ and $\operatorname{Voc}^-(I_i) \subseteq \operatorname{Voc}^+(\Gamma_i) \cap \operatorname{Voc}^-(\Delta_i)$ and;
- The partial solutions, that is, for each $1 \leq i \leq n$, derivations π_i^L (resp. π_i^R) of conclusion $\vdash \Gamma_i, I_i$ (resp. $\vdash I_i^{\perp}, \Delta_i$).



Figure 2 A PRIS– Proof-relevant Interpolation Situation

A PRIS as given by the above definition will be graphically represented as in Figure 2.

In order to make clear our methodology of exploiting PRIS, we define some special cases of PRIS which have a specific role in the forthcoming development:

Definition 9 (Initial, solved, and elementary PRIS). We define the following cases of PRIS:

and a n initial PRIS is a PRIS with <math>n = 0. It thus has the following form:

$$\overline{\pi} \\ \vdash \Gamma, \Delta$$

with π being a cut-free coherent decorated proof. This is the initial situation of an interpolation problem.

A solved PRIS is a PRIS with n = 1 and π being reduced to the trivial derivation constituted only of an open premise node $\vdash \Gamma, \Delta$, of the form:

$$\pi = \frac{\vdash \frac{\pi_1^L}{\Gamma, I_1} \quad \vdash \frac{\pi_1^R}{I_1^\perp, \Delta}}{\vdash \Gamma, \Delta} \quad \text{(Cut)}$$

A solved PRIS therefore corresponds to the solution of an interpolation problem, with I_1 being the interpolant and π_1^L, π_1^R being the interpolating proofs.

An elementary PRIS is a PRIS such that its goal is reduced to an instance of a single
 n-ary inference rule (r), together with n open premises: it is a PRIS where there remains
 a single inference rule to solve in order to obtain a solution to an interpolation problem.
 It thus has the form:

$$\pi = \frac{ \begin{array}{c} \pi_{1}^{L} & \pi_{1}^{R} \\ \Gamma_{1}, I_{1} & \Gamma_{1}^{L}, \Delta_{1} \\ \hline \\ \mu \\ \Gamma_{1}, \Delta_{1} \end{array} (Cut) \\ \hline \\ \mu \\ \Gamma_{n}, \Delta_{n} \end{array} + \frac{ \begin{array}{c} \pi_{n}^{L} \\ \Gamma_{n}, I_{n} \\ \hline \\ \mu \\ \Gamma_{n}, \Delta_{n} \end{array} (Cut) \\ \hline \\ \mu \\ \Gamma_{n}, \Delta_{n} \end{array} (Cut)$$

241 4.3 Solving PRIS

Our proof-relevant interpolation problem can therefore be rephrased as: *How to relate initial* and solved PRIS via cut-introduction? The crucial step lies in the following lemma:

XX:8 Interpolation as cut-introduction

Lemma 10. For any n-ary inference rule (r) and any elementary PRIS π of the form:

$$\frac{ \frac{\pi_{1}^{L}}{\vdash \Gamma_{1}, I_{1}} \quad \frac{\pi_{1}^{R}}{\vdash I_{1}^{\perp}, \Delta_{1}}}{\frac{\vdash \Gamma_{1}, \Delta_{1}}{\vdash \Gamma_{1}, \Delta_{1}}} \quad (Cut) \qquad \qquad \frac{\frac{\pi_{n}^{L}}{\vdash \Gamma_{n}, I_{n}} \quad \frac{\pi_{n}^{R}}{\vdash I_{n}^{\perp}, \Delta_{n}}}{\vdash \Gamma_{n}, \Delta_{n}} \quad (Cut)$$

there exist I, π^L, π^R such that $\pi' = \frac{\vdash \overset{\pi^L}{\Gamma}, I \quad \vdash \overset{\pi^R}{I^\perp}, \Delta}{\vdash \Gamma, \Delta}$ (Cut) is a solved PRIS and $\pi \longleftarrow_{\mathsf{cut}}^{\star} \pi'$.

Since, each application of the above lemma reduces by one inference the size of the *goal* part of an interpolation situation, each sequence of cut-introduction obtained by application of Lemma 10 terminates in a solved PRIS, that is an interpolated proof. As a conclusion, we obtain the following corollary:

▶ Corollary 11. Any initial PRIS can be reduced, by cut-expansions, to a solved PRIS.

As a conclusion, we obtain the expected main theorem (Theorem 5) by working in a *reversed way*, using cut-introduction:

Theorem 12. Let A, B be LL formulas and π be a cut-free LL proof of $A \vdash B$.

There exists a LL formula C such that $\operatorname{Voc}^+(C) \subseteq \operatorname{Voc}^+(A) \cap \operatorname{Voc}^+(B)$ and $\operatorname{Voc}^-(C) \subseteq \operatorname{Voc}^-(A) \cap \operatorname{Voc}^-(B)$ and two cut-free LL proofs π_1, π_2 of $A \vdash C$ and $C \vdash B$ respectively such that $A \stackrel{\pi_1}{\vdash} C \quad C \stackrel{\pi_2}{\vdash} B \xrightarrow[(Cut)]{} \longrightarrow_{cut}^{\star} \pi.$

256

257 4.4 Proof of Lemma 10

We now prove the main lemma (full details are given in Appendix D.1) which proceed by a case analysis on the elementary PRIS under consideration.

²⁶⁰ 4.4.1 Axiom case

 $If \pi = \overline{\vdash F, F^{\perp}}^{(Ax)}, take I = F^{\perp}, \pi_1^l = \overline{\vdash F, F^{\perp}}^{(Ax)} and \pi_1^r = \overline{\vdash F, F^{\perp}}^{(Ax)}.$ $The cut between \pi_1^l and \pi_1^r reduces to \pi by one cut-axiom reduction step.$ $If \pi = \overline{\vdash F, F^{\perp}}^{(Ax)}, one takes I = \perp, \pi_1^l = \frac{\pi}{\vdash F, F^{\perp}, \perp}^{(L)} and \pi_1^r = \overline{\vdash I}^{(1)}.$ $The cut of \pi_1^l and \pi_1^r reduces to \pi by a key 1/\perp case.$

²⁶⁵ The two other cases are treated similarly (see appendix).

²⁶⁶ 4.4.2 Logical rules

We analyze the possible cases for a logical rule involved in an elementary PRIS. Note that in each case, the principal formula may be part of the left or right part of the splitting; we treat only one case each time since the other is symmetrical be taking the dual interpolant and exchanging π^L and π^R .

If the last rule is (3), i.e. if
$$\pi = \frac{ \begin{pmatrix} \pi_1^L \\ \Gamma, I \end{pmatrix} + I^{\perp}, \Delta, A, B}{ \begin{pmatrix} \Gamma, \Delta, A, B \\ \hline \Gamma, \Delta, A, B \end{pmatrix}}$$
 (Cut) then taking $I' = I, \pi^L = \pi_1^L$

and $\pi^R = \frac{\vdash I^{\perp}, \Delta, A, B}{\vdash I^{\perp}, \Delta, A \otimes B}$ (8) we obtain a solved PRIS π' such that $\pi \leftarrow_{\mathsf{cut}} \pi'$ by a commutative reduction of (Cut). 272

273

then setting $I = I_1 \otimes I_2, \pi^L = \frac{\vdash \prod_{1,I_1}^{\pi_1} \vdash \prod_{2,I_2}^{\pi_2}}{\vdash \prod_{1,\Gamma_2,I_1 \otimes I_2} \mid (\otimes)} \text{ and } \pi^R = \frac{\vdash I_1^{\perp}, \Delta_1, A \quad \vdash I_2^{\perp}, \Delta_2, B}{\vdash I_1^{\perp}, I_2^{\perp}, \Delta_1, \Delta_2, A \otimes B} \quad (\otimes)$ 275

one gets a solved PRIS π' such that $\pi \leftarrow _{\mathsf{cut}}^{\star} \pi'$ by a commutative reduction of (Cut) and 276 a key $(\otimes)/(\otimes)$ case. 277

If the last rule is (!p), that is
$$\pi = \frac{\vdash F, ?\Gamma, I_1 \quad \vdash I_1^{\perp}, ?\Delta}{\vdash F, ?\Gamma, ?\Delta}$$
 (Cut), then by setting $I = ?I_1$,

 $\pi^{L} = \frac{\vdash F, ?\Gamma, I_{1}}{\vdash F, ?\Gamma, ?I_{1}} \xrightarrow{(?d)} \text{and } \pi^{R} = \frac{\vdash I_{1}^{\perp}, ?\Delta}{\vdash !I_{1}^{\perp}, ?\Delta} \xrightarrow{(!p)}, \text{ one gets that } \pi \text{ cut-expands to } \pi':$

$$\pi' = \frac{\vdash !F, \stackrel{\pi}{?}\stackrel{L}{\Gamma}, ?I_1 \qquad \vdash !I_1^{\stackrel{\pi}{\perp}}, ?\Delta}{\vdash !F, ?\Gamma, ?\Delta} \quad (Cut) \qquad \longrightarrow_{cut} \qquad \frac{\vdash \stackrel{\pi}{F}, ?\Gamma, I_1}{\vdash F, ?\Gamma, I} \quad (?d) \qquad \frac{\vdash I_1^{\stackrel{\perp}{\perp}}, ?\Delta}{\vdash I^{\perp}, ?\Delta} \quad (!p) \qquad \longrightarrow_{cut} \pi.$$

²⁸¹ If the last rule is (?c), ie.
$$\pi = \frac{\vdash ?F, ?F, \Gamma, I_1 \qquad \vdash I_1^{\perp}, \Delta}{\vdash ?F, ?F, \Gamma, \Delta}$$
 (Cut), by setting $I = I_1, \pi^R =$

282 π_1^R and $\pi^L = \frac{\vdash ?F, ?F, \Gamma, I_1}{\vdash ?F, \Gamma, I}$ one gets that π cut-expands to the solved PRIS π' :

$$\pi' = \frac{\vdash \stackrel{?}{F} \stackrel{\pi}{\Gamma} \stackrel{\Gamma}{\Gamma}, I \qquad \vdash \stackrel{\pi}{I} \stackrel{R}{\Gamma}, \Delta}{\vdash \stackrel{?}{F} \stackrel{\Gamma}{\Gamma}, \Lambda} \quad (Cut) \qquad \longrightarrow_{cut} \qquad \frac{\vdash \stackrel{?}{F} \stackrel{R}{\Gamma} \stackrel{\Gamma}{\Gamma}, \Gamma, I_1 \qquad \vdash \stackrel{n}{I} \stackrel{1}{\Gamma}, \Delta}{\stackrel{\vdash \stackrel{?}{F} \stackrel{R}{\Gamma}, \Gamma, \Lambda} \quad (Cut) = \pi.$$

If the last rule is (a), i.e. $\pi = \frac{\vdash F\{y/x\}, \Gamma, I_1 \qquad \stackrel{\pi_1^K}{\vdash I_1^{\perp}, \Delta}}{\stackrel{\vdash F\{y/x\}, \Gamma, \Delta}{(\exists)}}$ (Cut) with $\operatorname{Voc}(I_1) \subseteq \operatorname{Voc}(F\{y/x\}, \Gamma) \cap$

 $\operatorname{Voc}(\Delta)$. In this case, note that we treat only the case of a FO language containing no 285 function symbols. We reason by case on whether y occurs in Γ, Δ : 286

²⁸⁷ = If y occurs in both, then we simply take
$$I = I_1$$
 as interpolant, $\pi^L = \frac{\pi_1^L}{\vdash \exists x F, \Gamma}$ (\exists) and
²⁸⁸ $\pi^R = \pi_1^R$. Since $\operatorname{Voc}(I) = \operatorname{Voc}(I_1) \subseteq \operatorname{Voc}(F, \Gamma) \cap \operatorname{Voc}(\Delta) = \operatorname{Voc}(\exists x F, \Gamma) \cap \operatorname{Voc}(\Delta)$, we

XX:10 Interpolation as cut-introduction

have that
$$\pi' = \frac{\vdash \exists x_F^{\perp}, \Gamma, I \qquad \vdash I^{\perp}, \Delta}{\vdash \exists x_F, \Gamma, \Delta}$$
 (Cut) is a solved PRIS to which π cut-expands

via a cut-commutation rule. 290

289

 $= \text{ If } y \text{ occurs in } \Gamma \text{ but not in } \Delta, \text{ then we set } I = \exists y I_1, \pi^L = \frac{\pi_1^L}{\vdash \exists x F, \Gamma, I_1} \stackrel{(\exists)}{\vdash} \frac{\exists x F, \Gamma, I_1}{\vdash \exists x F, \Gamma, \exists y I_1} \stackrel{(\exists)}{\vdash}$ and

$$\pi^{R} = \frac{\vdash I_{1}^{\perp}, \Delta}{\vdash \forall y I_{1}^{\perp}, \Delta} \quad \text{(v) we have that } \pi' = \frac{\vdash \exists x F, \Gamma, I \quad \vdash I^{\perp}, \Delta}{\vdash \exists x F, \Gamma, \Delta} \quad \text{(Cut) is a solved}$$

PRIS to which π cut-expands via a cut-commutation rule and a key $(\exists)/(\forall)$ rule:

= If y occurs in Δ but not in Γ , then we set $I = \forall y I_1, \pi^L = \frac{\vdash F\{y/x\}, \Gamma, I_1}{\vdash \exists x F, \Gamma, I_1}$ (\exists) and (\forall) 291

$$\pi^{R} = \frac{\overset{\pi^{R}_{1}}{\vdash \exists y I_{1}^{\perp}, \Delta}}{\vdash \exists y I_{1}^{\perp}, \Delta} \quad (\exists) \quad \text{One gets that } \pi' = \frac{\vdash \exists x F, \Gamma, \forall y I_{1} \quad \vdash \exists y I_{1}^{\perp}, \Delta}{\vdash \exists x F, \Gamma, \Delta} \quad (\mathsf{Cut}) \quad \text{is a}$$

solved PRIS to which π cut-expands:

$$\begin{array}{ccc} {}_{294} & & \pi' \longrightarrow_{\mathsf{cut}} \frac{\vdash F\{y/x\}, \Gamma, I_1}{\vdash \exists xF, \Gamma, I_1} & (\exists) & \pi_1^R \\ & & \vdash I_1^{\perp}, \Delta \\ & & \vdash \exists xF, \Gamma, \Delta \end{array} & (\mathsf{Cut}) \end{array} \longrightarrow_{\mathsf{cut}} \frac{\vdash F\{y/x\}, \Gamma, I_1 & \vdash I_1^{\perp}, \Delta \\ & & \frac{\vdash F\{y/x\}, \Gamma, \Delta }{\vdash \exists xF, \Gamma, \Delta} & (\exists) \end{array} & (\mathsf{Cut})$$

 \mathbf{P}

292

293

Other cases are treated similarly (See details in Appendix D.1). 296

4.5 Some consequences and extensions 297

In the same way as in the previous section, a similar development of LK-PRIS and LJ-PRIS 298 can be done, either from scratch or by means of linear embeddings which relates linear PRIS 299 to classicial or intuitionistic ones, this results in the following proposition: 300

▶ Proposition 13. Any initial LK-PRIS (resp. LJ-PRIS) can be reduced, by cut-expansions, 301 to a solved LK-PRIS (resp. solved LJ-PRIS). 302

▶ Remark 14. In fact, the above theorem does not rely on the cut-freeness of π , only the 303 notion of interpolant does: the cut-introduction performed in solving PRIS can indeed be 304 extended to handle the cut-case, as long as one relaxes the conditions on the language for 305 the interpolant. This suggests an immediate generalization of proof-relevant Craig and 306 Lyndon's interpolation for proofs with cuts in which each (pair of) cut-formula is assigned 307 to the left/right component of a splitting (in a consistent way for a given cut inference): 308 the language of the interpolant is then constructed by taking into account the language of 309 the cut-formulas depending on the choice of splitting. In that case, an additional degree of 310 freedom appears in the choice of the interpolant when assigning each cut to a component of 311

the splitting: strategies for optimizing the language of the interpolant could therefore be investigated.

Remark 15. Complementing on the previous remark, analytic cuts (that is cut formulas
 that are subformula of some formula of the conclusion sequent) are of particular interest here.
 Indeed, the above described procedure to work with PRIS containing cuts can be readily
 applied to proofs with analytic cuts.

In a similar way, it would be interesting to consider extending our procedure to proofs with cuts which are globally analytic (ie. simply requiring that every cut-formula is a subformula of some formula in the conclusion sequent). While the vocabulary constraint on interpolant *is locally violated* when encountering a globally-analytic cut, it is globally satisfied when reaching a solved PRIS. Other weakenings on the conditions on cut-formulas car be considered, such as the semi-analyticity conditions by Jalali and Tabatakai [39].

324 4.6 Extension with cuts

Even though the approach developed in this paper naturally fits in logical frameworks where 325 one has a sequent proof system satisfying a cut-elimination theorem, it may be relevant to 326 consider what is the impact of the cut inference on the process of synthesizing interpolants. 327 Indeed, the cut inference being the only one that breaks the subformula property, it also 328 prevents an immediate extension of Maehara's proof technique since the interpolants generated 329 via the premises may well violate the vocabulary constraints that we require for interpolation. 330 This is why interpolation is usually presented as a consequence of the cut-elimination 331 theorem. 332

On the other hand, it is not strictly necessary to consider cut-free proofs to perform 333 Maehara's interpolation nor our cut-introduction method. There may be several reasons to 334 consider cut formulas while interpolating. First, performing cut-elimination may be costly, 335 in both time and space: one shall first eliminate all the cuts and then work with the cut-free 336 proof which may be much larger than the original proofs. That is a good reason not to 337 perform more cut-reduction than required. On the other hand, another reason lies in the fact 338 from the computational viewpoint, performing interpolation from the cut-free proof consist 339 in first reducing a program to its most explicit form, where all intermediate computations 340 have been removed, before trying to factor computation. This goes seemingly in the wrong 341 direction as performing this computation may end up duplicating some code or data in the 342 process of unveiling the value of the computation. 343

In contrast, in many computational situations, interpolation can be performed in presence of cuts: think for instance of a function that computes a data combining booleans and natural numbers while receiving an input made of those data. When intermediate computations rely on those base type, one can hope, thanks to the method described below, to factor the computation through an interpolant type built from those two base types. This will require further investigation to be made precise.

let us now outline how the results in the previous subsections and specifically Definition 8 and Lemma 10 can be extended to proofs with cuts. Note that what follows essentially amounts to a standard trick consisting in viewing a cut inference on C, C^{\perp} as a \otimes introducing $C \otimes C^{\perp}$ (or the corresponding conjunction when working in classical or intuitionistic logic). The only subtelty is to find a workaround to maintain the adequate notion of vocabulary: we introduce the notion of cut vocabularies for that purpose. Note that in what follows, we adapt, without details, the previous paragraphs method to Craig interpolation as the polarity

XX:12 Interpolation as cut-introduction

constraints induced by Lyndon interpolation cannot be satisfied as such in our development
 when cut are present.

Let us first remark that the notion of coherent decorated proofs is straightforwardly extended to derivations containing the cut-inference by asking that on a cut inference, the coloring is transferred consistently from the conclusion sequent to the ancestors in the premises and that the cut-inferences is given the same, arbitrary, color. In the following, for simplicity, we assume that all cut inferences are assigned to the left component but this can be relaxed (in a consistent way)

Definition 16 (Proof-relevant Craig Interpolation Situations with Cuts). A CPRIS for (Γ, Δ) is the data of:

the goal, that is an LK decorated proof π of conclusion $\vdash \Gamma, \Delta$ and with $n \geq 0$ open premises $(\vdash \Gamma_i, \Delta_i)_{1 \leq i \leq n}$ such that for each $1 \leq i \leq n$ the formulas in Γ_i (resp. Δ_i) are ancestors of formulas in Γ (resp. of Δ) or ancestors of a cut-formula;

The cut vocabularies, that is for each $1 \leq i \leq n$, a pair $(\mathbb{C}_i^L, \mathbb{C}_i^R)$ of sets of literals;

The partial interpolants, that is for each $1 \leq i \leq n$, a formula I_i st. $\operatorname{Voc}(I_i) \subseteq (\operatorname{Voc}(\Gamma_i) \cup \mathbb{C}_i^L) \cap (\operatorname{Voc}(\Delta_i) \cup \mathbb{C}_i^R)$ and;

The partial solutions, that is, for each $1 \leq i \leq n$, derivations π_i^L (resp. π_i^R) of conclusion $\vdash \Gamma_i, I_i$ (resp. $\vdash I_i^{\perp}, \Delta_i$) such that if \mathcal{C}^L (resp. \mathcal{C}^R) is the set of cut-formulas in π_i^L (resp. in π_i^R), then $\bigcup_{C \in \mathcal{C}^\epsilon} \mathbb{Voc}(C) = \mathbb{C}_i^\epsilon, \epsilon \in \{L, R\}.$

With the above definition, the notions of initial, solved, and elementary CPRIS is adapted trivially and one notices that a solved CPRIS when the cut-formulas are all (locally or globally) analytic is a solution to the interpolation problem if cut formulas are colored consistently with the conclusion formulas they are a subformula of.

Since all invariants of the proof of Lemma 10 hold for CPRIS, what remains to show is that the proof of Lemma 10 can be extended to the case of elementary CPRIS concluded with a cut-inference. It can easily be seen that such an extension directly follows for the case of the \otimes rule, the only difference is that instead of being part of the vocabulary of the context formulas, the cut-formulas contribute to the cut vocabularies.

If the last rule is cut, i.e. if
$$\pi = \frac{\vdash \prod_{1}^{\pi_{1}^{L}} I_{1} \qquad \vdash I_{1}^{\perp}, \Delta_{1}, C}{\vdash \prod_{1}, \Delta_{1}, C}$$
 (Cut) $\frac{\vdash \prod_{2}^{L}, I_{2} \qquad \vdash I_{2}^{\perp}, \Delta_{2}, C^{\perp}}{\vdash \prod_{2}, \Delta_{2}, C^{\perp}}$ (Cut),
 $\frac{\vdash \prod_{1}, \Delta_{1}, C}{\vdash \prod_{1}, \Gamma_{2}, \Delta_{1}, \Delta_{2}}$ (Cut) (Cut),
 $\frac{\vdash \prod_{1}^{L}, \Delta_{1}, C}{\vdash \prod_{1}^{L}, \Gamma_{2}, \Delta_{1}, \Delta_{2}}$ (Cut) (Cut)

then setting
$$I = I_1 \otimes I_2, \pi^L = \frac{\prod_{1=1}^{\pi_1^L} I_1 + \prod_{2=2}^{\pi_2^L} I_2}{\prod_{1=1}^{-1} \prod_{1=1}^{-1} \prod_{1=2}^{-1} I_1 \otimes I_2} \otimes \text{ and } \pi^R = \frac{\prod_{1=1}^{-1} I_1^\perp, \Delta_1, C + \prod_{2=1}^{-1} \Delta_2, C^\perp}{\prod_{1=1}^{-1} \prod_{1=2}^{-1} I_1 \otimes I_2} \otimes (Cut)$$

one gets a solved CPRIS π' such that $\pi \leftarrow_{cut}^* \pi'$ by a commutative reduction of (Cut) and a key (\otimes)/(\otimes) case.

$$\text{if } \pi = \frac{\vdash \prod_{1,C,I_{1}}^{\pi_{1}^{L}} \subset \prod_{1}^{\pi_{1}^{R}} \cap \prod_{1}^{\pi_{1}^{R}} \cap \prod_{1}^{\pi_{1}^{R}} \cap \prod_{1}^{\pi_{1}^{R}} \cap \prod_{1}^{\pi_{1}^{R}} \cap \prod_{1}^{\pi_{1}^{L}} \cap \prod$$

391

$$I_1 \otimes I_2, \pi^L = \frac{\vdash \Gamma_1, C, I_1 \quad \vdash \Gamma, C^{\perp}, I_2}{\underset{\vdash \Delta_1, \Delta_2, I_1 \otimes I_2}{\vdash \Delta_1, \Delta_2, I_1 \otimes I_2}} (\text{Cut}) \text{ and } \pi^R = \frac{\underset{I_1^{\perp}, \Delta_1}{\# I_1^{\perp}, \Delta_1} \quad \underset{I_2^{\perp}, \Delta_2}{\# I_2^{\perp}, \Delta_1, \Delta}}{\underset{\vdash (I_1 \otimes I_2)^{\perp}, \Delta_1, \Delta}} (\otimes) \text{ one }$$

gets a solved CPRIS π' such that $\pi \leftarrow \overset{*}{\operatorname{cut}} \pi'$ by a commutative reduction of (Cut) and a key (\otimes)/(\otimes) case.

³⁹⁴ 5 On the computational significance of the result: Interpolating ³⁹⁵ System L

т

Stated purely in terms of interpolating validity judgements, interpolation is not very meaningful computationally of course. On the other hand, the refined statement investigated in the present paper opens new perspectives: the interpolant formula can be viewed as a an interface type I through which a computation can from data type A to data type B be factored via type I. Let us focus on intuitionistic statement (say, in ILL or LJ):

If π proves $\Gamma, \Delta \vdash C$, there exists some formula I such that $\operatorname{Voc}(I) \subseteq \operatorname{Voc}(\Gamma) \cap \operatorname{Voc}(\Delta, C)$ and proofs π_1, π_2 of $\Gamma \vdash I$ and $I, \Delta \vdash C$ respectively, such that $(\operatorname{Cut})(\pi_1, \pi_2) \longrightarrow_{\operatorname{cut}}^{\star} \pi$.

Rephrased in terms of the λ -calculus, that would mean in particular that: for any closed term $\lambda x.t : A \to B$ there is a type C such that $\operatorname{Voc}(C) \subseteq \operatorname{Voc}(A) \cap \operatorname{Voc}(B)$ and terms $u : A \to C$ and $v : C \to B$ such that $\lambda x.t =_{\beta} \lambda x.(v(ux))$. This is the content of the analysis by Čubrić. Data type C can therefore be viewed as an interface between types A and Bwhile processing computation t, involving only pieces of data involved in both A and B.

Of course, this interface depends in t itself in the above proof, which would be different if 408 uniform interpolation is lifted to this proof-relevant framework. Indeed, in that case, the 409 data type for the interpolant only depend on A and the common vocabulary with the output 410 type. In that way, interpolation would express some generic preprocessing from A to the 411 interpolant type $UI(A, \mathcal{V})$ depending only on formula A and type variables \mathcal{V} , from which 412 any computation to a type B (sharing vocabulary \mathcal{V} with A) can be performed. In the same 413 spirit, more precise results on interpolation (especially on polarity of occurrences of relational 414 symbols, etc.) would provide more information on data usage. 415

We shall now outline a rephrasing of our previous results in a term calculus which corresponds to sequent calculus as the λ -calculus does to natural deduction. For uniformity, this version is essentially inspired from the calculus designed by Munch-Maccagnoni [29, 30], presented in a two-sided manner rather than one-sided to emphasize the input/output behaviour and term/context duality, that we restrict to the purely linear (ie multiplicative and additive setting). Some details are omitted below: see Appendix E for more details.

422 ▶ Definition 17 (Linear System L). As usual, the grammar and type system of linear system
 423 L consists of three syntactic categories: commands, terms and contexts, defined by mutual
 424 recursion:

 $\begin{array}{ll} {}_{425} & = c ::= \langle t \mid e \rangle \\ {}_{426} & = t, u ::= x \mid \mu \alpha. c \mid (s,t) \mid \lambda x.t \mid \mu[\alpha,\beta].c \mid () \mid \mu[].c \mid \iota_1(t) \mid \iota_2(t) \mid \mu(\pi_1(\alpha) \mapsto c \mid \pi_2(\beta) \mapsto d) \mid tp \\ {}_{427} & d) \mid tp \\ {}_{428} & = e, f ::= \alpha \mid \tilde{\mu} x.c \mid \tilde{\mu}(x,y).c \mid t \cdot e \mid [e,f] \mid \tilde{\mu}().c \mid [] \mid \pi_1(e) \mid \pi_2(e) \mid \tilde{\mu}(\iota_1(x) \mapsto c \mid \iota_2(x) \mapsto d) \mid stop \end{array}$

⁴³⁰ This gives rise to three kinds of typing judgments: $c : (\Gamma \vdash \Delta); \quad \Gamma \vdash t : A \mid \Delta; \quad \Gamma \mid e : A \vdash \Delta$ ⁴³¹ and the typing rules given in Figure 3.

Identity rules					
$\overline{x:A \vdash x:A \mid \emptyset}$ Axt	$\frac{c:(\Gamma \vdash \alpha: A, \Delta)}{\Gamma \vdash \mu \alpha. c: A \mid \Delta} \mu$	Term formation			
$\frac{1}{\emptyset \mid \alpha: A \vdash \alpha: A} \text{ Axc}$	$\frac{c:(\Gamma,x:A\vdash \Delta)}{\Gamma\mid \tilde{\mu}x.c:A\vdash \Delta} \ \tilde{\mu}$	Context formation			
$\frac{\Gamma \vdash t: A \mid \Delta \Gamma \mid e: A}{\langle t \mid e \rangle: (\Gamma, \Gamma' \vdash \Delta, \Delta}$	$\frac{\vdash \Delta'}{\prime)}$ cut	Command formation			
Multiplicative rules					
$\frac{\Gamma \vdash s: A \mid \Delta \Gamma' \vdash t: I}{\Gamma, \Gamma' \vdash (s, t): A \otimes B \mid \Delta}$	$\frac{B \mid \Delta'}{\Delta, \Delta'} \otimes^{r} \frac{\Gamma, x : A \vdash t : B \mid \Delta}{\Gamma \vdash \lambda x.t : A \multimap B \mid \Delta} \multimap^{r}$	Term formation			
$\frac{c:(\Gamma\vdash\alpha:A,\beta:B,\Delta)}{\Gamma\vdash\mu[\alpha,\beta].c:A{\mathord{ \otimes } } B\mid\Delta}$	$ \mathfrak{S}^r \overline{\emptyset \vdash (): 1 \mid \emptyset} \ 1^r \qquad \frac{c: (\Gamma \vdash \Delta)}{\Gamma \vdash \mu[].c: \bot \mid \Delta} \ \bot $	T			
$\frac{c:(\Gamma, x: A, y: B \vdash \Delta)}{\Gamma \mid \tilde{\mu}(x, y).c: A \otimes B \vdash \Delta}$	$\Gamma \otimes^{l} = rac{\Gamma \vdash t : A \mid \Delta \Gamma' \mid e : B \vdash \Delta'}{\Gamma, \Gamma' \mid t \cdot e : A \multimap B \vdash \Delta, \Delta'} \multimap^{l}$	Context formation			
$\frac{\Gamma \mid e: A \vdash \Delta \Gamma' \mid f: B}{\Gamma, \Gamma' \mid [e, f]: A \otimes B \vdash \Delta}$	$\frac{B\vdash\Delta'}{\Delta,\Delta'} \otimes^l \frac{c:(\Gamma\vdash\Delta)}{\Gamma\mid \tilde{\mu}().c:1\vdash\Delta} 1^l \overline{\emptyset\mid []:\bot\vdash}$	$\overline{\emptyset} \perp^l$			
Additive rules					
$\frac{\Gamma \vdash t : A \mid \Delta}{\Gamma \vdash \iota_1(t) : A \oplus B \mid \Delta} \oplus_1^t$	$\Gamma \qquad \frac{\Gamma \vdash t : B \mid \Delta}{\Gamma \vdash \iota_2(t) : A \oplus B \mid \Delta} \oplus_2^r$	Term formation			
$\frac{c:(\Gamma\vdash\alpha:A,\Delta) d}{\Gamma\vdash\mu(\pi_1(\alpha)\mapsto c\mid\pi_2(\beta))}$	$ \frac{:(\Gamma \vdash \beta : B, \Delta)}{\mapsto d) : A \otimes B \mid \Delta} \otimes^{r} \qquad \overline{\Gamma \vdash \mathtt{tp} : \top \mid \Delta} \ ^{\top r} $				
$\frac{\Gamma \mid e : A \vdash \Delta}{\Gamma \mid \pi_1(e) : A \otimes B \vdash \Delta} \otimes_1^l$	$\frac{\Gamma \mid e : B \vdash \Delta}{\Gamma \mid \pi_2(e) : A \otimes B \vdash \Delta} \otimes_2^l$	Context formation			
$\frac{c:(\Gamma, x: A \vdash \Delta) d:(\Gamma', x: B \vdash \Delta')}{\Gamma \mid \tilde{\mu}(\iota_1(x) \mapsto c \mid \iota_2(y) \mapsto d): A \oplus B \vdash \Delta} \oplus^l \overline{\Gamma \mid \mathtt{stop}: 0 \vdash \Delta} 0^l$					
Figure 3 Type derivati	ion rules for System L				

▶ Remark 18. Notice that we use very similar notation for term and context constructs $(\iota/\pi, (u, v)/[e, f], ()/[], tp/stop)$, in order to emphasize the deep symmetry of the System L framework we work with. This symmetry is exploited in the proof of our main theorem.

The calculus could actually be presented with one-sided typing judgments, see [29].

We provide some of the main reduction rules of this calculus, keeping in mind the invariant
 that one reduces *commands*:

 $\begin{aligned} & 438 \qquad = \langle \mu \alpha.c \mid e \rangle \longrightarrow_{\mu} c\{e/\alpha\} \\ & 439 \qquad = \langle t \mid \tilde{\mu}x.c \rangle \longrightarrow_{\tilde{\mu}} c\{t/x\} \\ & 440 \qquad = \langle \lambda x.t \mid u \cdot e \rangle \longrightarrow_{\lambda} \langle \tilde{\mu}x.\langle t \mid e \rangle \mid u \rangle \\ & 441 \qquad = \langle (t,u) \mid \tilde{\mu}(x,y).c \rangle \longrightarrow_{\otimes} c\{t/x,u/y\} \\ & 442 \qquad = \langle \mu[\alpha,\beta].c \mid [e,f] \rangle \longrightarrow_{\otimes} c\{e/\alpha,f/\beta\} \\ & 443 \qquad = \langle \iota_{j}(t) \mid \tilde{\mu}(\iota_{1}(x_{1}) \mapsto c_{1} \mid \iota_{2}(x_{2}) \mapsto c_{2}) \rangle \longrightarrow_{\oplus} c_{j}\{t/x_{j}\} \\ & 444 \qquad = \langle \mu(\pi_{1}(\alpha_{1}) \mapsto c_{1} \mid \pi_{2}(\alpha_{2}) \mapsto c_{2}) \mid \pi_{j}(e) \rangle \longrightarrow_{\&} c_{j}\{e/\alpha_{j}\} \end{aligned}$

435

445 = $\langle () | \tilde{\mu}().c \rangle \longrightarrow_1 c$ 446 = $\langle \mu [].c | [] \rangle \longrightarrow_\perp c$

Moreover, one shall consider in the development below the η rules for μ and $\tilde{\mu}$ (which respectively reduce terms and contexts):

The above rules satisfy the expected subject reduction property. Moreover, we need to reflect the splitting of sequents which will lead us to define a system L with a split typing system to be detailed now.

▶ Definition 19 (System L with split typing judgements). We introduce split typing judgements where types are enriched with a splitting information in the form of a label $l \in \{L, R\}$ carried by each type occurring in the judgment, which are then of one of the following shape:

460 with $k, l, k_1, \ldots, k_m, l_1, \ldots, l_n \in \{L, R\}.$

⁴⁶¹ By convention, to save space and ease the reading of judgements, we may use colors to ⁴⁶² represent the labels: a labelled type T^L (resp. U^R) may be written T (resp. U), shorthand ⁴⁶³ which will be extended to whole typing contexts sharing the same label: Γ or Γ .

⁴⁶⁴ ► Theorem 20. In what follows, t (resp. e, resp. c) is a normal L-term (resp. normal
 ⁴⁶⁵ L-context, resp. normal L-command). The following interpolating results hold:

1. If $c : (\Gamma_1, \Gamma_2 \vdash \Delta_1, \Delta_2)$, there exist a type $I \in \mathbb{V}oc(\Gamma_1, \Delta_1) \cap \mathbb{V}oc(\Gamma_2, \Delta_2)$ and t, e such that $\Gamma_1 \vdash t : I \mid \Delta_1$ and $\Gamma_2 \mid e : I \vdash \Delta_2$, and $\langle t \mid e \rangle \longrightarrow^* c$.

2. If $\Gamma_1, \Gamma_2 \vdash t : A \mid \Delta_1, \Delta_2$, there exist a type $I \in \mathbb{V}oc(\Gamma_1, \Delta_1, A) \cap \mathbb{V}oc(\Gamma_2, \Delta_2)$ and α, t', e' such that $\Gamma_1 \vdash t' : A \mid \alpha : I, \Delta_1$ and $\Gamma_2 \mid e' : I \vdash \Delta_2$, and $t'\{e'/\alpha\} \longrightarrow^* t$.

3. If $\Gamma_1, \Gamma_2 \mid e : A \vdash \Delta_1, \Delta_2$, there exist a type $I \in \mathbb{V}oc(\Gamma_1, \Delta_1, A) \cap \mathbb{V}oc(\Gamma_2, \Delta_2)$ and α, e', e'' such that $\Gamma_1 \mid e' : A \mid \alpha : I, \Delta_1$ and $\Gamma_2 \mid e'' : I \vdash \Delta_2$, and $e'\{e''/\alpha\} \longrightarrow^* e$.

472 **4.** If $\Gamma_1, \Gamma_2 \vdash t : A \mid \Delta_1, \Delta_2$, there exist a type $I \in \mathbb{V}oc(\Gamma_1, \Delta_1) \cap \mathbb{V}oc(\Gamma_2, \Delta_2, A)$ and α, t', t'' 473 such that $\Gamma_1 \vdash t'' : I \mid \Delta_1$ and $\Gamma_2, x : I \vdash t' : A \mid \Delta_2$, and $t'\{t''/x\} \longrightarrow^* t$.

474 **5.** If $\Gamma_1, \Gamma_2 \mid e : A \vdash \Delta_1, \Delta_2$, there exist a type $I \in \mathbb{V}oc(\Gamma_1, \Delta_1) \cap \mathbb{V}oc(\Gamma_2, \Delta_2, A)$ and x, t', e'475 such that $\Gamma_1 \vdash t' : I \mid \Delta_1$ and $\Gamma_2, x : I \mid e' : A \vdash \Delta_2$, and $e'\{t'/x\} \longrightarrow^* e$.

476 Proof sketch (Details are provided in Appendix E). The result is proved by mutual induc477 tion on the structure of terms, contexts and commands. We treat only cases 1–3, cases 4
478 and 5 being essentially similar to cases 3 and 2 respectively.

⁴⁷⁹ **Case 1.** If c is a command in normal form, it is either of the form $\langle z | e \rangle$ (with z being ⁴⁸⁰ declared in Γ_1 or Γ_2) or $\langle t | \beta \rangle$ (with α being declared in Δ_1 or Δ_2). Depending on the case ⁴⁸¹ and whether the variable is declared in the left or right component of the typing contexts, ⁴⁸² we apply induction hypotheses for one of cases 2–4.

Let us assume for instance that $c = \langle z \mid e \rangle$ and $\langle z \mid e \rangle : (\Gamma_1, \Gamma_2, z : A \vdash \Delta_1, \Delta_2)$. Then, *e* being structurally smaller than *c*, the induction hypothesis applied on $\Gamma_1, \Gamma_2 \mid e : A \vdash \Delta_1, \Delta_2$, ensures the existence of a type $I \in \mathbb{V}oc(\Gamma_1, \Delta_1) \cap \mathbb{V}oc(\Gamma_2, \Delta_2, A)$ as well as x, t', e' such that $\Gamma_1 \vdash t' : I \mid \Delta_1$ and $\Gamma_2, x : I \mid e' : A \vdash \Delta_2$, and $e'\{t'/x\} \longrightarrow^* e$.

XX:16 Interpolation as cut-introduction

⁴⁸⁷ Therefore $\langle t' | \tilde{\mu}x.\langle z | e' \rangle \rightarrow^{\star} c$ and we indeed have $\Gamma_1 \vdash t' : I \mid \Delta_1$ and $\Gamma_2, z : A \mid$ ⁴⁸⁸ $\tilde{\mu}x.\langle z \mid e' \rangle : I \vdash \Delta_2$. The other cases are similar.

489 **Case 2.** In that case, we make a case distinction based on the structure of t.

⁴⁹⁰ If $t = \mu \alpha.c$ and its typing judgment has shape $\Gamma_1, \Gamma_2 \vdash \mu \alpha.c : A \mid \Delta_1, \Delta_2$; It follows that c⁴⁹¹ has typing judgment $\Gamma_1, \Gamma_2 \vdash \alpha : A, \Delta_1, \Delta_2$ and therefore by induction hypothesis, there ⁴⁹² exist an interpolant type I, a term t and a context e such that $\Gamma_1 \vdash t : I \mid \alpha : A, \Delta_1$ and ⁴⁹³ $\Gamma_2 \mid e : I \vdash \Delta_2$ such that $\langle t \mid e \rangle \longrightarrow^* c$. Let us then set $I' = I, t' = \mu \alpha. \langle t \mid \beta \rangle$ and e' = e⁴⁹⁴ and one straightforwardly gets that $t' \{ e' / \beta \} \longrightarrow^* \mu \alpha.c$.

If $t = \lambda x.u$ and its typing judgment has shape $\Gamma_1, \Gamma_2 \vdash \lambda x.u : A \multimap B \mid \Delta_1, \Delta_2$.

⁴⁹⁶ By induction hypothesis we find an interpolant type I for u which can be used to ⁴⁹⁷ interpolate t as well: we have α, t', e' such that $\Gamma_1, x : A \vdash t' : B \mid \alpha : I, \Delta_1$ and ⁴⁹⁸ $\Gamma_2 \mid e' : I \vdash \Delta_2$, and $t'\{e'/\alpha\} \longrightarrow^* u$. Therefore $\lambda x.t'\{e'/\alpha\} \longrightarrow^* \lambda x.u = t$.

⁴⁹⁹ If t = (u, v) and its typing judgment has shape $\Gamma_1, \Gamma'_1, \Gamma_2, \Gamma'_2 \vdash (u, v) : A \otimes B \mid \Delta_1, \Delta'_1, \Delta_2, \Delta'_2$ ⁵⁰⁰ with $\Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2$ and $\Gamma'_1, \Gamma'_2 \vdash v : B \mid \Delta'_1, \Delta'_2$

By induction hypothesis we find an interpolant type J for u and K for v that we can combine to interpolate t as well: we have α, u', e' such that $\Gamma_1 \vdash u' : A \mid \alpha : J, \Delta_1$ and $\Gamma_2 \mid e' : J \vdash \Delta_2$, and $u'\{e'/\alpha\} \longrightarrow^* u$ and we have β, v', f' such that $\Gamma'_1 \vdash v' : B \mid \beta : K, \Delta'_1$ and $\Gamma'_2 \mid f' : K \vdash \Delta'_2$, and $v'\{f'/\alpha\} \longrightarrow^* v$. Therefore we set $t' = \mu\gamma.\langle\mu[\alpha, \beta].\langle(u', v') \mid \gamma\rangle \mid \delta\rangle$ and we have $t'\{[e', f'])/\delta\} = \mu\gamma.\langle\mu[\alpha, \beta].\langle(u', v') \mid \gamma\rangle \mid \delta\rangle$ $\delta\rangle\{[e', f'])/\delta\} \longrightarrow^* \mu\gamma.\langle(u'\{e'/\alpha\}, v'\{f'/\beta\}) \mid \gamma\rangle \longrightarrow^* \mu\gamma.\langle(u, v) \mid \gamma\rangle \longrightarrow \eta_{\mu} (u, v).$

⁵⁰⁸ Case 3. The context case is symmetrical to the previous one, due to the symmetry of linear ⁵⁰⁹ typing, but for the case of the applicative context which is detailed in appendix.

510 ► Remark 21. The above results could be extended to full LL, even though it would be more adequate formally to move to the original calculus by Curien and Herbelin and prove a corresponding result to proof-relevant interpolation for LK to avoid complexities in dealing with structural rules.

In Curien-Herbelin's setting, a particular attention should be paid to the reduction rules that are needed in order to investigate to what account one can restrict to specific evaluation strategies, such as call-by-name and call-by-value, in interpolating, in order to avoid the computational critical-pair. Note that in the above development, this critical pair is not problematic due to the linearity discipline that is enforced: the sequent calculus non confluence does not induce a trivial equational theory (as is known from proof nets for instance which precisely perform this quotient).

521 6 Conclusion

In this paper, we established a refined, proof-relevant, version of Craig-Lyndon interpolation 522 theorems for first-order linear logic and then deduced it, using completely standard tools of LL 523 proof theory, to LK and LJ. A most striking fact, in our opinion, is that the result was *almost* 524 there for decades, since the early proofs by Maehara (and its broad dissemination in proof 525 theory textbooks, not to speak of applications to broader logical frameworks) and Prawitz. 526 Borrowing Feferman's words, "though deceptively simple and plausible on the face of it", we 527 think that this approach to proof-relevant interpolation in sequent calculus emphasizes a deep 528 duality between interpolation and cut elimination : more specifically, the process of synthesis 529 of the interpolant and the two interpolating proofs is reformulated as a cut-introduction 530

⁵³¹ process. Finally, we considered the computational content of the results by an analysis of the ⁵³² results in system L.

While we think that interpolation as cut-introduction is both a new conceptual and 533 technical contribution of this work, a proof-relevant interpolation theorem has already been 534 established by Cubrić [9, 10] in the early 90's for propositional intuitionistic natural deduction 535 in the form of an interpolation for the typed λ -calculus and for bicartesian closed categories. 536 Our approach is similarly subject to a computational interpretation that we plan to develop 537 in a future work about interpolation in system L [11, 12]. In fact we also hope that our 538 interpolation-as-cut-introduction can pave the way for a broader analysis of the computational 539 content of interpolation as a manner to factor computation through interfacing (that is, 540 interpolating) types. Indeed, while the computational interpretation of Cubrić's result, stated 541 in the λ -calculus, is certainly more transparent than the sequent calculus that we presented 542 here, it has not been extended in more than 30 years, except once by Matthes [26]. A reason 543 for this might be that while both allows for a proof-relevant phrasing, Maehara's method 544 is more modular and easily extensible than Prawitz as it rests on a logical framework, the 545 sequent calculus, that is inherently more modular that natural deduction. For instance, we 546 conjecture that it is possible to state a computational version of interpolation in classical 547 System L, that is in a classical framework featuring continuations, while it is not clear how 548 Čubrić' results can be extended beyond simply typed λ -calculus, eg. to Parigot's $\lambda\mu$ -calculus. 549

Among other future works that we plan to tackle, one can list the following directions.

Extend our treatment of System L to full LL as well as LK (with a focus on evaluation strategies).

Extend proof-relevant interpolation to provide a treatment to circular proofs for LL with
 least and greatest fixed-points.

 In LL, proof nets are a proof system that satisfies canonicity properties akin to natural 555 deduction for intuitionistic logic. It seems that interpolation as cut-introduction in 556 (multiplicative) proof-nets can be reformulated in terms of the parsing correctness criterion. 557 - We hope to establish factorization properties for models of LL, similar to Čubrić's results. 558 An intrinsic advantage of sequent calculus over natural deduction, and therefore of our 559 approach over Čubrić's, is that many more logics can be formulated as sequent calculi than 560 in natural deduction. Can we extend our results to other logics having cut-elimination? 561 - An important question and clearly non-trivial question that we would like to explore 562 is whether such a proof-relevant approach to interpolation can be extended to uniform 563 interpolation. That would mean that all computations that can be performed from 564 a piece of data u in a type A to data sharing with A only a fixed set \mathcal{L} of primitive 565 datatypes can be factored through a program that computes a value v in the uniform 566 interpolant datatype build from \mathcal{L} such that everything that can be computed from u567 can be computed from v as well. 568

⁵⁶⁹ Interpolating proofs containing cuts can actually be useful as shown recently by Hetzl and Jalali [22]. We plan to follow the directions outlined in Remarks 14 and 15.

⁵⁷¹ = Finally, we plan to reconsider Čubrić's results from the cut-introduction perspective.

572 — References

Bahareh Afshari and Graham E. Leigh. Lyndon interpolation for modal μ-calculus. In Aybüke
 Özgün and Yulia Zinova, editors, Language, Logic, and Computation - 13th International
 Tbilisi Symposium, TbiLLC 2019, Batumi, Georgia, September 16-20, 2019, Revised Selected

XX:18 Interpolation as cut-introduction

- Papers, volume 13206 of Lecture Notes in Computer Science, pages 197–213. Springer, 2019.
- 577 doi:10.1007/978-3-030-98479-3_10.
- Michele Basaldella and Kazushige Terui. Infinitary completeness in ludics. In Proceedings of the 25th Annual IEEE Symposium on Logic in Computer Science, LICS 2010, 11-14 July 2010, Edinburgh, United Kingdom, pages 294–303. IEEE Computer Society, 2010. doi: 10.1109/LICS.2010.47.
- Michele Basaldella and Kazushige Terui. On the meaning of logical completeness. Log. Methods
 Comput. Sci., 6(4), 2010. doi:10.2168/LMCS-6(4:11)2010.
- Evert W Beth. On padoa's method in the theory of definition. Indagationes Mathematicae,
 15:330–339, 1953.
- James Brotherston and Rajeev Goré. Craig interpolation in displayable logics. In *TABLEAUX*, volume 6793 of *Lecture Notes in Computer Science*, pages 88–103. Springer, 2011.
- Alessandra Carbone. Interpolants, cut elimination and flow graphs for the propositional calculus. Ann. Pure Appl. Log., 83(3):249-299, 1997. doi:10.1016/S0168-0072(96)00019-X.
- William Craig. Linear reasoning. A new form of the Herbrand-Gentzen theorem. J. Symb.
 Log., 22(3):250-268, 1957. doi:10.2307/2963593.
- William Craig. Three uses of the herbrand-gentzen theorem in relating model theory and
 proof theory. J. Symb. Log., 22(3):269-285, 1957. doi:10.2307/2963594.
- ⁵⁹⁴ **9** Djordje Cubric. *Results in categorical proof theory*. PhD thesis, McGill University, 1993.
- ⁵⁹⁵ 10 Djordje Cubric. Interpolation property for bicartesian closed categories. Arch. Math. Log.,
 ⁵⁹⁶ 33(4):291-319, 1994. doi:10.1007/BF01270628.
- Pierre-Louis Curien and Hugo Herbelin. The duality of computation. In Martin Odersky and
 Philip Wadler, editors, Proceedings of the Fifth ACM SIGPLAN International Conference
 on Functional Programming (ICFP '00), Montreal, Canada, September 18-21, 2000, pages
 233-243. ACM, 2000. doi:10.1145/351240.351262.
- Pierre-Louis Curien and Guillaume Munch-Maccagnoni. The duality of computation under focus. In Cristian S. Calude and Vladimiro Sassone, editors, *Theoretical Computer Science* 603 - 6th IFIP TC 1/WG 2.2 International Conference, TCS 2010, Held as Part of WCC 2010, Brisbane, Australia, September 20-23, 2010. Proceedings, volume 323 of IFIP Advances in Information and Communication Technology, pages 165–181. Springer, 2010. doi:10.1007/ 978-3-642-15240-5_13.
- Vincent Danos, Jean-Baptiste Joinet, and Harold Schellinx. A new deconstructive logic: Linear
 logic. J. Symb. Log., 62(3):755-807, 1997.
- Jeremy E. Dawson, James Brotherston, and Rajeev Goré. Machine-checked interpolation
 theorems for substructural logics using display calculi. In *IJCAR*, volume 9706 of *Lecture Notes in Computer Science*, pages 452–468. Springer, 2016.
- Adrien Deloro. Cours de logique mathématique. Draft of a book on mathematical logic,
 2023. URL: https://webusers.imj-prg.fr/~adrien.deloro/index.php?topic=teaching&
 page=logique.
- 16 Kosta Došen. Logical Consequence: A Turn in Style, pages 289–311. Springer Netherlands,
 Dordrecht, 1997. doi:10.1007/978-94-017-0487-8_15.
- Solomon Feferman. Harmonious logic: Craig's interpolation theorem and its descendants.
 Synthèse, 164(3):341–357, 2008. URL: https://doi.org/10.1007/s11229-008-9354-2, doi:
 10.1007/S11229-008-9354-2.
- Melvin Fitting. Linear reasoning in modal logic. J. Symb. Log., 49(4):1363–1378, 1984.
 doi:10.2307/2274285.
- I9 Jean-Yves Girard. Linear logic. Theoretical Computer Science, 50(1):1 101, 1987. doi:
 10.1016/0304-3975(87)90045-4.
- ⁶²⁴ 20 Jean-Yves Girard. Proof theory and logical complexity. Bibliopolis, 1987.
- Thomas A. Henzinger, Ranjit Jhala, Rupak Majumdar, and Kenneth L. McMillan. Abstractions
 from proofs. SIGPLAN Not., 39(1):232–244, jan 2004. doi:10.1145/982962.964021.

- Stefan Hetzl and Raheleh Jalali. On the completeness of interpolation algorithms. In Pawel
 Sobocinski, Ugo Dal Lago, and Javier Esparza, editors, Proceedings of the 39th Annual
 ACM/IEEE Symposium on Logic in Computer Science, LICS 2024, Tallinn, Estonia, July
 8-11, 2024, pages 45:1-45:13. ACM, 2024. doi:10.1145/3661814.3662112.
- Makoto Kanazawa. Computing interpolants in implicational logics. Annals of Pure and Applied
 Logic, 142(1):125-201, 2006. URL: https://www.sciencedirect.com/science/article/pii/
 S0168007205001922, doi:10.1016/j.apal.2005.12.014.
- Roman Kuznets. Multicomponent proof-theoretic method for proving interpolation properties.
 Ann. Pure Appl. Log., 169(12):1369–1418, 2018. URL: https://doi.org/10.1016/j.apal.
 2018.08.007, doi:10.1016/J.APAL.2018.08.007.
- ⁶³⁷ **25** Shoji Maehara. On the interpolation theorem of Craig. Sûgaku, 12(4), 1960.
- Ralph Matthes. Interpolation for natural deduction with generalized eliminations. In Reinhard
 Kahle, Peter Schroeder-Heister, and Robert Stärk, editors, *Proof Theory in Computer Science*,
 pages 153–169, Berlin, Heidelberg, 2001. Springer Berlin Heidelberg.
- Kenneth L. McMillan. Interpolation and model checking. In *Handbook of Model Checking.*,
 pages 421–446. 2018. doi:10.1007/978-3-319-10575-8_14.
- ⁶⁴³ 28 Grigori Mints. Finite investigations of transfinite derivations. Journal of Soviet Mathematics, 10:548–596, 1978.
- Guillaume Munch-Maccagnoni. Focalisation and Classical Realisability. In Erich Grädel
 and Reinhard Kahle, editors, *Computer Science Logic '09*, volume 5771 of *Lecture Notes in Computer Science*, pages 409–423. Springer, Heidelberg, 2009.
- Guillaume Munch-Maccagnoni. Syntax and Models of a non-Associative Composition of Programs and Proofs. PhD thesis, Univ. Paris Diderot, 2013.
- ⁶⁵⁰ 31 Dag Prawitz. Natural Deduction: A Proof-Theoretical Study. Dover Publications, Mineola,
 ⁶⁵¹ N.Y., 1965.
- João Rasga, Walter Alexandre Carnielli, and Cristina Sernadas. Interpolation via translations.
 Math. Log. Q., 55(5):515-534, 2009.
- A. Robinson. A result of consistency and its application to the theory of definition. Indagationes
 Mathematicae, 18, 1956.
- ⁶⁵⁶ 34 Dirk Roorda. Interpolation in fragments of classical linear logic. The Journal of Symbolic
 ⁶⁵⁷ Logic, 59(2):419-444, 1994. URL: http://www.jstor.org/stable/2275398.
- Alexis Saurin. A linear perspective on cut-elimination for non-wellfounded sequent calculi with
 least and greatest fixed-points. In *TABLEAUX*, volume 14278 of *Lecture Notes in Computer Science*, pages 203–222. Springer, 2023.
- ⁶⁶¹ **36** Kurt Schütte. *Proof theory*, volume 85. Springer, 1977.
- ⁶⁶² 37 D. S. Shamkanov. Circular proofs for the Gödel-Löb provability logic. Mathematical Notes,
 ⁶⁶³ 96(3), 2014. URL: https://doi.org/10.1134/S0001434614090326.
- ⁶⁶⁴ 38 Lutz Strassburger. Linear Logic and noncommutativity in the Calculus of Structures. PhD
 ⁶⁶⁵ thesis, TU Dresden, 2003.
- Amirhossein Akbar Tabatabai and Raheleh Jalali. Universal proof theory: Semi-analytic
 rules and craig interpolation. Ann. Pure Appl. Log., 176(1):103509, 2025. URL: https:
 //doi.org/10.1016/j.apal.2024.103509, doi:10.1016/J.APAL.2024.103509.
- 669 40 Gaisi Takeuti. Proof theory, volume 81. Courier Corporation, 2013.
- A. S. Troelstra and H. Schwichtenberg. *Basic Proof Theory*. Cambridge Tracts in Theoretical
 Computer Science. Cambridge University Press, 2 edition, 2000.

- 672 **A** LL cut-elimination reduction
- ▶ Definition 22. Cut commutations or External reductions are defined in Figure 4.
- ▶ Definition 23. LL cut-reduction are the reduction rules given in Figure 4, Figure 5
- ⁶⁷⁵ together with the usual cut-elimination rule for first-order quantifiers and the following two
- ⁶⁷⁶ reductions (together with their symmatrical cases):

 \blacksquare the (Cut)/(Cut) commutation

 \blacksquare the (Cut)/(Ax) reduction

B Full proof of proof-relevant interpolation for LL refining Maehara's method

⁶⁷⁹ We shall prove a refined version for cut-free proofs from which Theorem 5 follows directly by ⁶⁸⁰ LL cut-elimination theorem⁴:

• **Theorem 24.** Let Γ , Δ be lists of LL formulas and $\pi \vdash \Gamma$, Δ be cut-free. There exists a LL formula C such that $\mathbb{Voc}(C) \subseteq \mathbb{Voc}(\Gamma) \cap \mathbb{Voc}(\Delta)$ and two cut-free proofs π_1, π_2 of $\vdash \Gamma, C$ and $\vdash C^{\perp}, \Delta$ respectively such that $\frac{\pi_1}{\vdash \Gamma, C} \xrightarrow{\pi_2}_{\vdash C^{\perp}, \Delta}_{(\mathsf{Cut})} \longrightarrow_{\mathsf{cut}}^{\star} \pi$.

⁶⁸⁴ **Proof.** By LL cut-elimination theorem, one can assume that π is cut-free and reason by ⁶⁸⁵ induction on the structure of π and by case on the last inference. We will proceed by ⁶⁸⁶ "introducing cuts" and build new interpolants in such a way as to preserve the denotational ⁶⁸⁷ equivalence of the *interpolated proof* with the proof being constructed.

In fact, we shall prove a slightly stronger result, that is the cut of the interpolating proofs reduces, by cut-elimination, to the interpolated proof (up to exchange rules which are neglected in the following).

⁶⁹¹ If $\pi = \overline{\vdash F, F^{\perp}}$ ^(Ax), $\Gamma = F$, one simply takes $C = F^{\perp} \pi_1 = \pi_2 = \overline{\vdash F, F^{\perp}}$ ^(Ax). (The ⁶⁹² case when $\Gamma = F^{\perp}$ is symmetrical, taking C = F.)

⁴ For simplicity, we omit the polarities used for Lyndon's statement of interpolation below, but this proof would work with the polarities as well, as in the cut-introduction approach in the body of the paper.

In the promotion commutation case, the premisse in $\vdash ! C^{\perp}, ? \Sigma$ is assumed to end with a promotion rule.

Figure 4 Cut-commutations rules, or External reduction rules, where r = (ext, F) and F is the principal occurrence.

$$\frac{\frac{\pi}{\vdash !F^{\perp}, ?\Delta} \stackrel{(!p)}{\vdash ?\Delta, \Gamma} \stackrel{\vdash ?F, \Gamma}{\vdash ?F, \Gamma}_{(\mathsf{Cut})} \stackrel{(?c)}{\longrightarrow} \frac{\pi}{\vdash !F^{\perp}, ?\Delta} \stackrel{(!p)}{\vdash !F^{\perp}, ?\Delta} \stackrel{(!p)}{\vdash !F^{\perp}, ?\Delta} \stackrel{(!p)}{\vdash ?\Delta, ?F, \Gamma}_{(\mathsf{Cut})} \stackrel{(\mathsf{Cut})}{\vdash ?\Delta, ?F, \Gamma}_{(\mathsf{Cut})}$$

$$\frac{\frac{\pi}{\vdash ! F^{\perp}, ? \Delta} \stackrel{(!p)}{\vdash ? \Delta, \Gamma} \xrightarrow{\vdash \Gamma} (?w)}{\vdash ? \Delta, \Gamma} \xrightarrow{(cut)} \stackrel{(?w)}{\longrightarrow} \frac{\vdash \Gamma}{\vdash ? \Delta, \Gamma} (?w) ^{\star}$$

$$\begin{array}{c|c} \displaystyle \frac{\vdash \Gamma}{\vdash \Gamma, \bot} & (\bot) & \displaystyle \frac{}{\vdash 1} & (1) \\ \displaystyle \\ \displaystyle \frac{}{\vdash \Gamma} & \Gamma & (\mathsf{Cut}) & \displaystyle \frac{}{r} & \vdash \Gamma \end{array}$$

Figure 5 Key cut-reduction rules, or Principal reductions, where $r = (\text{princ}, \{F, F'^{\perp}\})$ with $\{F, F'^{\perp}\}$ the principal occurrences that have been reduced.

XX:23

If $\pi = \overline{\vdash F, F^{\perp}}$ (Ax), $\Gamma = F, F^{\perp}$, one simply takes $C = \bot, \pi_{1} = \frac{\pi}{\Gamma, \bot}$ (\bot) and $\pi_{2} = \overline{\vdash 1}^{(1)}$. (The case when Γ is empty is symmetrical, taking C = 1.) (I) (The case when Γ is empty is symmetrical, taking C = 1.) (I) If the last rule is (\otimes), that is $\pi = \frac{\pi'}{\vdash F, \Gamma, \Delta'} = \frac{\pi''}{\vdash G, \Gamma', \Delta''}$ (\otimes), assuming $\Gamma = F \otimes$ (I) $\pi'_{1} \vdash F, \Gamma', C'$, (ii) $\pi'_{2} \vdash C'^{\perp}, \Delta'$, (iii) $\pi''_{1} \vdash G, \Gamma'', C''$ and (iv) $\pi''_{2} \vdash C''^{\perp}, \Delta''$ such that (I) $\pi'_{1} \vdash F, \Gamma', C'$, (ii) $\pi'_{2} \vdash C'^{\perp}, \Delta'$, (iii) $\pi''_{1} \vdash G, \Gamma'', C''$ and (iv) $\pi''_{2} \vdash C''^{\perp}, \Delta''$ such that (I) $\pi'_{1} \vdash F, \Gamma', C'$, (ii) $\pi'_{2} \vdash C'^{\perp}, \Delta'$, (iii) $\pi''_{1} \vdash G, \Gamma'', C''$ and (iv) $\pi''_{2} \vdash C''^{\perp}, \Delta''$ such that (I) $\pi'_{1} \vdash F, \Gamma', \Delta'$ (Cut) $\rightarrow_{cut}^{*} \pi'$ ($\pi''_{1} = \pi''_{2} = \pi''_{2} = \pi''_{2}$ (Cut) $\rightarrow_{cut}^{*} \pi'$ (Cut) $\rightarrow_{cut}^{*} \pi'$ (Cut) (I) $\pi'_{2} \vdash F \otimes G, \Gamma', \Gamma'', C', C''$ (\otimes) $\pi''_{2} = C'^{\perp} \otimes C''^{\perp}, \Delta', \Delta''$ (\otimes). (I) $\pi'_{2} \vdash F \otimes G, \Gamma', \Gamma'', C', C'' = \pi''_{2}$ (Cut) (I) $\pi'_{2} \vdash F \otimes G, \Gamma', \Gamma'', C', C'' = \pi''_{2}$ (Cut) (I) $\pi'_{2} \vdash F \otimes G, \Gamma', \Gamma'', C', C'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', C', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', \Delta', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', \Delta', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', \Delta', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', \Delta', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', \Delta', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', \Delta', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', \Delta', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', \Delta', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', \Delta', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', \Delta', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', \Delta', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', \Delta', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', \Delta', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F \otimes G, \Gamma', \Gamma'', \Delta', \Delta'' = \pi''_{2}$ (Cut) (I) $\pi''_{2} \vdash F$

If the last rule is (8), that is $\pi = \frac{\pi'}{\vdash F, G, \Gamma', \Delta}$ (8), assuming $\Gamma = F \otimes G, \Gamma'$. By induc- $\vdash F \otimes G, \Gamma', \Delta$ (8), that $\mathbb{Voc}(C') \subset \mathbb{Voc}(F, G, \Gamma') \cap \mathbb{Voc}(\Delta)$

tion hypothesis, there is an interpolant C' such that $\operatorname{Voc}(C') \subseteq \operatorname{Voc}(F, G, \Gamma') \cap \operatorname{Voc}(\Delta)$ as well as proofs $\pi'_1 \vdash F, G, \Gamma', C'$ and $\pi'_2 \vdash C'^{\perp}, \Delta$ such that

$$\underbrace{ \frac{\pi_1'}{\vdash F,G,\Gamma',C'} \quad \frac{\pi_2'}{\vdash C'^{\perp},\Delta'}}_{\vdash F,G,\Gamma',\Delta'} \quad \underset{(\mathsf{Cut})}{\longrightarrow^{\star}_{\mathsf{cut}}} \pi'.$$

XX:24 Interpolation as cut-introduction

If the last rule is (1) , that is $\pi = \overline{\vdash 1}$ (1) then the interpolant is trivially \perp or 1 depending 698 on whether $\Gamma = 1$ or not, one interpolating proof being an axiom and the other being π 699 itself: 700

If $\Gamma = 1$, let $C = \bot$, $\pi_1 = \overline{\vdash 1, \bot}$ (Ax) and $\pi_2 = \overline{\vdash 1}$ (1). One easily gets 701 $\frac{\pi_1 \quad \pi_2}{\vdash 1} \quad (\mathsf{Cut}) \quad \longrightarrow_{\mathsf{cut}} \pi.$

If Γ is empty, then let C = 1, $\pi_1 = \overline{\vdash 1}$ (1) and $\pi_2 = \overline{\vdash 1, \perp}$ (Ax). One easily gets 703 $\frac{\pi_1 \quad \pi_2}{\vdash 1} \quad (\mathsf{Cut}) \quad \longrightarrow_{\mathsf{cut}} \pi.$ 704

If the last rule is (\bot) , that is $\pi = \frac{\pi'}{\vdash \Gamma', \Delta}$ and assume $\Gamma = \bot, \Gamma'$. By induction $\Gamma' = \bot, \Gamma', \Delta$ and $\pi'_2 \vdash C'^{\bot}, \Delta$ such that

hypothesis, there is an interpolant
$$C'$$
 and proof $\pi'_1 \vdash \Gamma', C'$ and $\pi'_2 \vdash C'^{\perp}, \Delta$ such that

$$\frac{\frac{\pi_1'}{\vdash \Gamma', C'} \quad \frac{\pi_2'}{\vdash C'^{\perp}, \Delta'}}{\vdash \Gamma', \Delta'} \quad \underset{(\mathsf{Cut})}{\longrightarrow_{\mathsf{cut}}^{\star} \pi'}$$

By setting $C = C', \pi_1 = \frac{\pi'_1}{\vdash \bot, \Gamma', C'}$ (\bot) and $\pi_2 = \pi'_2$, one gets

$$\frac{\frac{\pi_{1}}{\vdash \bot, \Gamma', C} \quad \frac{\pi_{2}}{\vdash C^{\bot}, \Delta'}}{\vdash \bot, \Gamma', \Delta'} \quad (Cut) \quad \longrightarrow_{cut} \frac{\frac{\pi_{1}}{\vdash \Gamma', C} \quad \frac{\pi_{2}}{\vdash C^{\bot}, \Delta'}}{\frac{\vdash \Gamma', \Delta'}{\vdash \bot, \Gamma', \Delta'}} \quad (Cut) \quad \longrightarrow_{cut}^{\star} \pi_{cut}$$

If the last rule is (\top) , that is $\pi = \overline{\vdash \top, \Gamma', \Delta'}^{(\top)}$, assuming $\Gamma = \top, \Gamma'$. Set C = 0, $\pi'_1 = \overline{\vdash \top, \Gamma', 0}^{(\top)}$ and $\pi'_2 = \overline{\vdash \top, \Delta'}^{(\top)}$ In such a case

$$\frac{\overline{\vdash \top, \Gamma', 0} \quad \stackrel{(\top)}{\vdash \top, \Gamma', \Delta'} \quad \stackrel{(\top)}{\vdash \top, \Delta'} \quad \stackrel{(\top)}{\longrightarrow_{\mathsf{cut}}^{\star} \pi}$$

If the last rule is (a), that is $\pi = \frac{\pi'}{\vdash F, \Gamma', \Delta'} = \frac{\pi''}{\vdash G, \Gamma', \Delta'}$ (a), assuming $\Gamma = F \otimes G, \Gamma'$.

By induction hypothesis, there are interpolants C', C'', as well as interpolating proofs (i) $\pi'_1 \vdash F, \Gamma', C'$, (ii) $\pi'_2 \vdash C'^{\perp}, \Delta'$, (iii) $\pi''_1 \vdash G, \Gamma', C''$ and (iv) $\pi''_2 \vdash C''^{\perp}, \Delta'$ such that

$$\frac{\pi_1' \quad \pi_2'}{\vdash F, \Gamma', \Delta'} \quad (\operatorname{Cut}) \quad \longrightarrow_{\operatorname{cut}}^{\star} \pi' \qquad \frac{\pi_1'' \quad \pi_2''}{\vdash G, \Gamma', \Delta'} \quad (\operatorname{Cut}) \quad \longrightarrow_{\operatorname{cut}}^{\star} \pi'$$

Let $C = C' \oplus C''$ and let $\pi_1 = \frac{\pi_1'}{\vdash F, \Gamma', C' \oplus C''} \stackrel{(\oplus^1)}{\longrightarrow} \frac{\pi_1''}{\vdash G, \Gamma', C' \oplus C''} \stackrel{(\oplus^2)}{\overset{(\otimes)}{\mapsto}}$ and 706

$$\pi_{2} = \frac{\pi_{2}' - \pi_{2}''}{\vdash C'^{\perp} \otimes C''^{\perp}, \Delta'} \quad (\&) \; .$$

One observes that

If the last rule is $(\oplus^i), i \in \{1, 2\}$, that is $\pi = \frac{\pi'}{\vdash F_i, \Gamma', \Delta} (\oplus^i)$, assuming $\Gamma = F_1 \oplus F_2, \Gamma'$. By induction hypothesis, there is an interpolant C' such that $\operatorname{Voc}(C') \subseteq \operatorname{Voc}(F_i, \Gamma') \cap \operatorname{Voc}(\Delta)$ as well as proofs $\pi'_1 \vdash F_i, \Gamma', C'$ and $\pi'_2 \vdash C'^{\perp}, \Delta$ such that

$$\frac{\frac{\pi_1'}{\vdash F_i, \Gamma', C'} \quad \frac{\pi_2'}{\vdash C'^{\perp}, \Delta'}}{\vdash F_i, \Gamma', \Delta'} \quad \underset{(\mathsf{Cut})}{\longrightarrow_{\mathsf{cut}}^{\star} \pi'}.$$

Setting C = C', $\pi_1 = \frac{\pi'_1}{\vdash F_1 \oplus F_2, \Gamma', C}$ (\oplus^i) and $\pi_2 = \pi'_2$ we get the following cutreduction starting with a cut-commutation case:

$$\begin{array}{c|c} \frac{\pi_{1}}{\vdash F_{1} \oplus F_{2}, \Gamma', C} & \frac{\pi_{2}}{\vdash C^{\perp}, \Delta'} & (\operatorname{Cut}) & \longrightarrow_{\operatorname{cut}} \\ \hline \\ \frac{\pi_{1}'}{\vdash F_{1} \oplus F_{2}, \Gamma', \Delta'} & \frac{\pi_{2}'}{\vdash C'^{\perp}, \Delta'} \\ \frac{\pi_{1}'}{\vdash F_{i}, \Gamma', C'} & \frac{\pi_{2}'}{\vdash C'^{\perp}, \Delta'} \\ \hline \\ \frac{\Gamma_{i}, \Gamma', C'}{\vdash F_{i} \oplus F_{2}, \Gamma', \Delta'} & (\operatorname{Cut}) & \longrightarrow_{\operatorname{cut}}^{\star} & \pi. \end{array}$$

If the last rule is (?d) , that is $\pi = \frac{\pi'}{\vdash F, \Gamma', \Delta}$ assuming $\Gamma = ?F, \Gamma'$. By induction $\vdash ?F, \Gamma', \Delta$ (?d)

hypothesis, there is an interpolant C' such that $\mathbb{V}oc(C') \subseteq \mathbb{V}oc(F, \Gamma') \cap \mathbb{V}oc(\Delta)$ as well as proofs $\pi'_1 \vdash F, \Gamma', C'$ and $\pi'_2 \vdash C'^{\perp}, \Delta$ such that

$$\frac{\frac{\pi_1'}{\vdash F, \Gamma', C'} \quad \frac{\pi_2'}{\vdash C'^{\perp}, \Delta}}{\vdash F, \Gamma', \Delta} \quad \underset{(\mathsf{Cut})}{\longrightarrow} \overset{\star}{\longrightarrow} ^{\star}_{\mathsf{cut}} \pi'.$$

XX:26 Interpolation as cut-introduction

By setting
$$C = C', \pi_1 = \frac{\pi'_1}{\vdash ?F, \Gamma', C'}$$
 (?d), one gets:

$$\frac{\frac{\pi_{1}}{\vdash ?F,\Gamma',C} \quad \frac{\pi_{2}}{\vdash C^{\perp},\Delta}}{\vdash ?F,\Gamma',\Delta} \quad (\mathsf{Cut}) \quad \longrightarrow_{\mathsf{cut}} \frac{\frac{\pi_{1}'}{\vdash F,\Gamma',C'} \quad \frac{\pi_{2}'}{\vdash C'^{\perp},\Delta}}{\frac{\vdash F,\Gamma',\Delta}{\vdash ?F,\Gamma',\Delta} \quad (\mathsf{Cut})} \quad \longrightarrow_{\mathsf{cut}}^{\star} \pi.$$

If the last rule is (!p) , that is $\pi = \frac{\pi'}{\vdash F, ?\Gamma', ?\Delta'} \xrightarrow{(!p)} \text{assuming } \Gamma = !F, ?\Gamma' \text{ and } \Delta = ?\Delta'.$

By induction hypothesis, there is an interpolant C' such that $\operatorname{Voc}(C') \subseteq \operatorname{Voc}(F, ?\Gamma') \cap \operatorname{Voc}(\Delta)$ as well as proofs $\pi'_1 \vdash F, ?\Gamma', C'$ and $\pi'_2 \vdash {C'}^{\perp}, ?\Delta'$ such that

$$\frac{\frac{\pi_1'}{\vdash F, ?\,\Gamma', C'} \quad \frac{\pi_2'}{\vdash C'^{\perp}, ?\,\Delta'}}{\vdash F, ?\,\Gamma', ?\,\Delta'} \quad \underset{(\mathsf{Cut})}{\longrightarrow_{\mathsf{cut}}^{\star}} \pi'.$$

By setting $C = ?C', \pi_1 = \frac{\pi'_1}{\vdash F, ?\Gamma', ?C'} \xrightarrow{(?d)}_{(!p)}$ and $\pi_2 = \frac{\pi'_2}{\vdash !C'^{\perp}, ?\Delta'} \xrightarrow{(!p)}$, one gets:

$$\frac{\pi_{1}}{\vdash !F, ?\Gamma', ?\Delta'} \quad (Cut) \longrightarrow_{cut} \begin{array}{c} \frac{\pi_{1}'}{\vdash F, ?\Gamma', C'} & (?d) & \frac{\pi_{2}}{\vdash C'^{\perp}, ?\Delta'} \\ F, ?\Gamma', C & (?d) & \frac{\vdash C'^{\perp}, ?\Delta'}{\vdash C^{\perp}, ?\Delta'} \end{array} \\ (!p) & (Cut) \\ \frac{\vdash F, ?\Gamma', ?\Delta'}{\vdash !F, ?\Gamma', ?\Delta'} & (!p) \\ \xrightarrow{cut} & \frac{\pi_{1}' & \pi_{2}'}{\vdash !F, ?\Gamma', ?\Delta'} \end{array} \\ (Cut) & (Cut) \\ \frac{\tau_{1}' & \pi_{2}'}{\vdash !F, ?\Gamma', ?\Delta'} & (!p) \\ \xrightarrow{cut} & \pi. \end{array}$$

If the last rule is (?w) , that is $\pi = \frac{\pi'}{\vdash \Gamma', \Delta}$ assuming $\Gamma = ?F, \Gamma'$. By induction hypothesis, there is an interpolant C' such that $\operatorname{Voc}(C') \subseteq \operatorname{Voc}(\Gamma') \cap \operatorname{Voc}(\Delta)$ as well as

proofs $\pi'_1 \vdash \Gamma', C'$ and $\pi'_2 \vdash {C'}^{\perp}, \Delta$ such that

$$\frac{\overbrace{\vdash \Gamma', C'}^{\pi'_1} \quad \overbrace{\vdash C'^{\perp}, \Delta}^{\pi'_2}}{\vdash \Gamma', \Delta} \quad \underset{(\mathsf{Cut})}{\longrightarrow_{\mathsf{cut}}^{\star}} \pi'.$$

By setting $C = C', \ \pi_1 = \frac{\pi'_1}{\vdash ?F, \Gamma', C}$ (?w), one gets $\mathbb{V}oc(C) \subseteq \mathbb{V}oc(\Gamma) \cap \mathbb{V}oc(\Delta)$ and:

$$\frac{\frac{\pi_{1}}{\vdash ?F, \Gamma', C} \quad \frac{\pi_{2}}{\vdash C^{\perp}, \Delta}}{\vdash ?F, \Gamma', \Delta} \quad (Cut) \quad \longrightarrow_{cut} \frac{\frac{\pi_{1}'}{\vdash \Gamma', C'} \quad \frac{\pi_{2}'}{\vdash C'^{\perp}, \Delta}}{\frac{\vdash \Gamma', \Delta}{\vdash ?F, \Gamma', \Delta}} \quad (Cut) \quad \longrightarrow_{cut}^{\star} \pi.$$

If the last rule is (?c) , that is $\pi = \frac{\pi'}{\vdash ?F, ?F, \Gamma', \Delta}$ assuming $\Gamma = ?F, \Gamma'$. By induction hypothesis, there is an interpolant C' such that $\operatorname{Voc}(C') \subseteq \operatorname{Voc}(?F, ?F, \Gamma') \cap \operatorname{Voc}(\Delta)$ as

well as proofs $\pi'_1 \vdash ?F, ?F, \Gamma', C'$ and $\pi'_2 \vdash {C'}^{\perp}, \Delta$ such that

$$\frac{\frac{\pi_1'}{\vdash ?F, ?F, \Gamma', C'} \quad \overline{\vdash C'^{\perp}, \Delta}}{\vdash ?F, ?F, \Gamma', \Delta} \quad \xrightarrow{\mathsf{(Cut)}} \quad \longrightarrow_{\mathsf{cut}}^{\star} \pi'.$$

By setting C = C', $\pi_1 = \frac{\pi'_1}{\vdash ?F, \Gamma', C'}$ (?c) and $\pi_2 = \pi'_2$ one gets:

$$\frac{\begin{array}{cccc}
\frac{\pi_{1}}{\vdash ?F,\Gamma',C} & \frac{\pi_{2}}{\vdash C^{\perp},\Delta} \\
\frac{\pi_{1}'}{\vdash ?F,\Gamma',C} & \frac{\pi_{2}'}{\vdash C'^{\perp},\Delta} \\
\frac{\pi_{1}'}{\vdash ?F,?F,\Gamma',C'} & \frac{\pi_{2}'}{\vdash C'^{\perp},\Delta} \\
\frac{\mu ?F,?F,\Gamma',\Delta}{\vdash ?F,\Gamma',\Delta} & (Cut) & \longrightarrow_{cut}^{\star} & \pi.
\end{array}$$

If the last rule is (\forall) , that is $\pi = \frac{\pi'}{\vdash F, \Gamma', \Delta} \quad (\forall) \quad x \notin \mathsf{FV}(\Gamma', \Delta) \text{ assuming } \Gamma = \forall xF, \Gamma'.$

By induction hypothesis, there is an interpolant C' such that $\operatorname{Voc}(C') \subseteq \operatorname{Voc}(F, \Gamma') \cap \operatorname{Voc}(\Delta)$ as well as proofs $\pi'_1 \vdash F, \Gamma', C'$ and $\pi'_2 \vdash C'^{\perp}, \Delta$ such that

$$\frac{\frac{\pi_1'}{\vdash F, \Gamma', C'} \quad \frac{\pi_2'}{\vdash C'^{\perp}, \Delta}}{\vdash F, \Gamma', \Delta} \quad \underset{(\mathsf{Cut})}{\longrightarrow} \overset{\star}{\underset{\mathsf{cut}}} \pi'.$$

By setting $C = \exists x.C', \ \pi_1 = \frac{\pi'_1}{\vdash F, \Gamma', \exists x.C'} \xrightarrow{(\exists)} \text{and } \pi_2 = \frac{\pi'_2}{\vdash \forall xC'^{\perp}, \Delta} (\forall) \text{ one gets:}$

⁷⁰⁸ If the last rule is (I), that is $\pi = \frac{\pi'}{\vdash F\{y/x\}, \Gamma', \Delta}$ (I) assuming $\Gamma = \exists xF, \Gamma'$. (I)

XX:28 Interpolation as cut-introduction

In this case, Note that we treat only the case of a FO language containing no function 709 symbols. 710

By induction hypothesis, there is an interpolant C' such that $\operatorname{Voc}(C') \subseteq \operatorname{Voc}(F\{y/x\}, \Gamma') \cap$ $\mathbb{V}oc(\Delta)$ as well as proofs $\pi'_1 \vdash F\{y/x\}, \Gamma', C'$ and $\pi'_2 \vdash {C'}^{\perp}, \Delta$ such that

$$\frac{\frac{\pi'_1}{\vdash F\{y/x\}, \Gamma', C'} \quad \frac{\pi_2}{\vdash C'^{\perp}, \Delta}}{\vdash F\{y/x\}, \Gamma', \Delta} \quad \xrightarrow{\text{(Cut)}} \quad \longrightarrow_{\text{cut}}^{\star} \pi'.$$

In this case, we reason by case on whether y occurs in Γ', Δ : 711

712

If y occurs in both, then we simply take C = C' as interpolant, $\pi_1 = \frac{\pi'_1}{\vdash \exists xF, \Gamma'}$ (\exists) and $\pi_2 = \pi'_2$. and we have $\operatorname{Voc}(C) = \operatorname{Voc}(C') \subseteq \operatorname{Voc}(F, \Gamma') \cap \operatorname{Voc}(\Delta) = \operatorname{Voc}(\exists xF, \Gamma') \cap \operatorname{Voc}(\Delta)$ 713 $\mathbb{V}\mathsf{oc}(\Delta)$ 714

= If y occurs in Γ' but not in Δ , then we set $C = \exists y C', \pi_1 = \frac{\pi'_1}{\exists x F, \Gamma', C'} \stackrel{(\exists)}{\exists x F, \Gamma', \exists y C'}$ and 1

$$\pi_{2} = \frac{\pi_{2}}{\forall y C'^{\perp}, \Delta} \quad (\forall) \text{ one gets:}$$

$$\frac{\pi_{1}}{\vdash \exists x.F, \Gamma', \Delta} \quad (\mathsf{Cut}) \quad \longrightarrow_{\mathsf{cut}} \quad \frac{\pi_{1}'}{\vdash \exists xF, \Gamma', C'} \quad \stackrel{(\exists)}{=} \quad \pi_{2}' \quad (\mathsf{Cut}) \quad \longrightarrow_{\mathsf{cut}} \quad \frac{\pi_{1}' \quad \pi_{2}'}{\vdash \exists xF, \Gamma', \Delta} \quad \stackrel{(\mathsf{Cut})}{=} \quad \xrightarrow{\tau_{1}' \quad \pi_{2}'} \quad (\mathsf{Cut}) \quad \longrightarrow_{\mathsf{cut}} \quad \frac{\pi_{1}' \quad \pi_{2}'}{\vdash \exists xF, \Gamma', C'} \quad \stackrel{(\exists)}{=} \quad \xrightarrow{\tau_{1}' \quad \pi_{2}'} \quad (\mathsf{Cut}) \quad \xrightarrow{\tau_{1}' \quad \pi_{2}' \quad (\mathsf{Cut})} \quad \xrightarrow{\tau_{2}' \quad (\mathsf{Cut})} \quad \xrightarrow{\tau$$

= If y occurs in Δ but not in Γ' , then we set $C = \forall yC', \pi_1 = \frac{\pi'_1}{\vdash \exists xF, \Gamma', C'} \stackrel{(\exists)}{\vdash \exists xF, \Gamma', \forall y. C'}$ and /

$$\pi_{2} = \frac{\pi_{2}}{\vdash \exists y C'^{\perp}, \Delta} \quad (\exists) \text{ . One gets:}$$

$$\frac{\pi_{1} \quad \pi_{2}}{\vdash \exists x F, \Gamma', \Delta} \quad (\mathsf{Cut}) \quad \longrightarrow_{\mathsf{cut}} \quad \frac{\pi_{1}'}{\vdash \exists x F, \Gamma', C'} \quad (\exists) \quad \pi_{2}' \quad (\mathsf{Cut})$$

$$\longrightarrow_{\mathsf{cut}} \quad \frac{\pi_{1}' \quad \pi_{2}'}{\vdash F\{y/x\}, \Gamma', \Delta} \quad (\mathsf{Cut})$$

$$\longrightarrow_{\mathsf{cut}} \quad \pi.$$

715

716

717

С Details on linear embeddings of LJ and LK and proof-relevant interpolation for LJ and LK.

In the present section, we deduce our proof-relevant statement of interpolation for LJ and LK 718 as a direct corollary of the properties of the linear embeddings of classical and intuitionistic 719 logics into linear logic and the proof-relevant interpolation theorem for LL: 720

First one needs to extend the results of the previous section on proof-relevant interpolation

 $_{722}$ to two-sided LL, which is clear. The only additional care amount to dealing properly

with the notion of positive/negative subformula in order to obtain the refined Lyndoninterpolation result, but there is no difficulty in doing so and this simply amounts to the

⁷²⁵ usual notion of positive/negative suboccurrences and that linear embeddings (and the ⁷²⁶ reverse translation, skelettons, preseve the polarities of subformulas);

Second, consider some provable sequents $\Gamma \vdash_{\mathsf{LK}} \Delta$ (resp. $\Gamma \vdash_{\mathsf{LJ}} A$) and their respective cut-free proofs under consideration π_{LK} (resp. π_{LJ}) and consider a splitting of the sequents: $\Gamma' \vdash \Delta'$ and $\Gamma'' \vdash \Delta''$ (resp. $\Gamma' \vdash A$);

Consider the linear sequents and proofs corresponding to those sequents via linear
 translations which do not introduce additional cuts (see for instance [35] for such cut-free
 translations) and the linear translations of the splittings considered above.

Apply proof-relevant LL interpolation to obtain a formula C_k (resp. C_j) in the common vocabulary and proofs π_1^l, π_2^l interpolating *wrt*. C_k (resp. C_j);

By erasing all the linear information of C_k (resp. C_j) and π_1, π_2 (which is called their skeletons), this provides us with the expected solution. Indeed, a cut-reduction step in LL (resp. its intuitionistic fragment, ILL) can be simulated in LK (resp. LJ) via the skeleton translation.

More details are provided in the following paragraphs. Note that the following elements are essentially standard, from original results by Danos, Joinet and Schellinx [13].

Definition 25 (Skeleton). For A an LL formula, we define Sk(A) inductively:

 $\mathsf{Sk}(A \otimes B)$ = $\mathsf{Sk}(A) \wedge \mathsf{Sk}(B)$ $Sk(A \otimes B)$ = $\mathsf{Sk}(A) \lor \mathsf{Sk}(B)$ Sk(!A)Sk(A)= $\mathsf{Sk}(A) \lor \mathsf{Sk}(B)$ $\mathsf{Sk}(A) \wedge \mathsf{Sk}(B)$ $Sk(A \otimes B)$ $\mathsf{Sk}(A \oplus B)$ Sk(?A)Sk(A)== = 742 Sk(1)= $\mathsf{Sk}(\top) = \top$ $Sk(\perp)$ = Sk(0) = FSk(a)= a $\mathsf{Sk}(A \multimap B) =$ $\mathsf{Sk}(A) \Rightarrow \mathsf{Sk}(B)$

⁷⁴³ Let π be a two-sided LL proof of $\Gamma \vdash \Delta$. $\mathsf{Sk}(\pi)$ is the LK proof of $\mathsf{Sk}(\Gamma) \vdash \mathsf{Sk}(\Delta)$ obtained ⁷⁴⁴ by the following recursive process by case analysis on the last rule r of π : (i) if $r \in \{(1p), (?d)\}$, ⁷⁴⁵ then $\mathsf{Sk}(\pi)$ is the skeleton of the premise of π ; (ii) otherwise, apply the corresponding rule ⁷⁴⁶ with, for premises, the skeletons of the premises of π .

Proposition 26. For any LL proof π of s, $Sk(\pi)$ is a LK proof of Sk(s).

A standard result of LL proof theory, developed by Danos, Joinet and Schellinx [13], is that there exist linear decorations for LK:

Proposition 27. For any LK sequent s and any LK proof π , there is a linear decoration of π , that is a LL proof π^d such that $Sk(\pi^d) = \pi$.

Moreover, the skeleton maps cut-related LL-proofs to cut-related LK proofs (resp. LJ proofs): LL cut-reduction sequences can be simulated in LK (resp. LJ).

Eventually, proof relevant interpolation for LJ and LK is therefore a direct and simple
 corollary and the above theory of linear decorations together with proof-relevant interpolation
 theorem for LL.

⁷⁵⁷ **D** Details on interpolation as cut-introduction

⁷⁵⁸ We recall and provide further details on the notions of decorated and coherent proof:

XX:30 Interpolation as cut-introduction

Definition 28. \blacksquare A decorated proof is an LL proof st. each sequent is equipped with a splitting.

A coherent decorated proof is a decorated proof such that for each node, the splitting of the conclusion and of its premises is coherent wrt the ancestor relation: a formula belonging to the left (resp. right) component of the splitting has all its ancestors belonging to the left (resp. right) component of the splitting. More precisely:

- that each ancestor of an auxiliary formula belonging to the left (resp. right) component
 of the splitting belongs to the left (resp. right) component of the splitting of the
 corresponding premise;
- that each immediate subformula of a principal formula which belongs to the left (resp. right) component of the paritition, itself belongs to the left (resp. right) component of the splitting of its premises.
- In a two-sided calculus, the coherence condition can be refined as follows:

each ancestor of an auxiliary formula belonging to the antecedent (resp. succedent) of
left component of the splitting belongs to the antecedent (resp. succedent) of the left
component of the splitting of the corresponding premise;

- that each ancestor of an auxiliary formula belonging to the antecedent (resp. succedent) of right component of the splitting belongs to the antecedent (resp. succedent) of the right component of the splitting of the corresponding premise;
- that each positive immediate subformula of a principal formula which belongs to the antecedent (resp. succedent) of the left component of the splitting, itself belongs to the antecedent (resp. succedent) of the left component of the splitting of its premises;
- 4. that each positive immediate subformula of a principal formula which belongs to the antecedent (resp. succedent) of the right component of the splitting, itself belongs to the antecedent (resp. succedent) of the right component of the splitting of its premises;
- that each negative immediate subformula of a principal formula which belongs to the antecedent (resp. succedent) of the left component of the splitting, itself belongs to the succedent (resp. antecedent) of the left component of the splitting of its premisses;
- that each negative immediate subformula of a principal formula which belongs to the antecedent (resp. succedent) of the right component of the splitting, itself belongs to the succedent (resp. antecedent) of the right component of the splitting of its premisses.

⁷⁹⁰ D.1 Details on the Proof of the Main Lemma

⁷⁹¹ We now provide full details on the proof of Lemma 10 given in Section 4.4.

792 D.1.1 Axiom case

796

⁷⁹³ If $\pi = \overline{\vdash F, F^{\perp}}$ (Ax), one simply takes $I = F^{\perp}, \pi_1^l = \overline{\vdash F, F^{\perp}}$ (Ax) and $\pi_1^r = \overline{\vdash F, F^{\perp}}$ (Ax).

The cut between π_1^l and π_1^r reduces to π by one cut-axiom reduction step.

⁷⁹⁷ If $\pi = \overline{\vdash F, F^{\perp}}$ ^(Ax), the case is symmetrical taking I = F.

⁷⁹⁹ = If $\pi = \overline{\vdash F, F^{\perp}}$ ^(Ax), one takes $I = \perp, \pi_1^l = \frac{\pi}{\vdash F, F^{\perp}, \perp}$ ^(\perp) and $\pi_1^r = \overline{\vdash 1}$ ⁽¹⁾. ⁸⁰⁰ The cut of π_1^l and π_1^r reduces to π by a key $1/\perp$ case. ⁸⁰¹ ⁸⁰² = If $\pi = \overline{\vdash F, F^{\perp}}$ ^(Ax), the case is symmetrical to the previous one, taking I = 1: ⁸⁰³ $\pi_1^l = \overline{\vdash 1}$ ⁽¹⁾ and $\pi_1^r = \frac{\pi}{\vdash F, F^{\perp}, \perp}$ ^(\perp).

804 D.1.2 Logical rules

We analyze the possible cases for a logical rule involved in an elementary PRIS. Note that in each case, the principal formula may be part of the left or right part of the splitting; we treat only one case each time since the other is symmetrical be taking the dual interpolant and exchanging π^L and π^R .

⁸⁰⁹ If the last rule is (8), that is if
$$\pi = \frac{ \stackrel{\pi_1^L}{\Gamma} I \quad \vdash I^{\perp}, \stackrel{\pi_1^R}{\Delta, A, B}}{ \stackrel{\vdash \Gamma, \Delta, A, B}{\vdash \Gamma, \Delta, A \otimes B}}$$
 (Cut) then taking $I' = I$,

 $\pi^{L} = \pi_{1}^{L} \text{ and } \pi^{R} = \frac{\prod_{i=1}^{R} \pi_{1}^{R}}{\prod_{i=1}^{L} \Delta, A, B} \quad \text{(s)} \text{ we obtain a solved PRIS } \pi' \text{ such that } \pi \leftarrow_{\mathsf{cut}} \pi'$

812 If the last rule is (
$$\otimes$$
), that is if $\pi = \frac{\vdash \Gamma_1^{\perp}, I_1 \qquad \vdash I_1^{\perp}, \Delta_1, A}{\vdash \Gamma_1, \Delta_1, A}$ (Cut) $\frac{\vdash \Gamma_2, I_2 \qquad \vdash I_2^{\perp}, \Delta_2, B}{\vdash \Gamma_2, \Delta_2, B}$ (Cut) $\frac{\vdash \Gamma_2, \Delta_2, B}{\vdash \Gamma_2, \Delta_2, B}$ (Cut)

then setting
$$I = I_1 \otimes I_2, \pi^L = \frac{\prod_{1=1}^{L} \prod_{1=1}^{L} \prod_{1$$

one gets a solved PRIS π' such that $\pi \leftarrow_{cut}^* \pi'$ by a commutative reduction of (Cut) and a key (\otimes)/(\otimes) case.

⁸¹⁶ If the last rule is
$$(\perp)$$
, that is if $\pi = \frac{ \prod_{i=1}^{n} \prod_{j=1}^{n} \prod$

 $\underset{\text{sup}}{\overset{\text{H}}{\underset{\text{H}}}} \qquad \frac{\vdash \prod_{i=1}^{L} \prod_{i=1}^{L} (1)}{\vdash \bot, \Gamma, I_{1}} \quad \text{(L)} \quad \text{and} \quad \pi^{R} = \pi_{1}^{R}, \text{ one gets a solved PRIS } \pi' = \frac{\vdash \prod_{i=1}^{L} \prod_{i=1}^{L} \prod_{i=1}^{L} \prod_{i=1}^{R} \prod_{i=1}^{R} \prod_{i=1}^{R} \prod_{i=1}^{R} \prod_{i=1}^{R} \prod_{i=1}^{L} \prod_{i=1}^{L} \prod_{i=1}^{R} \prod_{i=1$

If the last rule is (1), that is if $\pi = \overline{\vdash 1}^{(1)}$ (which is indeed both an initial PRIS and an elementary PRIS), let I = 1 and $\pi_1^L = \overline{\vdash 1}^{(1)}$ and $\pi_1^R = \overline{\vdash \perp, 1}^{(A\times)}$. One gets a solved PRIS π' such that $\pi \leftarrow -\star_{cut} \pi'$ by a key $1/\bot$ case.

⁸²² If the last rule is
$$(\oplus^{i}), i \in \{1, 2\}$$
, that is if $\pi = \frac{ \prod_{i=1}^{\pi_{1}^{L}} \prod_{i=1}^{\pi_{1}^{L}} (Cut)}{ \prod_{i=1}^{L} \prod_{i=1}^{\pi_{1}^{L}} \prod_{i=1}^{L} (Cut)}$, then setting

XX:32 Interpolation as cut-introduction

$$I = I_{1}, \pi^{L} = \frac{\downarrow - F_{1}^{\pi L}, I_{1}}{\downarrow - F_{1} \oplus \mathcal{P}_{2}, \Gamma, I_{1}} (\oplus) \text{ and } \pi^{R} = \pi_{1}^{R}, \text{ one gets a solved PRIS } \pi' \text{ such that}$$

$$\pi \leftarrow _{\text{cut}}^{*} \pi' \text{ by a commutative reduction of (cut).}$$
If the last rule is (a), that is if
$$\pi = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{1}^{\pi L}, \Delta}{\vdash F_{1}, \Gamma, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{1}^{\pi L}, \Delta}{\vdash F_{1}, \Gamma, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{2}^{\pi L}, \Delta}{\vdash F_{1}, \Gamma, \Delta} (\text{cut})$$

$$\frac{\vdash F_{1}^{\pi L}, \Delta \to I_{2}^{\pi L}, \Delta}{\vdash I_{1}^{\pm}, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{2}^{\pi L}, \Delta}{\vdash F_{1}, \Gamma, I_{1} \to I_{2}} (\oplus) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{2}^{\pi L}, \Delta}{\vdash F_{1}^{\pm}, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{2}^{\pi L}, \Delta}{\vdash I_{1}^{\pm}, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{2}^{\pi L}, \Delta}{\vdash I_{1}^{\pm}, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{2}^{\pi L}, \Delta}{\vdash I_{1}^{\pm}, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{2}^{\pi L}, \Delta}{\vdash I_{1}^{\pm}, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{2}^{\pi L}, \Delta}{\vdash I_{1}^{\pm}, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{2}^{\pi L}, \Delta}{\vdash I_{1}^{\pm}, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{1}^{\pi L}, \Delta}{\vdash I_{1}^{\pm}, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{1}^{\pi L}, \Delta}{\vdash I_{1}^{\pm}, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{1}^{\pi L}, \Delta}{\vdash I_{1}^{\pm}, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{1}^{\pi L}, \Delta}{\vdash I_{1}^{\pi}, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1} \to I_{1}^{\pi L}, \Delta}{\vdash F_{1}^{\pi L}, I_{1} \to I_{1}^{\pi L}, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1}^{\pi L}, I_{1}^{\pi L}, I_{1}^{\pi L}, I_{1}^{\pi L}, \Delta}{\vdash F_{1}^{\pi L}, \Delta} (\text{cut}) = \frac{\vdash F_{1}^{\pi L}, I_{1}^{\pi L}, I_{1}^{$$

so that π indeed cut-expands to π' . If the last rule is (\top) , that is if $\pi = \overline{\vdash \top, \Gamma, \Delta}$ (\top) , by setting $I = 0, \pi^L = \overline{\vdash \top, \Gamma, 0}$ (\top) and $\pi^R = \overline{\vdash \top, \Delta}$ (\top) . Then $\pi' = \frac{\pi^L - \pi^R}{\vdash \top, \Gamma, \Delta}$ (Cut) is a solved PRIS to which π cut-expands by a (\top) commutation case. If the last rule is (?d), that is $\pi = \frac{\vdash \frac{\pi_L^T}{F, \Gamma, I_1} - \vdash I_1^{\frac{\pi_L}{1}}, \Delta}{\vdash P, \Gamma, \Delta}$ (Cut), then setting $I = I_1, \pi^L = \frac{\vdash F, \Gamma, \Delta}{\vdash P, \Gamma, \Delta}$ (?d)

$$\frac{\vdash \stackrel{\pi_1^L}{F, \Gamma, I_1}}{\vdash \stackrel{?F, \Gamma, I_1}{\vdash} (?d)} \text{ and } \pi^R = \pi_1^R \text{ one gets:}$$

$$\frac{ \vdash \stackrel{\pi}{}_{F,\Gamma,I}^{L} \vdash \stackrel{\pi}{}_{I^{\perp},\Delta}^{R} }{ \vdash \stackrel{?}{F,\Gamma,\Delta} }_{(\mathsf{Cut})} \xrightarrow{(\mathsf{Cut})} \frac{ \xrightarrow{\pi_{1}^{L}} \vdash \stackrel{\pi_{1}^{L}}{\vdash F,\Gamma,I_{1}} }_{\frac{\vdash F,\Gamma,\Delta}{\vdash \stackrel{?}{F},\Gamma,\Delta} (?\mathsf{d})} (\mathsf{Cut}) = \pi$$

so that π indeed cut-expands to π' .

If the last rule is (!p) , that is $\pi = \frac{\vdash F, ?\Gamma, I_1 \qquad \vdash I_1^{\perp}, ?\Delta}{\vdash F, ?\Gamma, ?\Delta}$ (Cut) , then by setting $I = ?I_1$,

$$\pi^{L} = \frac{\vdash F, ?\Gamma, I_{1}}{\vdash F, ?\Gamma, ?I_{1}} \xrightarrow{(?d)} \text{and} \pi^{R} = \frac{\pi_{1}^{R}}{\vdash I_{1}^{\perp}, ?\Delta} \xrightarrow{(!p)} , \text{ one gets:}$$

$$\begin{array}{ccc} \underbrace{\vdash !F, ?\Gamma, ?I_{1} & \vdash !I_{1}^{\pi_{\perp}^{R}}, ?\Delta}_{\vdash !F, ?\Gamma, ?\Delta} & (\mathrm{Cut}) & \longrightarrow_{\mathrm{cut}} & \underbrace{\vdash F, ?\Gamma, I_{1}}_{\vdash F, ?\Gamma, I} & (?\mathrm{d}) & \underbrace{\vdash I_{1}^{\perp}, ?\Delta}_{\vdash I_{1}^{\perp}, ?\Delta} & (!\mathrm{p}) \\ & \underbrace{\vdash F, ?\Gamma, ?\Delta}_{\vdash !F, ?\Gamma, ?\Delta} & (\mathrm{Cut}) \\ & \underbrace{\vdash F, ?\Gamma, ?\Delta}_{\vdash !F, ?\Gamma, ?\Delta} & (!\mathrm{p}) \\ & \underbrace{\vdash F, ?\Gamma, ?\Delta}_{\vdash !F, ?\Gamma, ?\Delta} & (\mathrm{Cut}) = \pi \end{array}$$

so that π indeed cut-expands to π' . 830

If the last rule is (?w) , that is $\pi = \frac{ \begin{pmatrix} \pi_1^L & \pi_1^R \\ \Gamma, I_1 & \Gamma \\ I_1^\perp, \Delta \\ \hline \begin{pmatrix} \Gamma, \Delta \\ \Gamma \\ \Gamma \\ \hline \Gamma \\ \hline \end{pmatrix} (Cut)$, then by setting $I = I_1$,

 $\pi^{L} = \frac{\vdash \prod_{i=1}^{n} I_{i}}{\vdash \frac{?}{F, \Gamma, I_{1}}} \quad \text{and} \ \pi^{R} = \pi_{1}^{R}, \text{ we ensure that, since } \mathbb{V}\mathsf{oc}(I) \subseteq \mathbb{V}\mathsf{oc}(\Gamma) \cap \mathbb{V}\mathsf{oc}(\Delta),$ $\pi' = \frac{\vdash ?F, \Gamma, I_1}{\vdash ?F, \Gamma, I} \stackrel{\pi^R}{\vdash I^{\perp}, \Delta} \quad \text{(Cut)} \text{ is indeed a solved PRIS and that}$ $L = \pi^R$

$$\pi' \longrightarrow_{\mathsf{cut}} \frac{\vdash \prod_{i=1}^{\pi_{1}} I_{1} \vdash I_{1}^{-1}, \Delta}{\vdash \prod_{i=1}^{r}, \prod_{i=1}^{r}, \Delta} (\mathsf{Cut}) = \pi.$$

by a cut commutation rule so that
$$\pi$$
 indeed cut-expands to π' as expected.

If the last rule is (?c) , that is $\pi = \frac{\vdash ?F, ?F, \Gamma, I_1 \quad \vdash I_1^{\perp}, \Delta}{\underbrace{\vdash ?F, ?F, \Gamma, \Delta}_{\vdash ?F, \Gamma, \Delta}}$ (Cut) by setting $I = I_1$,

$$\pi^{L} = \frac{\vdash ?F, ?F, \Gamma, I_{1}}{\vdash ?F, \Gamma, I} \quad (?c) \quad \text{and} \ \pi^{R} = \pi_{1}^{R} \text{ one gets:}$$

$$\pi' = \frac{\vdash ?F, \Gamma, I}{\vdash ?F, \Gamma, I} \vdash \frac{\pi^{R}}{I^{\perp}}, \Delta \quad (Cut) \quad \longrightarrow_{cut} \quad \frac{\vdash ?F, ?F, \Gamma, I_{1} \quad \vdash I_{1}^{\perp}, \Delta}{\vdash ?F, ?F, \Gamma, \Delta} \quad (Cut) = \pi$$

so that π' is indeed a solved PRIS to which π cut-expands. 832

XX:34 Interpolation as cut-introduction

If the last rule is
$$(\forall)$$
, that is $\pi = \frac{\left| \begin{array}{c} \prod_{i=1}^{n} \prod_{i=1}^{L} \prod_{i=1}^{n} \prod_{$

have that $\pi = \vdash \exists xF, \Gamma, \Delta$ via a cut-commutation rule. = If y occurs in Γ but not in Δ , then we set $I = \exists yI_1, \pi^L = \frac{\pi_1^L}{\vdash \exists xF, \Gamma, I_1}$ (\exists) and $\vdash \exists xF, \Gamma, \exists yI_1$ (\exists)

$$\pi^{R} = \frac{\overset{\pi^{R}}{\downarrow}}{\vdash \forall y I_{1}^{\perp}, \Delta} \quad (\forall) \text{ we have that } \pi' = \frac{\vdash \exists x F, \Gamma, I \qquad \vdash \Pi^{\perp}, \Delta}{\vdash \exists x F, \Gamma, \Delta} \quad (\mathsf{Cut}) \text{ is a solved}$$

PRIS to which π cut-expands via a cut-commutation rule and a key $(\exists)/(\forall)$ rule:

= If y occurs in Δ but not in Γ , then we set $I = \forall y I_1, \pi^L = \frac{\vdash F\{y/x\}, \Gamma, I_1}{\vdash \exists x F, \Gamma, I_1}$ (\exists) and (\forall)

846

$$\pi^{R} = \frac{\overset{\pi^{R}_{1}}{\vdash I_{1}^{\perp}, \Delta}}{\vdash \exists y I_{1}^{\perp}, \Delta} \quad (\exists) \text{ One gets that } \pi' = \frac{\vdash \exists x F, \Gamma, \forall y I_{1} \quad \vdash \exists y I_{1}^{\perp}, \Delta}{\vdash \exists x F, \Gamma, \Delta} \quad (\mathsf{Cut}) \text{ is a solved PRIS to which } \pi \text{ cut-expands:}$$

848 solved PRIS to which π cut-expands:

$$\pi' \longrightarrow_{\rm cut} \frac{\vdash F\{y/x\}, \Gamma, I_1}{\vdash \exists xF, \Gamma, I_1} \quad (\exists) \qquad \vdash I_1^{\perp}, \Delta \\ \vdash \exists xF, \Gamma, \Delta \quad ({\rm Cut}) \qquad \longrightarrow_{\rm cut} \frac{\vdash F\{y/x\}, \Gamma, I_1 \qquad \vdash I_1^{\perp}, \Delta}{\vdash \exists xF, \Gamma, \Delta} \quad ({\rm Cut}) \qquad = \pi$$

E Appendix on interpolating system L (Section 5) 849

▶ Proposition 29. In what follows, t (resp. e, resp. c) is a normal L-term (resp. normal 850 L-context, resp. normal L-command). The following interpolating results hold: 851

1. If $c: (\Gamma_1, \Gamma_2 \vdash \Delta_1, \Delta_2)$, there exist a type $I \in \mathbb{V}oc(\Gamma_1, \Delta_1) \cap \mathbb{V}oc(\Gamma_2, \Delta_2)$ and t, e such 852 that $\Gamma_1 \vdash t : I \mid \Delta_1 \text{ and } \Gamma_2 \mid e : I \vdash \Delta_2, \text{ and } \langle t \mid e \rangle \longrightarrow^* c.$ 853

2. If $\Gamma_1, \Gamma_2 \vdash t : A \mid \Delta_1, \Delta_2$, there exist a type $I \in \mathbb{V}oc(\Gamma_1, \Delta_1, A) \cap \mathbb{V}oc(\Gamma_2, \Delta_2)$ and α, t', e' 854 such that $\Gamma_1 \vdash t' : A \mid \alpha : I, \Delta_1 \text{ and } \Gamma_2 \mid e' : I \vdash \Delta_2, \text{ and } t'\{e'/\alpha\} \longrightarrow^* t.$ 855

3. If $\Gamma_1, \Gamma_2 \mid e: A \vdash \Delta_1, \Delta_2$, there exist a type $I \in \mathbb{V}oc(\Gamma_1, \Delta_1, A) \cap \mathbb{V}oc(\Gamma_2, \Delta_2)$ and α, e', e'' 856 such that $\Gamma_1 \mid e' : A \mid \alpha : I, \Delta_1 \text{ and } \Gamma_2 \mid e'' : I \vdash \Delta_2, \text{ and } e'\{e''/\alpha\} \longrightarrow^* e.$ 857

4. If $\Gamma_1, \Gamma_2 \vdash t : A \mid \Delta_1, \Delta_2$, there exist a type $I \in \mathbb{V}oc(\Gamma_1, \Delta_1) \cap \mathbb{V}oc(\Gamma_2, \Delta_2, A)$ and x, t', t''858 such that $\Gamma_1 \vdash t'' : I \mid \Delta_1$ and $\Gamma_2, x : I \vdash t' : A \mid \Delta_2$, and $t'\{t''/x\} \longrightarrow^* t$. 859

5. If $\Gamma_1, \Gamma_2 \mid e : A \vdash \Delta_1, \Delta_2$, there exist a type $I \in \mathbb{V}oc(\Gamma_1, \Delta_1) \cap \mathbb{V}oc(\Gamma_2, \Delta_2, A)$ and x, t', e'860 such that $\Gamma_1 \vdash t' : I \mid \Delta_1$ and $\Gamma_2, x : I \mid e' : A \vdash \Delta_2$, and $e'\{t'/x\} \longrightarrow^* e$. 861

Proof sketch. The result is proved by mutual induction on the structure of terms, contexts 862 and commands. We treat only cases 1-3, cases 4 and 5 being essentially similar to cases 3 863 and 2 respectively. 864

1. If c is a command in normal form, it is either of the form $\langle z \mid e \rangle$ (with z being declared 865 in Γ_1 or Γ_2) or $\langle t \mid \beta \rangle$ (with α being declared in Δ_1 or Δ_2). Depending on the case and 866 whether the variable is declared in the left or right component of the typing contexts, we 867 apply induction hypotheses for one of cases 2–4. 868

 $c = \langle z \mid e \rangle$. Let us assume for instance that $c = \langle z \mid e \rangle$ and $\langle z \mid e \rangle : (\Gamma_1, \Gamma_2, z : A \vdash$ 869 Δ_1, Δ_2). 870

Then, e being structurally smaller than c, the induction hypothesis applied on Γ_1, Γ_2 871 $e: A \vdash \Delta_1, \Delta_2$, ensures the existence of a type $I \in \mathbb{V}oc(\Gamma_1, \Delta_1) \cap \mathbb{V}oc(\Gamma_2, \Delta_2, A)$ as 872 well as x, t', e' such that $\Gamma_1 \vdash t' : I \mid \Delta_1$ and $\Gamma_2, x : I \mid e' : A \vdash \Delta_2$, and $e'\{t'/x\} \longrightarrow^* e$. 873 Therefore $\langle t' \mid \tilde{\mu}x \langle z \mid e' \rangle \longrightarrow^{\star} c$ and we indeed have $\Gamma_1 \vdash t' : I \mid \Delta_1$ and $\Gamma_2, z : A \mid A$ 874 $\tilde{\mu}x.\langle z \mid e' \rangle : I \vdash \Delta_2.$ 875

On the other hand, if $\langle z \mid e \rangle : (\Gamma_1, \Gamma_2, z : A \vdash \Delta_1, \Delta_2)$, e being structurally smaller 876 than c, the induction hypothesis applied on $\Gamma_1, \Gamma_2 \mid e : A \vdash \Delta_1, \Delta_2$, ensures the 877 existence of a type $I \in \mathbb{V}oc(\Gamma_1, \Delta_1) \cap \mathbb{V}oc(\Gamma_2, \Delta_2, A)$ as well as α, e', e'' such that 878 $\Gamma_1 \mid e' : \mathbf{A} \vdash \alpha : \mathbf{I}, \Delta_1 \text{ and } \Gamma_2 \mid e'' : \mathbf{I} \vdash \Delta_2, \text{ and } e'\{e''/\alpha\} \longrightarrow^* e.$ 879

Therefore $\langle \mu \alpha . \langle z \mid e' \rangle \mid e'' \rangle \longrightarrow^{\star} c$ and we indeed have $\Gamma_1, z : A \vdash \mu \alpha . \langle z \mid e' \rangle : I \mid \Delta_1$ 880 and $\Gamma_2, z : A \mid e'' : I \vdash \Delta_2$. 881

 $c = \langle t \mid \beta \rangle$. The other case is similar:

- Let us assume that $c = \langle t \mid \beta \rangle$ and $\langle t \mid \beta \rangle : (\Gamma_1, \Gamma_2 \vdash \Delta_1, \Delta_2, \beta : A).$
- Then, t being structurally smaller than c, the induction hypothesis applied on $\Gamma_1, \Gamma_2 \vdash$
- $t: A \mid \Delta_1, \Delta_2, \text{ ensures the existence of a type } I \in \mathbb{V}oc(\Gamma_1, \Delta_1) \cap \mathbb{V}oc(\Gamma_2, \Delta_2, A) \text{ as well}$ as x, t', t'' such that $\Gamma_1 \vdash t'': I \mid \Delta_1$ and $\Gamma_2, x: I \vdash t': A \mid \Delta_2, \text{ and } t'\{t''/x\} \longrightarrow^* t.$
- Therefore $\langle t'' \mid \tilde{\mu}x.\langle t' \mid \beta \rangle \rangle \longrightarrow^* c$ and we indeed have $\Gamma_1 \vdash t'': I \mid \Delta_1$ and $\Gamma_2, z: A \mid c$
- Therefore $\langle t'' \mid \mu x. \langle t' \mid \beta \rangle \longrightarrow^* c$ and we indeed have $\Gamma_1 \vdash t'' : I \mid \Delta_1$ and $\Gamma_2, z : A$
- $\tilde{\mu}x.\langle t' \mid \beta \rangle : I \vdash \Delta_2.$
- The last case is similar.
- ⁸⁹⁰ 2. In that case, we make a case distinction based on the structure of t. We detail the λ case:
- If $t = \mu \alpha . c$ and its typing judgment has shape $\Gamma_1, \Gamma_2 \vdash \mu \alpha . c : A \mid \Delta_1, \Delta_2$; It follows that 891 c has typing judgment $\Gamma_1, \Gamma_2 \vdash \alpha : A, \Delta_1, \Delta_2$ and therefore by induction hypothesis, 892 there exist an interpolant type I, a term t and a context e such that $\Gamma_1 \vdash t : I \mid \alpha : A, \Delta_1$ 893 and $\Gamma_2 \mid e : I \vdash \Delta_2$ such that $\langle t \mid e \rangle \longrightarrow^* c$. 894 Let us then set I' = I, $t' = \mu \alpha \langle t \mid \beta \rangle$ and e' = e and one straightforwardly gets that 895 $t'\{e'/\beta\} \longrightarrow^{\star} \mu\alpha.c.$ 896 If $t = \lambda x \cdot u$ and its typing judgment has shape $\Gamma_1, \Gamma_2 \vdash \lambda x \cdot u : A \multimap B \mid \Delta_1, \Delta_2$. 897 By induction hypothesis we find an interpolant type I for u which can be used to 898 interpolate t as well: we have α, t', e' such that $\Gamma_1, x : A \vdash t' : B \mid \alpha : I, \Delta_1$ and 899 $\Gamma_2 \mid e' : I \vdash \Delta_2$, and $t'\{e'/\alpha\} \longrightarrow^* u$. 900 Therefore $\lambda x.t'\{e'/\alpha\} \longrightarrow^* \lambda x.u = t.$ 901 If t = (u, v) and its typing judgment has shape $\Gamma_1, \Gamma'_1, \Gamma_2, \Gamma'_2 \vdash (u, v) : A \otimes B \mid \Delta_1, \Delta'_1, \Delta_2, \Delta'_2$ 902 with $\Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2$ and $\Gamma'_1, \Gamma'_2 \vdash v : B \mid \Delta'_1, \Delta'_2$ 903 By induction hypothesis we find an interpolant type J for u and K for v that we can 904 combine to interpolate t as well: we have α, u', e' such that $\Gamma_1 \vdash u' : A \mid \alpha : J, \Delta_1$ and 905 $\Gamma_2 \mid e': J \vdash \Delta_2$, and $u'\{e'/\alpha\} \longrightarrow^* u$ and we have β, v', f' such that $\Gamma'_1 \vdash v': B \mid \beta$: 906 $K, \Delta'_1 \text{ and } \Gamma'_2 \mid f' : K \vdash \Delta'_2, \text{ and } v'\{f'/\alpha\} \longrightarrow^* v$. 907 Therefore we set $t' = \mu \gamma \langle \mu[\alpha, \beta] \langle (u', v') \mid \gamma \rangle \mid \delta \rangle$ and we have $t'\{[e', f']\} = t'$ 908 $\mu\gamma.\langle\mu[\alpha,\beta].\langle(u',v') \mid \gamma\rangle \mid \delta\rangle\{[e',f'])/\delta\} \longrightarrow^* \mu\gamma.\langle(u'\{e'/\alpha\},v'\{f'/\beta\}) \mid \gamma\rangle \longrightarrow^*$ 909 $\mu\gamma.\langle (u,v) \mid \gamma \rangle \longrightarrow_{\eta\mu} (u,v).$ 910 If $t = \mu[\alpha, \beta].c$ and its typing judgment has shape $\Gamma_1, \Gamma_2 \vdash \mu[\alpha, \beta].c : A \otimes B \mid \Delta_1, \Delta_2$ 911 with $c: (\Gamma_1, \Gamma_2 \vdash \alpha : A, \beta : B, \Delta_1, \Delta_2)$. By induction hypothesis, we find an interpolant 912 type $I \in \mathbb{V}oc(\Gamma_1, \Delta_1) \cap \mathbb{V}oc(\Gamma_2, \Delta_2)$ and u, e such that $\Gamma_1 \vdash u : I \mid \alpha : A, \beta : B, \Delta_1$ and 913 $\Gamma_2 \mid e : I \vdash \Delta_2$, and $\langle u \mid e \rangle \longrightarrow^* c$. 914 Therefore, we set $t' = \mu[\alpha, \beta] \langle u \mid \gamma \rangle$ and have $\Gamma_1 \vdash u' : A \otimes B \mid \Delta_1, \gamma : I$ and 915 $t'\{e/\gamma\} = \mu[\alpha, \beta]. \langle u \mid e \rangle \longrightarrow^* t.$ 916 If t = () and its typing judgment has shape $\vdash () : 1 \mid \text{We set } t' = \mu \beta \langle \mu \mid \langle () \mid \beta \rangle \mid \alpha \rangle$ 917 such that $\vdash t' : 1 \mid \alpha : \bot$ and e = [], which ensures that $\mid e : \bot \vdash$. 918 Therefore, we have: $t'\{e'/\alpha\} \longrightarrow \mu\beta.\langle () \mid \beta \rangle \longrightarrow_{\eta_{\mu}} t.$ 919 If $t = \mu[].c$ and its typing judgment has shape $\Gamma_1, \Gamma_2 \vdash \mu[].c : \perp \mid \Delta_1, \Delta_2$. In particular 920 $c: (\Gamma_1, \Gamma_2 \vdash \Delta_1, \Delta_2)$ so, by induction hypothesis there exist an interpolant type 921 $I \in \mathbb{V}oc(\Gamma_1, \Delta_1) \cap \mathbb{V}oc(\Gamma_2, \Delta_2)$ and u, e such that $\Gamma_1 \vdash u : I \mid \Delta_1$ and $\Gamma_2 \mid e : I \vdash \Delta_2$, 922 and $\langle u \mid e \rangle \longrightarrow^* c$. 923 Setting $t' = \mu[].\langle u \mid \alpha \rangle$, we have $t'\{e/\alpha\} = \mu[].\langle u \mid e \rangle \longrightarrow^* \mu[].c$ as expected. 924 If $t = \iota_j(u)$ and its typing judgment has shape $\Gamma_1, \Gamma_2 \vdash \iota_j(u) : A_1 \oplus A_2 \mid \Delta_1, \Delta_2$, we 925 have $\Gamma_1, \Gamma_2 \vdash u : A_i \mid \Delta_1, \Delta_2$, so that by induction hypothesis, we find an interpolant 926 type I, a term u', a context e and a context variable α such that $\Gamma_1 \vdash u' : A_i \mid \Delta_1, \alpha : I$ 927 and $\Gamma_2 \mid e : I \vdash \Delta_2$ with $u'\{e/\alpha\} \longrightarrow^* u$. 928

929		Setting $t' = \iota_j(u')$, we have $\Gamma_1 \vdash t' : A_1 \oplus A_2 \mid \Delta_1, \alpha : I$ and $t'\{e/\alpha\} = \iota_j(u'\{e/\alpha\}) \longrightarrow^*$
930		$\iota_j(u') = t.$
931	lf	$f t = \mu(\pi_1(\alpha) \mapsto c \mid \pi_2(\beta) \mapsto d)$ and its typing judgment has shape $\Gamma_1, \Gamma_2 \vdash \mu(\pi_1(\alpha) \mapsto d)$
932		$c \mid \pi_2(\beta) \mapsto d) : A \otimes B \mid \Delta_1, \Delta_2.$
933		We have $c: (\Gamma_1, \Gamma_2 \vdash \alpha : A, \Delta_1, \Delta_2)$ and $d: (\Gamma_1, \Gamma_2 \vdash \beta : B, \Delta_1, \Delta_2)$. By applying
934		twice the induction hypothesis, we get:
935		I, J satisfying the interpolation constraints wrt c and d ;
936		$= \Gamma_1 \vdash u : I \mid \alpha : A, \Delta_1;$
937		$= \Gamma_1 \vdash v : J \mid \beta : B, \Delta_1;$
938		$\Gamma_2 \mid e: I \vdash \Delta_2$ and
939		$= \Gamma_2 \mid f: J \vdash \Delta_2$
940		such that $\langle u \mid e \rangle \longrightarrow^* c$ and $\langle v \mid f \rangle \longrightarrow^* d$.
941		Since $\langle x \mid e \rangle$: $(\Gamma_2, x : I \vdash \Delta_2)$ and $\langle y \mid f \rangle$: $(\Gamma_2, y : J \vdash \Delta_2)$, then by setting
942		$e' = \tilde{\mu}(\iota_1(x) \mapsto \langle x \mid e \rangle \mid \iota_2(y) \mapsto \langle y \mid f \rangle) \text{ and } t' = \mu(\pi_1(\alpha) \mapsto \langle \iota_1(u) \mid \gamma \rangle \mid \pi_2(\beta) \mapsto dt'$
943		$\langle \iota_1(v) \mid \gamma \rangle$, we have: $\Gamma_1 \vdash t' : A \otimes B \mid \Delta_1, \gamma : I \oplus J \text{ and } \Gamma_2 \mid e : I \oplus J \vdash \Delta_2$ so that
944		$u'\{e'/\gamma\} \longrightarrow \mu(\pi_1(\alpha) \mapsto \langle \iota_1(u) \mid e'\rangle \mid \pi_2(\beta) \mapsto \langle \iota_2(v) \mid e'\rangle) \longrightarrow \mu(\pi_1(\alpha) \mapsto \langle u \mid e\rangle \mid a)$
945		$\pi_2(\beta) \mapsto \langle v \mid f \rangle) \longrightarrow^* \mu(\pi_1(\alpha) \mapsto c \mid \pi_2(\beta) \mapsto d)$
946	lf	$\Gamma t = \text{tp}$ and its typing judgment has shape $\Gamma_1, \Gamma_2 \vdash \text{tp} : \top \mid \Delta_1, \Delta_2$.
947		One also has $\Gamma_1, \Gamma_2 \vdash tp : \top \mid \Delta_1, \Delta_2, \alpha : 0.$
948		Then for $I = 0$ and $e = \text{stop}$, we have $t\{e/\alpha\} = t$.
	з т	he context case is symmetrical to the previous one, due to the symmetry of linear typing
949	J. 1 Т	the only case to consider is that of the applicative context, all the other case can be
950	r d	irectly retrieve from the previous item dualizing terms and contexts
111-1	u	
951		
951 952	lf	$f = u \cdot f$ and its typing judgment has shape $\Gamma_1, \Gamma'_1, \Gamma_2, \Gamma'_2 \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta'_1, \Delta_2, \Delta'_2$.
951 952 953	lf	$f = u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma'_1, \Gamma_2, \Gamma'_2 \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta'_1, \Delta_2, \Delta'_2.$ Then we have $\Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma'_1, \Gamma'_2 \mid f : B \vdash \Delta'_1, \Delta'_2.$
951 952 953 954	lf	$f = u \cdot f$ and its typing judgment has shape $\Gamma_1, \Gamma'_1, \Gamma_2, \Gamma'_2 \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta'_1, \Delta_2, \Delta'_2$. Then we have $\Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2$ and $\Gamma'_1, \Gamma'_2 \mid f : B \vdash \Delta'_1, \Delta'_2$. By induction hypothesis we find an interpolant type J , together with u', g, α for u and
951 952 953 954 955	lf	$f e = u \cdot f$ and its typing judgment has shape $\Gamma_1, \Gamma'_1, \Gamma_2, \Gamma'_2 \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta'_1, \Delta_2, \Delta'_2$. Then we have $\Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2$ and $\Gamma'_1, \Gamma'_2 \mid f : B \vdash \Delta'_1, \Delta'_2$. By induction hypothesis we find an interpolant type J , together with u', g, α for u and an interpolant type K , together with f', f'', β for f which satisfy:
951 952 953 954 955 956	If	$\begin{split} f &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma_1', \Gamma_2, \Gamma_2' \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta_1', \Delta_2, \Delta_2'. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma_1', \Gamma_2' \mid f : B \vdash \Delta_1', \Delta_2'. \\ \text{By induction hypothesis we find an interpolant type } J, \text{ together with } u', g, \alpha \text{ for } u \text{ and} \\ \text{an interpolant type } K, \text{ together with } f', f'', \beta \text{ for } f \text{ which satisfy:} \\ &= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \end{split}$
951 952 953 954 955 956 957	If	$\begin{split} f &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma_1', \Gamma_2, \Gamma_2' \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta_1', \Delta_2, \Delta_2'. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma_1', \Gamma_2' \mid f : B \vdash \Delta_1', \Delta_2'. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy:&= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \end{split}$
951 952 953 954 955 956 957 958	If	$\begin{split} f &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma_1', \Gamma_2, \Gamma_2' \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta_1', \Delta_2, \Delta_2'. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma_1', \Gamma_2' \mid f : B \vdash \Delta_1', \Delta_2'. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy:&= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \\ &= \Gamma_1' \mid f' : B \vdash \Delta_1', \beta : K; \end{split}$
951 952 953 954 955 956 957 958 959	If	$\begin{split} \vec{F} &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma_1', \Gamma_2, \Gamma_2' \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta_1', \Delta_2, \Delta_2'. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma_1', \Gamma_2' \mid f : B \vdash \Delta_1', \Delta_2'. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy:&= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \\ &= \Gamma_1' \mid f' : B \vdash \Delta_1', \beta : K; \\ &= \Gamma_2' \mid f'' : K \vdash \Delta_2'; \end{split}$
951 952 953 954 955 956 957 958 959 960	If	$\begin{aligned} f &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma_1', \Gamma_2, \Gamma_2' \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta_1', \Delta_2, \Delta_2'. \end{aligned}$ Then we have $\Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma_1', \Gamma_2' \mid f : B \vdash \Delta_1', \Delta_2'. \end{aligned}$ By induction hypothesis we find an interpolant type J , together with u', g, α for u and an interpolant type K , together with f', f'', β for f which satisfy: $\begin{aligned} &= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \\ &= \Gamma_1' \mid f' : B \vdash \Delta_1', \beta : K; \\ &= \Gamma_2' \mid f'' : K \vdash \Delta_2'; \\ &= u' \{g/\alpha\} \longrightarrow^* u; \end{aligned}$
951 952 953 954 955 956 957 958 959 960 961	If	$\begin{split} f &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma_1', \Gamma_2, \Gamma_2' \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta_1', \Delta_2, \Delta_2'. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma_1', \Gamma_2' \mid f : B \vdash \Delta_1', \Delta_2'. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy:&= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \\ &= \Gamma_1' \mid f' : B \vdash \Delta_1', \beta : K; \\ &= \Gamma_2' \mid f'' : K \vdash \Delta_2'; \\ &= u' \{g/\alpha\} \longrightarrow^* u; \\ &= f' \{f''/\beta\} \longrightarrow^* f. \end{split}$
951 952 953 954 955 956 957 958 959 960 961 962	If	$\begin{split} f &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma_1', \Gamma_2, \Gamma_2' \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta_1', \Delta_2, \Delta_2'. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma_1', \Gamma_2' \mid f : B \vdash \Delta_1', \Delta_2'. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy:&= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \\ &= \Gamma_1' \mid f' : B \vdash \Delta_1', \beta : K; \\ &= \Gamma_2' \mid f'' : K \vdash \Delta_2'; \\ &= u'\{g/\alpha\} \longrightarrow^* u; \\ &= f'\{f''/\beta\} \longrightarrow^* f. \\ \text{Therefore, we have } \Gamma_1, \Gamma_1', \mid u' \cdot f' : A \multimap B \vdash \Delta_1, \Delta_1', \alpha : J, \beta : K \text{ and } \Gamma_2, \Gamma_2' \mid [g, f''] : \end{split}$
951 952 953 954 955 956 957 958 959 960 961 962 962	If	$\begin{split} f &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma_1', \Gamma_2, \Gamma_2' \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta_1', \Delta_2, \Delta_2'. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma_1', \Gamma_2' \mid f : B \vdash \Delta_1', \Delta_2'. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy:&= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \\ &= \Gamma_1' \mid f' : B \vdash \Delta_1', \beta : K; \\ &= \Gamma_2' \mid f'' : K \vdash \Delta_2'; \\ &= u' \{g/\alpha\} \longrightarrow^* u; \\ &= f' \{f''/\beta\} \longrightarrow^* f. \\ \text{Therefore, we have } \Gamma_1, \Gamma_1', \mid u' \cdot f' : A \multimap B \vdash \Delta_1, \Delta_1', \alpha : J, \beta : K \text{ and } \Gamma_2, \Gamma_2' \mid [g, f''] : J \otimes K \vdash \Delta_2, \Delta_2'. \end{split}$
951 952 953 954 955 956 957 958 959 960 961 962 963 964	If	$\begin{split} f &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma_1', \Gamma_2, \Gamma_2' \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta_1', \Delta_2, \Delta_2'. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma_1', \Gamma_2' \mid f : B \vdash \Delta_1', \Delta_2'. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy:&= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \\ &= \Gamma_1' \mid f' : B \vdash \Delta_1', \beta : K; \\ &= \Gamma_2' \mid f'' : K \vdash \Delta_2'; \\ &= u' \{g/\alpha\} \longrightarrow^* u; \\ &= f' \{f''/\beta\} \longrightarrow^* f. \\ \text{Therefore, we have } \Gamma_1, \Gamma_1' \mid u' \cdot f' : A \multimap B \vdash \Delta_1, \Delta_1', \alpha : J, \beta : K \text{ and } \Gamma_2, \Gamma_2' \mid [g, f''] : \\ J \otimes K \vdash \Delta_2, \Delta_2'. \\ \text{As a consequence, we have } \Gamma_1, \Gamma_1' \mid \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle : A \multimap B \vdash \Delta_1, \Delta_1', \gamma : \\ \end{bmatrix}$
951 952 953 954 955 956 957 958 959 960 961 962 963 964	If	$\begin{split} f &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma'_1, \Gamma_2, \Gamma'_2 \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta'_1, \Delta_2, \Delta'_2. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma'_1, \Gamma'_2 \mid f : B \vdash \Delta'_1, \Delta'_2. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy:&= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \\ &= \Gamma'_1 \mid f' : B \vdash \Delta'_1, \beta : K; \\ &= \Gamma'_2 \mid f'' : K \vdash \Delta'_2; \\ &= u' \{g/\alpha\} \longrightarrow^* u; \\ &= f' \{f''/\beta\} \longrightarrow^* f. \\ \text{Therefore, we have } \Gamma_1, \Gamma'_1, \mid u' \cdot f' : A \multimap B \vdash \Delta_1, \Delta'_1, \alpha : J, \beta : K \text{ and } \Gamma_2, \Gamma'_2 \mid [g, f''] : J \otimes K \vdash \Delta_2, \Delta'_2. \\ \text{As a consequence, we have } \Gamma_1, \Gamma'_1 \mid \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle : A \multimap B \vdash \Delta_1, \Delta'_1, \gamma : J \otimes K \\ We find that for the derivative for a field of the derivative for the fie$
951 952 953 955 955 955 955 958 959 960 961 962 963 964 965 966	If	$\begin{split} F &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma'_1, \Gamma_2, \Gamma'_2 \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta'_1, \Delta_2, \Delta'_2. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma'_1, \Gamma'_2 \mid f : B \vdash \Delta'_1, \Delta'_2. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy:&= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \\ &= \Gamma'_1 \mid f' : B \vdash \Delta'_1, \beta : K; \\ &= \Gamma'_2 \mid f'' : K \vdash \Delta'_2; \\ &= u' \{g/\alpha\} \longrightarrow^* u; \\ &= f' \{f''/\beta\} \longrightarrow^* f. \\ \text{Therefore, we have } \Gamma_1, \Gamma'_1, \mid u' \cdot f' : A \multimap B \vdash \Delta_1, \Delta'_1, \alpha : J, \beta : K \text{ and } \Gamma_2, \Gamma'_2 \mid [g, f''] : J \otimes K \vdash \Delta_2, \Delta'_2. \\ \text{As a consequence, we have } \Gamma_1, \Gamma'_1 \mid \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle : A \multimap B \vdash \Delta_1, \Delta'_1, \gamma : J \otimes K \\ \text{We thus set } I = J \otimes K, e' = \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle \text{ and } e'' = [g, f''] \text{ and we observe} \end{split}$
951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 966	If	$\begin{aligned} F &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma'_1, \Gamma_2, \Gamma'_2 \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta'_1, \Delta_2, \Delta'_2. \end{aligned}$ Then we have $\Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma'_1, \Gamma'_2 \mid f : B \vdash \Delta'_1, \Delta'_2. \end{aligned}$ By induction hypothesis we find an interpolant type J , together with u', g, α for u and an interpolant type K , together with f', f'', β for f which satisfy: $&= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \\ &= \Gamma'_1 \mid f' : B \vdash \Delta'_1, \beta : K; \\ &= \Gamma'_2 \mid f'' : K \vdash \Delta'_2; \\ &= u' \{g/\alpha\} \longrightarrow^* u; \\ &= f'\{f''/\beta\} \longrightarrow^* f. \end{aligned}$ Therefore, we have $\Gamma_1, \Gamma'_1, \mid u' \cdot f' : A \multimap B \vdash \Delta_1, \Delta'_1, \alpha : J, \beta : K \text{ and } \Gamma_2, \Gamma'_2 \mid [g, f''] : J \otimes K \vdash \Delta_2, \Delta'_2. \end{aligned}$ As a consequence, we have $\Gamma_1, \Gamma'_1 \mid \tilde{\mu}x.\langle \mu[\alpha, \beta].\langle x \mid u' \cdot f' \rangle \mid \gamma \rangle : A \multimap B \vdash \Delta_1, \Delta'_1, \gamma : J \otimes K \\$ We thus set $I = J \otimes K, e' = \tilde{\mu}x.\langle \mu[\alpha, \beta].\langle x \mid u' \cdot f' \rangle \mid \gamma \rangle$ and $e'' = [g, f'']$ and we observe that
951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968	If	$\begin{split} \vec{r} &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma_1', \Gamma_2, \Gamma_2' \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta_1', \Delta_2, \Delta_2'. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma_1', \Gamma_2' \mid f : B \vdash \Delta_1', \Delta_2'. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy:&= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \\ &= \Gamma_1' \mid f' : B \vdash \Delta_1', \beta : K; \\ &= \Gamma_2' \mid f'' : K \vdash \Delta_2'; \\ &= u'\{g/\alpha\} \longrightarrow^* u; \\ &= f'\{f''/\beta\} \longrightarrow^* f. \\ \text{Therefore, we have } \Gamma_1, \Gamma_1', \mid u' \cdot f' : A \multimap B \vdash \Delta_1, \Delta_1', \alpha : J, \beta : K \text{ and } \Gamma_2, \Gamma_2' \mid [g, f''] : J \otimes K \vdash \Delta_2, \Delta_2'. \\ \text{As a consequence, we have } \Gamma_1, \Gamma_1' \mid \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle : A \multimap B \vdash \Delta_1, \Delta_1', \gamma : J \otimes K \\ \text{We thus set } I = J \otimes K, e' = \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle \text{ and } e'' = [g, f''] \text{ and we observe that} \\ e'\{e''/\gamma\} \longrightarrow^* \tilde{\mu}x. \langle x \mid u'\{g/\alpha\} \cdot f'\{f''/\beta\} \longrightarrow^* \tilde{\mu}x. \langle x \mid u \cdot f \rangle \longrightarrow_{\eta_{\bar{\mu}}} u \cdot f = e. \\ \end{aligned}$
951 952 953 955 956 957 958 959 960 961 962 963 964 965 966 966 967 968	If ,	$\begin{split} \vec{r} &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma_1', \Gamma_2, \Gamma_2' \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta_1', \Delta_2, \Delta_2'. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma_1', \Gamma_2' \mid f : B \vdash \Delta_1', \Delta_2'. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy:&= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \\ &= \Gamma_1' \mid f' : B \vdash \Delta_1', \beta : K; \\ &= \Gamma_2' \mid f'' : K \vdash \Delta_2'; \\ &= u' \{g/\alpha\} \longrightarrow^* u; \\ &= f' \{f''/\beta\} \longrightarrow^* f. \\ \text{Therefore, we have } \Gamma_1, \Gamma_1', \mid u' \cdot f' : A \multimap B \vdash \Delta_1, \Delta_1', \alpha : J, \beta : K \text{ and } \Gamma_2, \Gamma_2' \mid [g, f''] : J \otimes K \vdash \Delta_2, \Delta_2'. \\ \text{As a consequence, we have } \Gamma_1, \Gamma_1' \mid \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle : A \multimap B \vdash \Delta_1, \Delta_1', \gamma : J \otimes K \\ \text{We thus set } I = J \otimes K, e' = \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle \text{ and } e'' = [g, f''] \text{ and we observe that} \\ e' \{e''/\gamma\} \longrightarrow^* \tilde{\mu}x. \langle x \mid u' \{g/\alpha\} \cdot f' \{f''/\beta\} \longrightarrow^* \tilde{\mu}x. \langle x \mid u \cdot f \rangle \longrightarrow_{\eta_{\tilde{\mu}}} u \cdot f = e. \\ \\ \text{The other cases are similar to cases already treated in item 2. \\ \end{aligned}$
951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 966 966 966 969 970	If 4. T	$\begin{split} \vec{r} &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma_1', \Gamma_2, \Gamma_2' \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta_1', \Delta_2, \Delta_2'. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma_1', \Gamma_2' \mid f : B \vdash \Delta_1', \Delta_2'. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy:&= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \\ &= \Gamma_1' \mid f' : B \vdash \Delta_1', \beta : K; \\ &= \Gamma_2' \mid f'' : K \vdash \Delta_2'; \\ &= u'\{g/\alpha\} \longrightarrow^* u; \\ &= f'\{f''/\beta\} \longrightarrow^* f. \\ \text{Therefore, we have } \Gamma_1, \Gamma_1', \mid u' \cdot f' : A \multimap B \vdash \Delta_1, \Delta_1', \alpha : J, \beta : K \text{ and } \Gamma_2, \Gamma_2' \mid [g, f''] : J \otimes K \vdash \Delta_2, \Delta_2'. \\ \text{As a consequence, we have } \Gamma_1, \Gamma_1' \mid \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle : A \multimap B \vdash \Delta_1, \Delta_1', \gamma : J \otimes K \\ \text{We thus set } I = J \otimes K, e' = \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle \text{ and } e'' = [g, f''] \text{ and we observe that} \\ e'\{e''/\gamma\} \longrightarrow^* \tilde{\mu}x. \langle x \mid u'\{g/\alpha\} \cdot f'\{f''/\beta\} \longrightarrow^* \tilde{\mu}x. \langle x \mid u \cdot f \rangle \longrightarrow_{\eta_{\tilde{\mu}}} u \cdot f = e. \\ \text{The other cases are similar to cases already treated in item 2. \\ \text{hese cases are treated similarly to 2, but for the fact that the interpolant lives in the \\ \end{array}$
951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971	If 4. T w	$\begin{split} \hat{f} & e = u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma_1', \Gamma_2, \Gamma_2' \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta_1', \Delta_2, \Delta_2'. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma_1', \Gamma_2' \mid f : B \vdash \Delta_1', \Delta_2'. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy: = \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ = \Gamma_2 \mid g : J \vdash \Delta_2; \\ = \Gamma_1' \mid f' : B \vdash \Delta_1', \beta : K; \\ = \Gamma_2' \mid f'' : K \vdash \Delta_2'; \\ = u' \{g/\alpha\} \longrightarrow^* u; \\ = f'\{f''/\beta\} \longrightarrow^* f. \\ \text{Therefore, we have } \Gamma_1, \Gamma_1' \mid u' \cdot f' : A \multimap B \vdash \Delta_1, \Delta_1', \alpha : J, \beta : K \text{ and } \Gamma_2, \Gamma_2' \mid [g, f''] : J \otimes K \vdash \Delta_2, \Delta_2'. \\ \text{As a consequence, we have } \Gamma_1, \Gamma_1' \mid \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle : A \multimap B \vdash \Delta_1, \Delta_1', \gamma : J \otimes K \\ \text{We thus set } I = J \otimes K, e' = \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle \text{ and } e'' = [g, f''] \text{ and we observe that} \\ e'\{e''/\gamma\} \longrightarrow^* \tilde{\mu}x. \langle x \mid u'\{g/\alpha\} \cdot f'\{f''/\beta\} \rangle \longrightarrow^* \tilde{\mu}x. \langle x \mid u \cdot f \rangle \longrightarrow_{\eta_{\tilde{\mu}}} u \cdot f = e. \\ The other cases are similar to cases already treated in item 2. \\ \\ \text{hese cases are treated similarly to 2, but for the fact that the interpolant lives in the orld of terms rather than of contexts but since we did not use connective \multimap to build an$
951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 966 967 968 969 970 971 972	If 4. T wir	$\begin{split} \hat{f} & e = u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma_1', \Gamma_2, \Gamma_2' \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta_1', \Delta_2, \Delta_2'. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma_1', \Gamma_2' \mid f : B \vdash \Delta_1', \Delta_2'. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy: = \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ = \Gamma_2 \mid g : J \vdash \Delta_2; \\ = \Gamma_1' \mid f' : B \vdash \Delta_1', \beta : K; \\ = \Gamma_2' \mid f'' : K \vdash \Delta_2'; \\ = u' \{g/\alpha\} \longrightarrow^* u; \\ = f'\{f''/\beta\} \longrightarrow^* f. \\ \text{Therefore, we have } \Gamma_1, \Gamma_1' \mid u' \cdot f' : A \multimap B \vdash \Delta_1, \Delta_1', \alpha : J, \beta : K \text{ and } \Gamma_2, \Gamma_2' \mid [g, f''] : J \otimes K \vdash \Delta_2, \Delta_2'. \\ \text{As a consequence, we have } \Gamma_1, \Gamma_1' \mid \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle : A \multimap B \vdash \Delta_1, \Delta_1', \gamma : J \otimes K \\ \text{We thus set } I = J \otimes K, e' = \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle \text{ and } e'' = [g, f''] \text{ and we observe that} \\ e'\{e''/\gamma\} \longrightarrow^* \tilde{\mu}x. \langle x \mid u'\{g/\alpha\} \cdot f'\{f''/\beta\} \rangle \longrightarrow^* \tilde{\mu}x. \langle x \mid u \cdot f \rangle \longrightarrow_{\eta_{\tilde{\mu}}} u \cdot f = e. \\ \\ \text{The other cases are similar to cases already treated in item 2. \\ \\ \text{hese cases are treated similarly to 2, but for the fact that the interpolant lives in the orld of terms rather than of contexts but since we did not use connective \multimap to build an atterpolant in 2, the symmetry of system L does the job. \\ \end{cases}$
951 952 953 955 955 956 957 958 959 960 961 962 963 964 965 966 966 967 966 967 968 969 970 971 972 973	If 4. T w 5. T	$\begin{split} \mathbf{r} &= u \cdot f \text{ and its typing judgment has shape } \Gamma_1, \Gamma'_1, \Gamma_2, \Gamma'_2 \mid u \cdot f : A \multimap B \vdash \Delta_1, \Delta'_1, \Delta_2, \Delta'_2. \\ \text{Then we have } \Gamma_1, \Gamma_2 \vdash u : A \mid \Delta_1, \Delta_2 \text{ and } \Gamma'_1, \Gamma'_2 \mid f : B \vdash \Delta'_1, \Delta'_2. \\ \text{By induction hypothesis we find an interpolant type } J, together with u', g, \alpha for u and an interpolant type K, together with f', f'', \beta for f which satisfy:&= \Gamma_1 \vdash u' : A \mid \Delta_1, \alpha : J; \\ &= \Gamma_2 \mid g : J \vdash \Delta_2; \\ &= \Gamma'_1 \mid f' : B \vdash \Delta'_1, \beta : K; \\ &= \Gamma'_2 \mid f'' : K \vdash \Delta'_2; \\ &= u' \{g/\alpha\} \longrightarrow^* u; \\ &= f'\{f''/\beta\} \longrightarrow^* f. \\ \text{Therefore, we have } \Gamma_1, \Gamma'_1, \mid u' \cdot f' : A \multimap B \vdash \Delta_1, \Delta'_1, \alpha : J, \beta : K \text{ and } \Gamma_2, \Gamma'_2 \mid [g, f''] : J \approx K \vdash \Delta_2, \Delta'_2. \\ \text{As a consequence, we have } \Gamma_1, \Gamma'_1 \mid \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle : A \multimap B \vdash \Delta_1, \Delta'_1, \gamma : J \approx K \\ \text{We thus set } I = J \approx K, e' = \tilde{\mu}x. \langle \mu[\alpha, \beta]. \langle x \mid u' \cdot f' \rangle \mid \gamma \rangle \text{ and } e'' = [g, f''] \text{ and we observe that} \\ e'\{e''/\gamma\} \longrightarrow^* \tilde{\mu}x. \langle x \mid u'\{g/\alpha\} \cdot f'\{f''/\beta\} \rangle \longrightarrow^* \tilde{\mu}x. \langle x \mid u \cdot f \rangle \longrightarrow_{\eta_{\tilde{\mu}}} u \cdot f = e. \\ The other cases are similar to cases already treated in item 2. \\ \text{hese cases are treated similarly to 2, but for the fact that the interpolant lives in the orld of terms rather than of contexts but since we did not use connective \multimap to build an tterpolant in 2, the symmetry of system L does the job. \\ \text{hese cases are treated similarly to 3, but for the fact that the interpolant lives in the orld of terms rather than of contexts but since we did not use connective \multimap to build an tterpolant in 2, the symmetry of system L does the job. \\ \text{hese cases are treated similarly to 3, but for the fact that the interpolant lives in the orld of terms rather than of contexts but since we did not use connective \multimap to build an tterpolant in 2, the symmetry of system L does the job. \\ \text{hese cases are treated similarly to 3, but for the fact that the interpolant lives in the orld of terms rather than of contexts but since we did not use connective \multimap to build an tterpolant in 2, the symmetry of system L does the job. \\ \text{hese $

975 applies.

XX:38 Interpolation as cut-introduction

976

977	С	ontents	
978 979	1	"– Why Not a Proof-Relevant Interpolation Theorem? – Introduce Cuts, Of Course!"	1
980	2	Background on LL proof theory	3
981	3	Proof-relevant interpolation theorem	4
982 983 984 985 986 987 988	4	Interpolation as cut-introduction4.1Refining Maehara's method	5 6 7 8 10 11
989	5	On the computational significance of the result: Interpolating System L	13
990	6	Conclusion	
991	\mathbf{A}	LL cut-elimination reduction	
992	в	Full proof of proof-relevant interpolation for LL refining Maehara's method	
993 994	С	C Details on linear embeddings of LJ and LK and proof-relevant interpolation for LJ and LK.	
995 996	D	Details on interpolation as cut-introduction D.1 Details on the Proof of the Main Lemma	29 30
997	\mathbf{E}	Appendix on interpolating system L (Section 5)	35