

q-Rook placements and Jordan forms of upper-triangular nilpotent matrices

Martha Yip

University of Pennsylvania, 209 South 33rd St., Philadelphia PA 19104, USA

Abstract. The set of n by n upper-triangular nilpotent matrices with entries in a finite field \mathbb{F}_q has Jordan canonical forms indexed by partitions $\lambda \vdash n$. We study a connection between these matrices and non-attacking q -rook placements, which leads to a combinatorial formula for the number $F_\lambda(q)$ of matrices of fixed Jordan type as a weighted sum over rook placements.

Résumé. L'ensemble des matrices triangulaires supérieures nilpotentes d'ordre n sur un corps fini \mathbb{F}_q a des formes canoniques de Jordan indexées par les partitions $\lambda \vdash n$. Nous étudions une connexion entre ces matrices et les placements de tours, et nous présentons une formule combinatoire pour le nombre $F_\lambda(q)$ des matrices comme une somme sur les placements de tours.

Keywords: q -rook placements, Jordan canonical form, nilpotent matrices, set partitions

1 Introduction

In the beautiful paper *Variations on the Triangular Theme*, Kirillov (1995) studied various structures on the set of triangular matrices. Denote by $G_n(\mathbb{F}_q)$, the group of n by n upper-triangular matrices over the field \mathbb{F}_q having q elements, and let $\mathfrak{g}_n(\mathbb{F}_q) = \text{Lie}(G_n(\mathbb{F}_q))$ denote the corresponding Lie algebra of n by n upper-triangular nilpotent matrices over \mathbb{F}_q . It is known, for example, that the conjugacy classes of $G_n(\mathbb{F}_q)$ are in bijection with the adjoint orbits in $\mathfrak{g}_n(\mathbb{F}_q)$. To study the adjoint orbits we consider the Jordan canonical form. Each matrix $X \in \mathfrak{g}_n(\mathbb{F}_q)$ is similar to a block diagonal matrix consisting of elementary Jordan blocks with eigenvalue zero:

$$J_i = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ 0 & 0 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & 0 & \cdots & 0 & 0 \end{pmatrix}_{i \times i} .$$

If the Jordan canonical form of X has block sizes $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n \geq 0$, then X is said to have *Jordan type* λ , where λ is a partition of n . The Jordan type of X depends only on its adjoint orbit, so the similarity classes of nilpotent matrices are indexed by the partitions of n .

Let $\mathfrak{g}_{n,\lambda}(\mathbb{F}_q) \subseteq \mathfrak{g}_n(\mathbb{F}_q)$ be the set of matrices of fixed Jordan type λ . It was shown by Springer that $\mathfrak{g}_{n,\lambda}(\mathbb{F}_q)$ an algebraic manifold with f^λ irreducible components, each of which has the same dimension $\binom{n}{2} - n_\lambda$. Here, f^λ is the number of standard Young tableaux of shape λ , and n_λ is given in Equation 9. Let

$$F_\lambda(q) = |\mathfrak{g}_{n,\lambda}(\mathbb{F}_q)| \tag{1}$$

be the number of matrices of Jordan type λ . We note that $\sum_{\lambda \vdash n} F_\lambda(q) = |\mathfrak{g}_n(\mathbb{F}_q)| = q^{\binom{n}{2}}$. The cases $F_{(1^n)}(q) = 1$ and $F_{(n)}(q) = (q-1)^{n-1} q^{\binom{n-1}{2}}$ are readily computed, since the matrix of Jordan type (1^n) is the matrix of rank zero, and the matrices of Jordan type (n) are the matrices of rank $n-1$.

In **Section 2**, we present a simple recurrence equation for $F_\lambda(q)$ (see Proposition 2). As a consequence of the recurrence equation, it follows that $F_\lambda(q)$ is a polynomial in q with nonnegative integer coefficients, $\deg F_\lambda(q) = \binom{n}{2} - n_\lambda$, and the coefficient of the highest degree term in $F_\lambda(q)$ is f^λ .

A connection with q -Rook placements

In their study of a formula of Frobenius, Garsia and Remmel (1986) introduced the q -rook polynomial

$$R_k(q, B) = \sum_{c \in \mathcal{C}(B, k)} q^{\text{inv}(c)}, \tag{2}$$

which is a sum over the set $\mathcal{C}(B, k)$ of non-attacking placements of k rooks on the Ferrers board B , and $\text{inv}(c)$, defined in Equation 11, is the number of inversions of c . In the case when $B = \delta_n$ is the staircase-shaped board, Garsia and Remmel showed that $R_k(q, \delta_n) = S_{n, n-k}(q)$ is a q -Stirling number of the second kind. These numbers are defined by the recurrence

$$S_{n,k}(q) = q^{k-1} S_{n-1, k-1}(q) + [k]_q S_{n-1, k}(q) \quad \text{for } 0 \leq k \leq n, \tag{3}$$

with initial conditions $S_{0,0}(q) = 1$, and $S_{n,k}(q) = 0$ for $k < 0$ or $k > n$.

It was shown by Solomon (1990) that non-attacking placements of k rooks on rectangular $m \times n$ boards are naturally associated to m by n matrices with rank k over \mathbb{F}_q . By identifying a Ferrers board B inside an n by n grid with the entries of an n by n matrix, Haglund (1998) generalized Solomon's result to the case of non-attacking placements of k rooks on Ferrers boards, and obtained a formula for the number of n by n matrices with rank k whose support is contained in the Ferrers board region. As a special case of Haglund's formula, the number of nilpotent matrices of rank k is

$$P_k(q) = (q-1)^k q^{\binom{n}{2}-k} R_k(q^{-1}, \delta_n). \tag{4}$$

Now, a matrix in $\mathfrak{g}_{n,\lambda}(\mathbb{F}_q)$ has rank $n - \ell(\lambda)$, where $\ell(\lambda)$ is the number of parts of λ , so the number of matrices in $\mathfrak{g}_n(\mathbb{F}_q)$ with rank k is

$$P_k(q) = \sum_{\lambda \vdash n: \ell(\lambda)=n-k} F_\lambda(q). \tag{5}$$

Given Equations 4 and 5, it is natural to ask whether it is possible to partition the placements $\mathcal{C}(\delta_n, k)$ into disjoint subsets so that the sum over each subset of placements gives $F_\lambda(q)$. The goal of this paper is to study the connection between upper-triangular nilpotent matrices over \mathbb{F}_q and non-attacking q -rook placements on the staircase-shaped board δ_n .

Section 3 forms the heart of this paper. Inspired by Equation 7 for $F_\lambda(q)$, we define a graph \mathcal{Z} closely related to Young’s lattice. The main result is Theorem 9, which states that there is a weight-preserving bijection between rook placements and paths in \mathcal{Z} . As a result, we obtain a formula for $F_\lambda(q)$ as a sum over certain weighted q -rook placements (see Corollary 10), which can be viewed as a generalization of Haglund’s formula in Equation 4.

There is a well-known bijection between rook placements on the staircase-shaped board δ_n with k rooks, and set partitions of $\{1, \dots, n\}$ with $n - k$ blocks. In **Section 4**, we describe how paths in \mathcal{Z} gives a new bijection between these sets, and how this gives a definition of a lattice of compositions which appears to be new. Finally, in **Section 5**, we mention some further problems to pursue. In this article, the proofs are either omitted or briefly sketched. Full details can be found in the preprint Yip (2013).

2 The recurrence equation for $F_\lambda(q)$

We define a *partition* λ of a nonnegative integer n , denoted by $\lambda \vdash n$, is a non-increasing sequence of nonnegative integers $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0$ with $|\lambda| = \sum_{i=1}^n \lambda_i = n$. If λ has k positive parts, write $\ell(\lambda) = k$. Represent a partition λ by its Ferrers diagram in the English notation, which is an array of λ_i boxes in row i , with the boxes justified upwards and to the left. Let λ'_j denote the size of the j th column of λ .

Example 1 *The partition*

$$\lambda = (4, 2, 2, 1) \vdash 9 \quad \text{has diagram} \quad \begin{array}{cccc} \square & \square & \square & \square \\ \square & \square & & \\ \square & \square & & \\ \square & & & \end{array}$$

and columns $\lambda'_1 = 4, \lambda'_2 = 3, \lambda'_3 = 1, \lambda'_4 = 1$.

Young’s lattice \mathcal{Y} is the lattice of partitions ordered by the inclusion of their Ferrers diagrams. In particular, write $\mu \prec \lambda$ if $\mu \subseteq \lambda$ and $|\lambda| = |\mu| + 1$. In other words, μ is covered by λ in \mathcal{Y} if the Ferrers diagram of λ can be obtained by adding a box to the Ferrers diagram of μ . If this box is added in the i th row and j th column of the diagram, assign a weight $c_{\mu\lambda}(q)$ to the edge between μ and λ , where

$$c_{\mu\lambda}(q) = \begin{cases} q^{|\mu| - \mu'_j}, & \text{if } j = 1, \\ q^{|\mu| - \mu'_{j-1}} (q^{\mu'_{j-1} - \mu'_j} - 1), & \text{if } j \geq 2. \end{cases} \tag{6}$$

See Figure 1 for an illustration.

The following recurrence formula for $F_\lambda(q)$ can be found in Borodin (1995), where he considers the matrices as particles of a certain mass. An elementary proof of a different flavour is outlined below.

Proposition 2 *Let $\lambda \vdash n$. The number of n by n upper-triangular nilpotent matrices over \mathbb{F}_q of Jordan type λ is*

$$F_\lambda(q) = \sum_{\mu \prec \lambda} c_{\mu\lambda}(q) F_\mu(q), \quad \text{with} \quad F_\emptyset(q) = 1.$$

Proof: Proceed by induction on n . Given $\lambda \vdash n$, first notice that any matrix in $\mathfrak{g}_{n,\lambda}(\mathbb{F}_q)$ has a leading principal submatrix of type μ where $\mu \prec \lambda$. Furthermore, let J_μ denote the Jordan matrix which is the




Fig. 1: Young’s lattice with edge weights $c_{\mu\lambda}(q)$, up to $n = 4$.

direct sum of elementary nilpotent Jordan blocks of sizes μ_1, \dots, μ_k . There are $c_{\mu\lambda}(q)$ matrices of Jordan type λ having J_μ as its leading principal submatrix, and by similarity, this result continues to hold if the matrix J_μ is replaced by any matrix Y of Jordan type μ . Summing over all $\mu \prec \lambda$ gives the desired formula. \square

The formula for $F_\lambda(q)$ in Proposition 2 can be rephrased as a sum over the set $\mathcal{P}(\lambda)$ of paths in the Young lattice \mathcal{Y} from the empty partition \emptyset to λ . Suppose

$$\varpi : \emptyset \xrightarrow{\epsilon_1} \lambda(1) \xrightarrow{\epsilon_2} \lambda(2) \xrightarrow{\epsilon_3} \dots \xrightarrow{\epsilon_n} \lambda(n) = \lambda$$

is a path in \mathcal{Y} , where the weight of the path

$$w(\varpi) = \prod_{r=1}^n c_{\lambda^{(r-1)}\lambda^{(r)}}(q) \tag{7}$$

is the product of the weights on its edges. Then Proposition 2 is equivalent to the statement

$$F_\lambda(q) = \sum_{\varpi \in \mathcal{P}(\lambda)} w(\varpi). \tag{8}$$

Example 3 *There are two partitions of 4 with 2 parts, namely $(3, 1)$ and $(2, 2)$. There are three paths from \emptyset to $(3, 1)$, giving*

$$\begin{aligned} F_{(3,1)}(q) &= (q-1) \cdot (q-1)q \cdot q^2 + (q-1) \cdot q \cdot (q-1)q^2 + (q^2-1) \cdot (q-1)q^2 \\ &= (q-1)^2 (3q^3 + q^2), \end{aligned}$$

and there are two paths from \emptyset to $(2, 2)$, giving

$$\begin{aligned} F_{(2,2)}(q) &= (q - 1) \cdot q \cdot (q - 1)q + (q^2 - 1) \cdot (q - 1)q \\ &= (q - 1)^2(2q^2 + q). \end{aligned}$$

Two observations about $F_\lambda(q)$ now follow readily from Proposition 2. For $\lambda \vdash n$, let

$$n_\lambda = \sum_{i \geq 1} (i - 1)\lambda_i. \tag{9}$$

Suppose $\varpi : \emptyset \longrightarrow \lambda^{(1)} \longrightarrow \lambda^{(2)} \longrightarrow \cdots \longrightarrow \lambda^{(n)} = \lambda$ is a path in \mathcal{Y} such that $\lambda^{(r)}$ is obtained by adding a box to $\lambda^{(r-1)}$ in row i and column j . Then $\deg c_{\lambda^{(r-1)}\lambda^{(r)}}(q) = r - i$, and therefore,

$$\deg w(\varpi) = \sum_{r=1}^n \deg c_{\lambda^{(r-1)}\lambda^{(r)}}(q) = \sum_{r=1}^n r - \sum_{i \geq 1} i\lambda_i = \binom{n}{2} - n_\lambda.$$

In particular, each polynomial $w(\varpi)$ arising from a path $\varpi \in \mathcal{P}(\lambda)$ has the same degree, so

$$\deg F_\lambda(q) = \binom{n}{2} - n_\lambda. \tag{10}$$

Moreover, each $w(\varpi)$ is monic, so the coefficient of the highest degree term in $F_\lambda(q)$ is the number of paths in \mathcal{Y} from \emptyset to λ , which is the number f^λ of standard Young tableaux of shape λ .

Second, the edge weight $c_{\lambda^{(r-1)}\lambda^{(r)}}(q)$ corresponding to the r th step in the path ϖ contributes a factor of $q - 1$ to $w(\varpi)$ if and only if the r th box added along the path is in column $j \geq 2$. Therefore, the multiplicity of $q - 1$ in each $w(\varpi)$ is $n - \lambda'_1 = n - \ell(\lambda)$, and so, the multiplicity of $q - 1$ in $F_\lambda(q)$ is $n - \ell(\lambda)$.

3 Jordan canonical forms and q -rook polynomials

A board B is a subset of an n by n grid of squares. We follow Haglund (1998), and index the squares following the convention for the entries of a matrix. A Ferrers board is a board B where if a square $s \in B$, then all squares lying north and/or east of s is also in B . Let δ_n denote the staircase-shaped board with n columns of sizes $0, 1, \dots, n - 1$. Let $\text{area}(B)$ be the number of squares in B , so $\text{area}(\delta_n) = \binom{n}{2}$ in particular.

A placement of k rooks on a board B is *non-attacking* if there is at most one rook in each row and each column of B . Let $\mathcal{C}(B, k)$ be the set of non-attacking placements of k rooks on B . For a placement $\gamma \in \mathcal{C}(B, k)$, let $\text{ne}(\gamma)$ be the number of squares in B lying directly north or directly east of a rook. Also define the *inversion* of the placement to be the number


$$\text{inv}(\gamma) = \text{area}(B) - k - \text{ne}(\gamma). \tag{11}$$

See Example 4 for an illustration. As noted in Garsia and Remmel (1986), the statistic $\text{inv}(\gamma)$ is a generalization of the number of inversions of a permutation, since permutations can be identified with non-attacking placements of rooks on a square-shaped board. In terms of the rook placement, $\text{inv}(\gamma)$ is the number of squares left blank.

Define the *weight* of a rook placement $\gamma \in \mathcal{C}(B, k)$ by

$$w(\gamma) = (q - 1)^k q^{\text{ne}(\gamma)}. \tag{12}$$

Example 4 We use \times to mark a rook and use \bullet to mark squares lying directly north or directly east of a rook. (These squares shall be referred to as the north-east squares of the placement.) The following illustration is a non-attacking placement of four rooks on the staircase-shaped board δ_7 .



This rook placement has $\text{ne}(\gamma) = 11$, $\text{inv}(\gamma) = 6$, and weight $w(\gamma) = (q - 1)^4 q^{11}$.

For $k \geq 0$, the q -rook polynomial of a Ferrers board B is defined in (Garsia and Remmel, 1986, I.4) by

$$R_k(q, B) = \sum_{c \in \mathcal{C}(B, k)} q^{\text{inv}(c)}. \tag{13}$$

The following result is due to (Haglund, 1998, Theorem 1).

Proposition 5 If B is a Ferrers board, then the number $P_{B,k}(q)$ of n by n matrices of rank k with support contained in B is



$$P_{B,k}(q) = (q - 1)^k q^{\text{area}(B) - k} R_k(q^{-1}, B).$$

□

Looking ahead, it will be convenient to consider Theorem 5 in the following equivalent form:

$$P_{B,k}(q) = \sum_{\gamma \in \mathcal{C}(B, k)} (q - 1)^k q^{\text{ne}(\gamma)} = \sum_{\gamma \in \mathcal{C}(B, k)} w(\gamma). \tag{14}$$

Example 6 There are seven non-attacking placements on δ_4 with two rooks:



This gives $P_2(q) = (q - 1)^2 (3q^3 + 3q^2 + q)$.

Recall that if $\lambda \vdash n$ is obtained from μ by adding a box in row i and column j , then the edge in \mathcal{Y} from μ to λ has weight

$$c_{\mu\lambda}(q) = \begin{cases} q^{|\mu|-\mu'_j}, & \text{if } j = 1, \\ (q-1)q^{|\mu|-\mu'_{j-1}} + \dots + (q-1)q^{|\mu|-\mu'_j}, & \text{if } j \geq 2. \end{cases} \tag{17}$$

Thus each edge weight is a sum of terms of the form $(q-1)^d q^e$ where $d = 1 - \delta_{j1}$ (δ_{j1} denotes the Kronecker delta function), and $e \in \mathbb{Z}_{\geq 0}$. This observation inspires the following definitions.

Based on the Young lattice \mathcal{Y} , we construct a graph \mathcal{Z} (see Figure 2). The vertices of \mathcal{Z} are partitions. If there is an edge from μ to λ in \mathcal{Y} of weight $q^{|\mu|-\mu'_{j-1}}(q^{\mu'_{j-1}-\mu'_j} - 1)$, then in \mathcal{Z} the edge is replaced by $\mu'_{j-1} - \mu'_j$ edges with weights

$$(q-1)q^{|\mu|-\mu'_{j-1}}, \dots, (q-1)q^{|\mu|-\mu'_j}.$$

A primitive path π

$$\emptyset \xrightarrow{\epsilon_1} \pi^{(1)} \xrightarrow{\epsilon_2} \pi^{(2)} \xrightarrow{\epsilon_3} \dots \xrightarrow{\epsilon_n} \pi^{(n)}$$

is a path in the graph \mathcal{Z} . The weight $w(\pi)$ of a primitive path is the product of its edge weights.

Let $\mathcal{PP}(\lambda)$ denote the set of primitive paths from \emptyset to λ . Then each path in Young's lattice corresponds to a set of primitive paths, and

$$F_\lambda(q) = \sum_{\pi \in \mathcal{PP}(\lambda)} w(\pi). \tag{18}$$

Remark 7 Let $\mu \vdash n-1$ where $\ell(\mu) = \ell$. Let $\pi' : \emptyset \longrightarrow \lambda^{(1)} \longrightarrow \lambda^{(2)} \longrightarrow \dots \longrightarrow \lambda^{(n-1)} = \mu$ be a primitive path. If λ is obtained by adding a box to the first column of μ , then there is a unique way to extend the primitive path π' by one edge, and by Equation 17, that edge has weight

$$q^{|\mu|-\ell}.$$

Furthermore, consider all possible λ which can be obtained by adding a box to μ in a column $j \geq 2$. Then by Equation 17, there are

$$\sum_{j=2}^{\ell(\mu')+1} \mu'_{j-1} - \mu'_j = \ell$$

ways to extend the primitive path π' by one edge.

In summary, the out-degree of μ in \mathcal{Z} is $\ell + 1$. Moreover, the weights

$$q^{|\mu|-\ell}, (q-1)q^{|\mu|-\ell}, (q-1)q^{|\mu|-\ell+1}, \dots, (q-1)q^{|\mu|-1},$$

for the edges have unique degrees $|\mu| - \ell \leq d \leq |\mu|$. This observation is crucial for the proof of the next lemma.

Lemma 8 Let $n \geq 1$ and $0 \leq k \leq n - 1$. Suppose $\gamma \in \mathcal{C}(\delta_n, k)$ is a rook placement with columns $\gamma^{(1)}, \dots, \gamma^{(n)}$. Then the sequence of weights $w(\gamma^{(1)}), \dots, w(\gamma^{(n)})$ determines a unique primitive path $\pi : \emptyset \xrightarrow{\epsilon_1} \pi^{(1)} \xrightarrow{\epsilon_2} \pi^{(2)} \xrightarrow{\epsilon_3} \dots \xrightarrow{\epsilon_n} \pi^{(n)}$ such that $\epsilon_r = w(\gamma^{(r)})$. Moreover, $\ell(\pi^{(n)}) = n - k$.




Fig. 2: The graph \mathcal{Z} , up to $n = 4$.

Proof: Proceed by induction on $n + k$. Suppose $\gamma \in \mathcal{C}(\delta_n, k)$ and let γ' be the placement consisting of the first $n - 1$ columns of γ . By induction, the sequence $\epsilon_1 = w(\gamma^{(1)}), \dots, \epsilon_{n-1} = w(\gamma^{(n-1)})$ determines a unique primitive path $\pi' : \emptyset \xrightarrow{\epsilon_1} \pi^{(1)} \xrightarrow{\epsilon_2} \dots \xrightarrow{\epsilon_{n-1}} \pi^{(n-1)}$. There are two cases to consider; γ' has either k or $k - 1$ rooks.

If γ' has k rooks, then the column $\gamma^{(n)}$ then has k north-east squares only and weight $q^k = q^{|\mu|-\ell}$. By the previous remark, there is a unique way to extend the primitive path corresponding to γ' by adding a box to the first column of $\pi^{(n-1)}$, and that edge has weight $q^{|\mu|-\ell}$.

In the second case, if γ' has $k - 1$ rooks, then there are $n - 1 - k$ available boxes in column $\gamma^{(n)}$ to place a rook. The placement of the rook uniquely determines the degree of the weight $w(\gamma^{(n)})$ of the column, which ranges from $k + 1, \dots, n - 1$. From the remark, there are precisely $\ell = n - 1 - k$ edges emanating from μ in \mathcal{Z} with weights having degrees $|\mu| - \ell + 1, \dots, |\mu|$. \square

Let $\mathcal{PP}(n, n - k) = \{\pi \in \mathcal{PP}(\lambda) \mid \lambda \vdash n \text{ and } \ell(\lambda) = n - k\}$ be the set of primitive paths in \mathcal{Z} from \emptyset to a partition with $n - k$ parts. Define a map $\Theta : \mathcal{C}(\delta_n, k) \rightarrow \mathcal{PP}(n, n - k)$ as follows. Suppose a rook placement γ has columns $\gamma^{(1)}, \dots, \gamma^{(n)}$. Let $\Theta(\gamma)$ be the primitive path

$$\emptyset \xrightarrow{\epsilon_1} \pi^{(1)} \xrightarrow{\epsilon_2} \pi^{(2)} \xrightarrow{\epsilon_3} \dots \xrightarrow{\epsilon_n} \pi^{(n)} \quad \text{with} \quad \epsilon_r = w(\gamma^{(r)}).$$

Theorem 9 The map $\Theta : \mathcal{C}(\delta_n, k) \rightarrow \mathcal{PP}(n, n - k)$ is a weight-preserving bijection. \square


It follows from Theorem 9 that we may associate a partition type to each rook placement on δ_n . The *partition type* of a rook placement γ is the partition at the endpoint of the primitive path $\Theta(\gamma)$. Let $\mathcal{C}(\lambda)$ be the set of rook placements of partition type λ . As a Corollary, we obtain a formula for $F_\lambda(q)$ as a sum over rook placements, and the equation can be viewed as a generalization of Haglund’s result in Equation 4.

Corollary 10 *Let $\lambda \vdash n$ be a partition with $\ell(\lambda) = n - k$ parts. Then*


$$F_\lambda(q) = \sum_{\gamma \in \mathcal{C}(\lambda)} (q - 1)^k q^{\text{ne}(\gamma)}.$$

Proof: The result follows from Equation 18 and the bijection Θ in Theorem 9. □

Example 11 *There are four primitive paths from \emptyset to $\lambda = (3, 1)$.*



Respectively, they correspond to the following rook placements, each having partition type $(3, 1)$.



Therefore,

$$F_{(3,1)}(q) = (q - 1)^2 (3q^3 + q^2).$$

4 A refinement to compositions

A *set partition* is a set $\sigma = \{B_1, \dots, B_k\}$ of nonempty disjoint subsets of \mathcal{N}_n such that $\bigcup_{i=1}^k B_i = \mathcal{N}_n$. The B_i s are the *blocks* of σ . Let $\Pi(n, k)$ be the set of set partitions of \mathcal{N}_n with k blocks. We adopt the convention of listing the blocks in order so that $\min B_i < \min B_j$ if $i < j$. This allows us to represent a set partition with a diagram similar to that of a Young tableau; the i th row of the diagram consists of the elements in block B_i listed in increasing order. A *composition* α of a nonnegative integer n is a sequence of positive integers $(\alpha_1, \dots, \alpha_k)$ such that $|\alpha| = \sum_{i=1}^k \alpha_i = n$. If α has k positive parts, write $\ell(\alpha) = k$. A set partition $\sigma = \{B_1, \dots, B_k\}$ has *composition type* α if $\alpha = (|B_1|, \dots, |B_k|)$.

The number of set partitions of \mathcal{N}_n with k blocks is the Stirling number $S_{n,n-k}(1)$ (see Equation 3). In addition, $S_{n,n-k}(1)$ is also the number of placements of k rooks on the staircase board δ_n . This follows from the following well-known bijection (see Stanley (1999)); given a placement $\gamma \in \mathcal{C}(\delta_n, k)$, construct

a set partition of \mathcal{N}_n where the integers i and j are in the same block if and only if there is a rook in square $(i, j) \in \gamma$.

We shall give another bijection $\Psi : \mathcal{C}(\delta_n, k) \rightarrow \Pi(n, n - k)$ that arises from the primitive paths in the graph \mathcal{Z} . Let $\gamma \in \mathcal{C}(\delta_n, k)$ be a rook placement. Construct a diagram for a set partition using the following procedure (also see Example 14).

- Let $\lambda^{(1)}$ be the diagram with a single box labelled 1 placed in the first row and the first column.
- For $k \geq 2$, if the weight of the k th column in γ has degree d , then place the box labelled k in the $(k - d)$ th row of $\lambda^{(k-1)}$, and rearrange the rows of the diagram into a partition shape $\lambda^{(k)}$, so that the rows of the same length have first column entries in increasing order.

Note that if $\pi = \Theta(\gamma)$ is the primitive path in \mathcal{Z} which corresponds to the placement γ , then $\pi^{(k)} = \lambda^{(k)}$. Let $\lambda^{(n)}$ be the partition shape of the diagram after the n th box has been placed. Let $\text{order}(\lambda^{(n)})$ be the diagram of the set partition resulting from ordering the rows of $\lambda^{(n)}$ so that the first column entries are increasing. Define $\Psi(\gamma) = \text{order}(\lambda^{(n)})$.

Proposition 12 $\Psi : \mathcal{C}(\delta_n, k) \rightarrow \Pi(n, n - k)$ is a bijection. □

The *composition type* of a rook placement $\gamma \in \mathcal{C}(\delta_n, k)$ is the composition type of $\Psi(\gamma)$. Let $\mathcal{C}(\alpha)$ be the set of rook placements with composition type α . For a composition α of n with $\ell(\alpha) = n - k$ parts, define

$$F_\alpha(q) = \sum_{c \in \mathcal{C}(\alpha)} (q - 1)^k q^{\text{ne}(c)}. \tag{19}$$

Let $\text{rearr}(\alpha)$ be the partition resulting from the rearrangement of the parts of the composition α so that they are nondecreasing.


Corollary 13 Let $\lambda \vdash n$. Then

$$F_\lambda(q) = \sum_{\lambda = \text{rearr}(\alpha)} F_\alpha(q). \tag{20}$$


□

We extend the definition of n_λ to compositions and let $n_\alpha = \sum_{i \geq 1} (i - 1)\alpha_i$. Then $\deg F_\alpha(q) = \binom{n}{2} - n_\alpha$, the multiplicity of the factor $q - 1$ in $F_\alpha(q)$ is $n = \ell(\alpha)$, and the coefficient of the highest degree term is the number of set partitions whose diagrams are increasing along rows and columns.

Example 14 Consider the rook placement $\gamma \in \mathcal{C}(\delta_8, 5)$:



The set partition diagrams $\lambda^{(1)}, \dots, \lambda^{(n)}$ correspond to the following primitive path in \mathcal{Z} .



The edge weights of this primitive path are the same as the column weights in the rook placement. Ordering the rows of the diagram of the endpoint of the primitive path so that the entries in the first column are increasing gives the set partition

$$\Psi(\gamma) = \text{order} \cdot \begin{array}{|c|c|c|} \hline 1 & 5 & 6 \\ \hline 3 & 7 & 8 \\ \hline 2 & 4 & \\ \hline \end{array} = \begin{array}{|c|c|c|} \hline 1 & 5 & 6 \\ \hline 2 & 4 & \\ \hline 3 & 7 & 8 \\ \hline \end{array}$$

Therefore, the rook placement γ has partition type $\lambda = (3, 3, 2)$ and composition type $\alpha = (3, 2, 3)$.

Remark 15 We can define a lattice \mathcal{X} of compositions by requiring that each path from \emptyset to α encodes a rook configuration of composition type α . The paper of Björner and Stanley (2005) considers two different lattices of compositions, but \mathcal{X} is different from the two presented in their paper. It may be interesting to investigate the combinatorial properties of \mathcal{X} , particularly as paths in \mathcal{X} are equivalent to set partitions, and they are known to play a crucial role in the supercharacter theory of unipotent upper-triangular matrices (see Thiem (2010) for example).

5 Closing remarks

5.1 Inverse Kostka-polynomials

Let $P_\lambda(x; t)$ denote the Hall-Littlewood function indexed by the partition λ , and let $m_\mu(x)$ denote the monomial symmetric function indexed by μ . See (Macdonald, 1995, Ch. III) for definitions. For $\lambda, \mu \vdash n$, the transition coefficients $L_{\lambda, \mu}(t)$ are defined by

$$P_\lambda(x; t) = \sum_{\mu} L_{\lambda, \mu}(t) m_\mu(x). \tag{20}$$

The recurrence formula for $F_\lambda(q)$ in Proposition 2 is essentially the same as the one for $L_{\lambda, 1^n}(t)$ (Macdonald, 1995, Equation 5.9'), so that $L_{\lambda, 1^n}(t) = t^{\binom{n}{2} - n\lambda} F_\lambda(t^{-1})$. It would be interesting to see if other entries in the transition matrix can be obtained as sums over rook placements on boards of another shape.

5.2 Matrices satisfying $X^2 = 0$

Kirillov and Melnikov (1995) considered the number $A_n(q)$ of n by n upper-triangular matrices over \mathbb{F}_q satisfying $X^2 = 0$. In their first characterization of these polynomials, they considered the number $A_n^r(q)$ of matrices of a given rank r , so that $A_n(q) = \sum_{r \geq 0} A_n^r(q)$, and observed that $A_n^r(q)$ satisfies the recurrence

$$A_n^r(q) = q^r A_{n-1}^r(q) + (q^{n-r} - q^r) A_n^r(q), \quad A_n^0(q) = 1.$$

We may think of $A_n(q)$ as the sum of $F_\lambda(q)$ over $\lambda \vdash n$ with at most two columns, so Proposition 2 is, in a sense, a generalization of this recurrence.

It was also conjectured by Kirillov and Melnikov that the same sequence of polynomials arise in a number of different ways. Ekhad and Zeilberger (1996) proved that one of the conjectured alternate definitions of $A_n(q)$, namely

$$C_n(q) = \sum_s c_{n+1,s} q^{\frac{n^2}{4} + \frac{1-s^2}{12}},$$

is a sum over all $s \in [-n-1, n+1]$ which satisfy $s \equiv n+1 \pmod{2}$ and $s \equiv (-1)^n \pmod{3}$, and $c_{n+1,s}$ are entries in the signed Catalan triangle, is indeed the same as $A_n(q)$. It would be interesting to see what other combinatorics may arise from considering the sum of $F_\lambda(q)$ over $\lambda \vdash n$ with at most k columns for a fixed k .

Acknowledgements

I would like to thank Jim Haglund and Alexandre Kirillov for their invaluable guidance.

References

- A. Björner and R. P. Stanley. An analogue of young's lattice for compositions. 2005.
- A. M. Borodin. Limit jordan normal form of large triangular matrices over a finite field. *Funct. Anal. Appl.*, 29(4):279–281, 1995.
- S. Ekhad and D. Zeilberger. The number of solutions of $x^2 = 0$ in triangular matrices over $gf(q)$. *Elec. J. Comb.*, 3, 1996.
- A. Garsia and J. B. Remmel. q -counting rook configurations and a formula of frobenius. *J. Combin. Theory Ser. A*, 41:246–275, 1986.
- J. Haglund. q -rook polynomials and matrices over finite fields. *Adv. in Appl. Math.*, 20:450–487, 1998.
- A. A. Kirillov. Variations on the triangular theme. *Amer. Math. Soc. Transl.*, 169(2):43–73, 1995.
- A. A. Kirillov and A. Melnikov. On a remarkable sequence of polynomials. *Sémin. Congr. 2, Soc. Math. France*, pages 35–42, 1995.
- I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford University Press, 1995.
- L. Solomon. The bruhat decomposition, tits system and iwahori ring for the monoid of matrices over a finite field. *Geom. Dedicata.*, 36:15–49, 1990.
- R. P. Stanley. *Enumerative Combinatorics*. Cambridge University Press, 1999.
- N. Thiem. Branching rules in the ring of superclass functions of unipotent upper-triangular matrices. *J. Algebraic. Combin.*, 2:267–298, 2010.
- M. Yip. q -rook placements and jordan forms of upper-triangular nilpotent matrices. 2013.