

On the diameter of random planar graphs


Guillaume Chapuy, CNRS & LIAFA, Paris

joint work with

Éric Fusy, Paris,
Omer Giménez, ex-Barcelona,
Marc Noy, Barcelona.


Planar graphs and maps

- **Planar graph** = (connected) graph on $V = \{1, 2, \dots, n\}$ that can be drawn in the plane without edge crossing.




Planar graphs and maps


- **Planar graph** = (connected) graph on $V = \{1, 2, \dots, n\}$ that can be drawn in the plane without edge crossing.




- **Planar map** = planar graph + **planar drawing of this graph** (up to continuous deformation)



=




≠




same graph
different maps

Planar graphs and maps


- **Planar graph** = (connected) graph on $V = \{1, 2, \dots, n\}$ that can be drawn in the plane without edge crossing.




- **Planar map** = planar graph + **planar drawing of this graph** (up to continuous deformation)



=



≠



same graph
different maps

- Note: the number of embeddings depends on the graph...


Uniform random planar map \neq Uniform random planar graph!

Some known results for maps (stated approximately)

- **Thm** [Chassaing-Schaeffer '04], [Marckert, Miermont '06], [Ambjörn-Budd '13]

In a uniform random map M_n of size n , distances are of order $n^{1/4}$.

For example one has $\frac{\text{Diam}(M_n)}{n^{1/4}} \rightarrow$ some real random variable




Some known results for maps (stated approximately)

- **Thm** [Chassaing-Schaeffer '04], [Marckert, Miermont '06], [Ambjörn-Budd '13]

In a uniform random map M_n of size n , distances are of order $n^{1/4}$.

For example one has $\frac{\text{Diam}(M_n)}{n^{1/4}} \rightarrow$ some real random variable




Some known results for maps (stated approximately)

- **Thm** [Chassaing-Schaeffer '04], [Marckert, Miermont '06], [Ambjörn-Budd '13]

In a uniform random map M_n of size n , distances are of order $n^{1/4}$.

For example one has $\frac{\text{Diam}(M_n)}{n^{1/4}} \rightarrow$ some real random variable



A lot of (very strong) things are known – very active field of research since 2004 [Bouttier, Di Francesco, Guitter, Le Gall, Miermont, Paulin, Addario-Berry, Albenque...]

Our main result: diameter of random planar GRAPHS

- **Thm** [C, Fusy, Giménez, Noy 2010+]

Let G_n be the uniform random planar graph with n vertices.

Then $\text{Diam}(G_n) = n^{1/4+o(1)}$ w.h.p.

More precisely $\mathbb{P}\left(\text{Diam}(G_n) \notin [n^{1/4-\epsilon}, n^{1/4+\epsilon}]\right) = O(e^{-n^{\Theta(\epsilon)}})$.

Our main result: diameter of random planar GRAPHS

- **Thm** [C, Fusy, Giménez, Noy 2010+]

Let G_n be the uniform random planar graph with n vertices.

Then $\text{Diam}(G_n) = n^{1/4+o(1)}$ w.h.p.

More precisely $\mathbb{P}\left(\text{Diam}(G_n) \notin [n^{1/4-\epsilon}, n^{1/4+\epsilon}]\right) = O(e^{-n^{\Theta(\epsilon)}})$.

- This is some kind of large deviation result. We also conjecture convergence in law:

$$\frac{\text{Diam}(G_n)}{n^{1/4}} \rightarrow \text{some real random variable}$$

- Note: for random trees,


$$\frac{\text{Diam}(T_n)}{n^{1/2}} \rightarrow \text{some real random variable}$$

$$\mathbb{P}\left(\text{Diam}(T_n) \notin [n^{1/2-\epsilon}, n^{1/2+\epsilon}]\right) = O(e^{-n^{\Theta(\epsilon)}})$$


[Flajolet et al '93]

(0) Connectivity in graphs


General




Connected
(1-connected)



2-Connected




3-Connected



A graph is k -connected if one needs to remove at least k vertices to disconnect it.


(0) Connectivity in graphs

General




A graph is k -connected if one needs to remove at least k vertices to disconnect it.



Connected (1-connected)



2-Connected



3-Connected




- [Tutte'66]:
- a connected graph decomposes into 2-connected components
 - a 2-connected graph decomposes into 3-connected components


[Whitney]: A 3-connected planar graph has a UNIQUE embedding

(0) Connectivity in graphs


General




Connected
(1-connected)




2-Connected



3-Connected




A graph is k -connected if one needs to remove at least k vertices to disconnect it.



- [Tutte'66]: - a connected graph decomposes into 2-connected components
- a 2-connected graph decomposes into 3-connected components

[Whitney]: A 3-connected planar graph has a UNIQUE embedding


[Tutte 60s], [Bender, Gao, Wormald'02], [Giménez, Noy'05] followed this path carrying counting results along the scheme → exact counting of planar graphs!
Here we follow the same path and carry deviations statements for the diameter.



(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)



- To simplify the exposition we consider a **quadrangular planar map** (faces have degree 4)

1. Label vertices by their **graph-distance** to some root vertex



(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- To simplify the exposition we consider a **quadrangular planar map** (faces have degree 4)
 1. Label vertices by their **graph-distance** to some root vertex
 2. Observe that there are only **two types** of faces (since bipartite)





(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- To simplify the exposition we consider a **quadrangular planar map** (faces have degree 4)

1. Label vertices by their **graph-distance** to some root vertex

2. Observe that there are only **two types** of faces (since bipartite)



3. Apply **Schaeffer rules**:



(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- To simplify the exposition we consider a **quadrangular planar map** (faces have degree 4)



1. Label vertices by their **graph-distance** to some root vertex
2. Observe that there are only **two types** of faces (since bipartite)
3. Apply **Schaeffer rules**:



(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- To simplify the exposition we consider a **quadrangular planar map** (faces have degree 4)



1. Label vertices by their **graph-distance** to some root vertex
2. Observe that there are only **two types** of faces (since bipartite)
3. Apply **Schaeffer rules**:



(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- To simplify the exposition we consider a **quadrangular planar map** (faces have degree 4)



1. Label vertices by their **graph-distance** to some root vertex
2. Observe that there are only **two types** of faces (since bipartite)
3. Apply **Schaeffer rules**:



(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- To simplify the exposition we consider a **quadrangular planar map** (faces have degree 4)



1. Label vertices by their **graph-distance** to some root vertex
2. Observe that there are only **two types** of faces (since bipartite)
3. Apply **Schaeffer rules**:



(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- To simplify the exposition we consider a **quadrangular planar map** (faces have degree 4)


1. Label vertices by their **graph-distance** to some root vertex
2. Observe that there are only **two types** of faces (since bipartite)
3. Apply **Schaeffer rules**:




(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- To simplify the exposition we consider a **quadrangular planar map** (faces have degree 4)

1. Label vertices by their **graph-distance** to some root vertex
2. Observe that there are only **two types** of faces (since bipartite)
3. Apply **Schaeffer rules**:




Fact: the blue map is a **tree**.




(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- To simplify the exposition we consider a **quadrangular planar map** (faces have degree 4)

1. Label vertices by their **graph-distance** to some root vertex
2. Observe that there are only **two types** of faces (since bipartite)
3. Apply **Schaeffer rules**:




Fact: the blue map is a **tree**.




(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- To simplify the exposition we consider a **quadrangular planar map** (faces have degree 4)

1. Label vertices by their **graph-distance** to some root vertex
2. Observe that there are only **two types** of faces (since bipartite)
3. Apply **Schaeffer rules**:




Fact: the blue map is a **tree**.




(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- To simplify the exposition we consider a **quadrangular planar map** (faces have degree 4)

1. Label vertices by their **graph-distance** to some root vertex
2. Observe that there are only **two types** of faces (since bipartite)
3. Apply **Schaeffer rules**:




Fact: the blue map is a **tree**.




(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- To simplify the exposition we consider a **quadrangular planar map** (faces have degree 4)

1. Label vertices by their **graph-distance** to some root vertex
2. Observe that there are only **two types** of faces (since bipartite)
3. Apply **Schaeffer rules**:




Fact: the blue map is a **tree**.



If one remembers the labels the construction is **bijective**!


(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- A **well-labelled tree** is a plane tree together with a mapping $l : V \rightarrow \mathbb{Z}_{>0}$ such that
 - if $v \sim v'$ then $|l(v) - l(v')| \leq 1$
 - $\min_v l(v) = 1$



(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- A **well-labelled tree** is a plane tree together with a mapping $l : V \rightarrow \mathbb{Z}_{>0}$ such that
 - if $v \sim v'$ then $|l(v) - l(v')| \leq 1$
 - $\min_v l(v) = 1$
- **Thm** [Cori-Vauquelin'81;Schaeffer'99]
There is a bijection between **quadrangular planar maps** with a pointed vertex and $n + 1$ vertices and **well-labelled trees** with n vertices. The labels in the tree correspond to **distances to the root in the map**.



(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)


- A **well-labelled tree** is a plane tree together with a mapping $l : V \rightarrow \mathbb{Z}_{>0}$ such that
 - if $v \sim v'$ then $|l(v) - l(v')| \leq 1$
 - $\min_v l(v) = 1$

- **Thm** [Cori-Vauquelin'81;Schaeffer'99]

There is a bijection between **quadrangular planar maps** with a pointed vertex and $n + 1$ vertices and **well-labelled trees** with n vertices. The labels in the tree correspond to **distances to the root in the map**.

Corollary: $\text{Diam}(M_n) = n^{1/4+o(1)}$

indeed: - the height of a random tree is $\approx n^{1/2+o(1)}$ w.h.p



(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)


- A **well-labelled tree** is a plane tree together with a mapping $l : V \rightarrow \mathbb{Z}_{>0}$ such that

- if $v \sim v'$ then $|l(v) - l(v')| \leq 1$

- $\min_v l(v) = 1$

- **Thm** [Cori-Vauquelin'81; Schaeffer'99]

There is a bijection between **quadrangular planar maps** with a pointed vertex and $n + 1$ vertices and **well-labelled trees** with n vertices. The labels in the tree correspond to **distances to the root in the map**.



Corollary: $\text{Diam}(M_n) = n^{1/4+o(1)}$

indeed: - the height of a random tree is $\approx n^{1/2+o(1)}$ w.h.p


- the labelling function behaves as a random walk along branches of the tree so $l(v) \approx \sqrt{n^{1/2+o(1)}} = n^{1/4+o(1)}$

(1) Maps: the Cori-Vauquelin-Schaeffer bijection (1981-1999-2008+)

- A **well-labelled tree** is a plane tree together with a mapping $l : V \rightarrow \mathbb{Z}_{>0}$ such that
 - if $v \sim v'$ then $|l(v) - l(v')| \leq 1$
 - $\min_v l(v) = 1$

- **Thm** [Cori-Vauquelin'81; Schaeffer'99]


There is a bijection between **quadrangular planar maps** with a pointed vertex and $n + 1$ vertices and **well-labelled trees** with n vertices. The labels in the tree correspond to **distances to the root in the map**.




Corollary: $\text{Diam}(M_n) = n^{1/4+o(1)}$

indeed: - the height of a random tree is $\approx n^{1/2+o(1)}$ w.h.p
- the labelling function behaves as a random walk along branches of the tree so $l(v) \approx \sqrt{n^{1/2+o(1)}} = n^{1/4+o(1)}$

[Chassaing-Schaeffer'04]




(2) Decomposition into 2-connected components



A connected map is “a tree of 2-connected maps”


(2) Decomposition into 2-connected components



A connected map is “a tree of 2-connected maps”

One can write this in terms of generating functions.

(2) Decomposition into 2-connected components



A connected map is “a tree of 2-connected maps”


One can write this in terms of generating functions.

The generating function of connected maps is explicitly known (e.g. using bijections)

→ deduce the g.f. of **2-connected maps** from the one of connected maps.

[Tutte 60's].

(2) Decomposition into 2-connected components



A connected map is “a tree of 2-connected maps”

One can write this in terms of generating functions.

The generating function of connected maps is explicitly known (e.g. using bijections)

→ deduce the g.f. of 2-connected maps from the one of connected maps.

[Tutte 60's].


(2) Decomposition into 2-connected components

Thm

The largest 2-connected component has size $\frac{n}{3} + n^{2/3}A$ where A converges to an explicit law.

The second-largest component has size $O(n^{2/3})$.

[Gao, Wormald'99] [Banderier, Flajolet, Schaeffer, Soria '01]



(2) Decomposition into 2-connected components


Thm

The largest 2-connected component has size $\frac{n}{3} + n^{2/3}A$ where A converges to an explicit law.

The second-largest component has size $O(n^{2/3})$.

[Gao, Wormald'99] [Banderier, Flajolet, Schaeffer, Soria '01]

tools to prove this: (very) fine singularity analysis of generating functions. In particular exact expressions for counting functions are mandatory!



(2) Decomposition into 2-connected components

Thm

The largest 2-connected component has size $\frac{n}{3} + n^{2/3}A$ where A converges to an explicit law.

The second-largest component has size $O(n^{2/3})$.


[Gao, Wormald'99] [Banderier, Flajolet, Schaeffer, Soria '01]

tools to prove this: (very) fine singularity analysis of generating functions. In particular exact expressions for counting functions are mandatory!

“Corollary”:

$$\text{Diam}(B_{n/3}) \sim \text{Diam}(M_n) = n^{1/4+o(1)}$$

random 2-conn. map of size $n/3$



(2) Decomposition into 2-connected components

Thm

The largest 2-connected component has size $\frac{n}{3} + n^{2/3}A$ where A converges to an explicit law.

The second-largest component has size $O(n^{2/3})$.

[Gao, Wormald'99] [Banderier, Flajolet, Schaeffer, Soria '01]


tools to prove this: (very) fine singularity analysis of generating functions. In particular exact expressions for counting functions are mandatory!

“Corollary”:

$$\text{Diam}(B_{n/3}) \sim \text{Diam}(M_n) = n^{1/4+o(1)}$$

random 2-conn. map of size $n/3$

indeed: $\text{Diam}(X_n) \leq \text{Diam}(M_n) \leq \text{Diam}(X_n) + 2 \max_i \text{Diam}(H_i)$



(2) Decomposition into 2-connected components

Thm

The largest 2-connected component has size $\frac{n}{3} + n^{2/3}A$ where A converges to an explicit law.

The second-largest component has size $O(n^{2/3})$.

[Gao, Wormald'99] [Banderier, Flajolet, Schaeffer, Soria '01]


tools to prove this: (very) fine singularity analysis of generating functions. In particular exact expressions for counting functions are mandatory!

“Corollary”:

$$\text{Diam}(B_{n/3}) \sim \text{Diam}(M_n) = n^{1/4+o(1)}$$

random 2-conn. map of size $n/3$

indeed: $\text{Diam}(X_n) \leq \text{Diam}(M_n) \leq \text{Diam}(X_n) + 2 \max_i \text{Diam}(H_i) \leq (n^{2/3})^{1/4+\epsilon}$ w.h.p.



(2) Decomposition into 2-connected components

Thm

The largest 2-connected component has size $\frac{n}{3} + n^{2/3}A$ where A converges to an explicit law.

The second-largest component has size $O(n^{2/3})$.

[Gao, Wormald'99] [Banderier, Flajolet, Schaeffer, Soria '01]

tools to prove this: (very) fine singularity analysis of generating functions. In particular exact expressions for counting functions are mandatory!


“Corollary”:

$$\text{Diam}(B_{n/3}) \sim \text{Diam}(M_n) = n^{1/4+o(1)}$$


random 2-conn. map of size $n/3$


indeed: $\text{Diam}(X_n) \leq \text{Diam}(M_n) \leq \text{Diam}(X_n) + 2 \max_i \text{Diam}(H_i)$

and X_n is essentially a random 2-conn. map of size $n/3$.




$$\leq (n^{2/3})^{1/4+\epsilon} \text{ w.h.p.}$$





(3) Decomposition into 3-connected components



Again one can write everything **in terms of generating functions**.

→ deduce the g.f. of **3-conn. maps** from the one of 2-connected maps. [Tutte 60's].

→ deduce the g.f. of **2-conn. graphs** from the one of 3-connected graphs [Bender, Gao, Wormald'02].


\textcircled{T} = 3-connected component

\textcircled{R} = series composition



\textcircled{M} = parallel composition

(3) Decomposition into 3-connected components

Prop A random 2-connected planar graph with n edges has diameter $n^{1/4+o(1)}$ with high probability.




a 2-conn. graph B_n



RMT tree

(3) Decomposition into 3-connected components

Prop A random 2-connected planar graph with n edges has diameter $n^{1/4+o(1)}$ with high probability.




Same idea:

- there exists a T -component Y_n of linear size w.h.p.

(3) Decomposition into 3-connected components

Prop A random 2-connected planar graph with n edges has diameter $n^{1/4+o(1)}$ with high probability.



Same idea:

- there exists a T -component Y_n of linear size w.h.p.
- the diameter of the RMT-tree is $n^{o(1)}$ w.h.p.
- The extra-length due the edge substitution is also $n^{o(1)}$

Conclusion (I)

- **Thm** [C, Fusy, Giménez, Noy 2010+]

Let G_n be the uniform random planar graph with n vertices.

Then $\text{Diam}(G_n) = n^{1/4+o(1)}$ w.h.p.

More precisely $\mathbb{P}\left(\text{Diam}(G_n) \notin [n^{1/4-\epsilon}, n^{1/4+\epsilon}]\right) = O(e^{-n^{\Theta(\epsilon)}})$.

- The proof relies both on exact generating functions and magical bijections: we couldn't do anything without this (or maybe something much weaker like $O(\sqrt{n})$?)
- The general picture is quite clear but the analysis is a bit tedious... (need to work with bivariate generating functions and prove estimates with enough uniformity)
- No way to obtain the convergence of $\frac{\text{Diam}(G_n)}{n^{1/4}}$ - even for planar maps this is very difficult!
- Same result for the uniform random graph with n vertices and $\lfloor \mu n \rfloor$ edges for $1 < \mu < 3$.

Conclusion (II)

- We generalized the Giménez-Noy enumeration result to **graphs embeddable on a surface of genus $g \geq 0$**

Thm [C, Fusy, Giménez, Mohar, Noy 2011] [Bender-Gao 2011]

$$\#\{n\text{-vertex genus } g \text{ graphs}\} \sim c_g \cdot n! \cdot \gamma^n \cdot n^{\frac{5}{2}g - 7/2} \quad \gamma \approx 27. \dots$$

Same kind of proof but Whitney's theorem (uniqueness of embedding) now requires that **there is no short non-contractible cycle**.


(but we could prove that)

The result on the diameter **should be the same** but this is not (and won't be) written.

The fact that non-contractible cycles are small imply the following:

Thm [C, Fusy, Giménez, Mohar, Noy 2011]

Fix $g \geq 1$. The random graph of genus g and size n has **chromatic number in $\{4, 5\}$** and **list chromatic number 5** w.h.p.



Thank you!