

1 Categorifying Non-Idempotent Intersection Types

2 Giulio Guerrieri 

3 University of Bath, Department of Computer Science, Bath, United Kingdom.

4 g.guerrieri@bath.ac.uk

5 Federico Olimpieri

6 Institut de Mathématiques de Marseille (I2M), Aix-Marseille Université, Marseille, France.

7 federico.olimpieri@univ-amu.fr

8 Abstract

9 Non-idempotent intersection types can be seen as a syntactic presentation of a well-known denota-
10 tional semantics for the lambda-calculus, the category of sets and relations. Building on previous
11 work, we present a categorification of this line of thought in the framework of the bang calculus,
12 an untyped version of Levy’s call-by-push-value. We define a bicategorical model for the bang
13 calculus, whose syntactic counterpart is a suitable category of types. In the framework of distributors,
14 we introduce intersection type distributors, a bicategorical proof relevant refinement of relational
15 semantics. Finally, we prove that intersection type distributors characterize normalization at depth 0.

16 **2012 ACM Subject Classification** Theory of computation → Lambda calculus; Theory of computa-
17 tion → Linear logic; Theory of computation → Categorical semantics

18 **Keywords and phrases** linear logic, bang calculus, non-idempotent intersection types, distributors,
19 relational semantics, combinatorial species, symmetric sequences, bicategory, categorification

20 **Digital Object Identifier** 10.4230/LIPIcs.CSL.2021.38

21 **Funding** This work is partially supported by EPSRC Project EP/R029121/1 *Typed lambda-calculi*
22 *with sharing and unsharing*

23 **Acknowledgements** The authors thank Lionel Vaux Auclair for insightful discussions and comments.

24 1 Introduction

25 Since Girard’s introduction of *linear logic* [32], the notion of linearity has played a central
26 role in the Logic-in-Computer-Science community. A program is linear when it uses its
27 inputs only *once* during computation (inputs cannot be copied or deleted); while a non-linear
28 program may call its inputs at will. Via the exponential modalities ! and ?, linear logic gives
29 a logical status to the operations of erasing and copying data.

30 Another way to study linearity is provided by some type systems. *Intersection types* were
31 introduced by Coppo and Dezani [14, 15] as an extension of simple types by means of the
32 (associative, commutative and idempotent) intersection connective $a \cap b$: a term of type
33 $a \cap b$ can be seen as a program of both type a and type b . This kind of type systems have
34 proven to be very useful to characterize various notion of normalization in the λ -calculus
35 [37]. If we impose *non-idempotency* to the intersection [31, 16] (i.e. $a \cap a \neq a$), we get a
36 “resource-sensitive” intersection type system, in the sense that the arrow type encodes the
37 *exact* number of times that a term needs its input during computation: intuitively, a term
38 typed $a \cap a \cap b$ can be used twice as a program of type a and once as a program of type
39 b . Non-idempotent intersection types allow *combinatorial* characterization of normalization
40 properties and of the execution time of programs [9, 16, 4] and proof-nets [19, 20]. Also,
41 De Carvalho’s non-idempotent intersection type system \mathcal{R} is a syntactic presentation of the
42 categorical semantics of λ -calculus given in the category of sets and relations [16, 17]. There
43 is a strong connection between linear logic and non-idempotent intersection types [18].



© Giulio Guerrieri, Federico Olimpieri;
licensed under Creative Commons License CC-BY

29th EACSL Annual Conference on Computer Science Logic (CSL 2021).

Editors: Christel Baier and Jean Goubault-Larrecq; Article No. 38; pp. 38:1–38:24

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

44 Inspired by [34, 41, 47, 43], we propose here a *categorification* of this kind of semantics.
 45 Roughly, categorification consists in replacing set-theoretic notions with category-theoretic
 46 ones. In general, this process gives both more fine-grained structures and general points of
 47 view. Mellies and Zeilberger [42] followed this approach to present a categorical definition of
 48 what a type system is: a type system is a *functor* between a category of type derivations and
 49 a category of terms. Since we are interested in categorical semantics with an intersection type
 50 presentation, the first natural thing to do is replacing the category of sets and relations with the
 51 bicategory of *distributors* [6, 10]. Distributor-induced semantics of programming languages
 52 were already presented in [12, 27]. In particular, Fiore, Gambino, Hyland and Winkler
 53 introduced the bicategory of *generalized species of structure* [27], a very rich framework that
 54 generalizes both relational semantics and Joyal’s *combinatorial species* [35, 27, 30, 47]. As
 55 shown in [12, 29], distributors can also lead to a generalization of Scott’s semantics.

56 Mazza, Pellissier and Vial [41], inspired by [42] and Hyland’s project of categorification
 57 of the theory of the λ -calculus [34], presented a general approach to intersection types rooted
 58 in the notion of multicategory. In their framework, the λ -calculus is seen as a 2-operad,
 59 where 2-cells consist of reduction paths. Intersection type systems are seen as a special
 60 kind of *fibrations*. Via a Grothendieck construction, with these fibrations they associate
 61 an *approximation presheaf* that interprets terms as *discrete distributors*. Thanks to this
 62 categorical approach, they are able to prove a parametric normalization theorem for a class
 63 of intersection type systems in a modular and elegant way. Their method relies on a Curry-
 64 Howard style correspondence between intersection type derivations and a kind of λ -terms
 65 *approximants*, the *polyadic terms*. However, their approach does not provide a denotational
 66 model and it does not support subtyping for intersection types. This latter feature is strictly
 67 linked to the fact that approximation presheaves action on types is restricted to discrete
 68 categories [41]. It is then natural to ask what happens when we take the standpoint of
 69 denotational semantics and we take into account categories with non-trivial morphisms.

70 Recently, Tsukada, Asada and Ong [47, 48] presented the *rigid Taylor expansion*¹ se-
 71 mantics for an η -expanded fragment of non-deterministic simply-typed λ -calculus with fixed
 72 point combinator, then extended to probabilistic and quantum computation: the linear
 73 approximants are still polyadic terms. Tsukada, Asada and Ong proved that this semantics
 74 is naturally isomorphic to the generalized species semantics. This time, the standpoint is
 75 well the one of denotational semantics and distributors ranges over groupoids. The groupoid
 76 structure of the model gives the possibility to define an *action* of type isomorphisms on
 77 polyadic terms. A quotient induced by this action guarantees the preservation, up to iso-
 78 morphism, of the semantics under reduction. Concretely one has that $\llbracket M \rrbracket \cong \llbracket N \rrbracket$ whenever
 79 $M \rightarrow N$ and the natural isomorphism is given by *reduction* of polyadic terms.

80 Inspired by these lines of thought, Olimpieri [43] introduced *intersection type distributors*,
 81 a categorized version of intersection type disciplines, where subtyping and denotational
 82 semantics are both taken into account. Intersection type distributors are a *syntactic present-*
 83 *ation* of bicategorical denotational semantics for the λ -calculus given by Kleisli bicategories
 84 of distributors for suitable pseudomonads. Each pseudomonad taken into account gives rise
 85 to a notion of intersection type, with specific resource behavior². The semantics obtained by

¹ The rigid Taylor expansion is a deterministic variant of Ehrhard and Regnier’s Taylor expansion [25, 26].

² It is worth noting that this new semantic setting is not a special case of [41], as standard polyadic terms fail dramatically subject reduction for intersection type distributors. The failure of subject reduction happens because standard polyadic terms [47, 41] cannot encode all the *qualitative* information produced by the *subtyping* feature of intersection type distributors. The interested reader can check a counterexample in Appendix A. However, we believe that from [43] and the present work one can

86 this method is *proof relevant*: given a term M , a type context Δ and a type a , we set

$$87 \quad \mathbb{T}_U(M)(\Delta, a) = \left\{ \begin{array}{c} \tilde{\pi} \\ \vdots \\ \Delta \vdash M : a \end{array} \mid \pi \text{ is a type derivation for } M \right\}$$

88 where $\mathbb{T}_U(M)$ is the intersection type distributor that interprets M in an appropriate
89 category U of types, and $\tilde{\pi}$ is an equivalence class of derivations. The equivalence relation
90 on derivations is induced by the composition of distributors, which generalizes the quotient
91 of [47]. We have that, if $M \rightarrow N$, then $\mathbb{T}_U(M) \cong \mathbb{T}_U(N)$. Categorification then allows us to
92 pass from a semantics of *types* to a semantics of *derivations*. Note that, in our setting, the
93 semantics of a term M associates with every type context Δ and type a the set of derivations
94 for M with conclusion $\Delta \vdash M : a$; while more coarse-grained models such as relational
95 semantics can only say if *there is* a type derivation for M with conclusion $\Delta \vdash M : a$.

96 In the present paper, we introduce *non-idempotent* intersection type distributors in an
97 untyped call-by-push-value setting [33, 24, 39, 46], the *bang calculus*. The call-by-push-value
98 paradigm subsumes call-by-name (CbN) and call-by-value (CbV), from both the operational
99 and denotational semantics standpoints [39, 33]. In this respect, our work is more general
100 than [43] (which considers only the CbN λ -calculus). Moreover, inspired by linear logic, the
101 bang calculus internalizes in the syntax the !-operator, which semantically corresponds to
102 the monadic operator to handle resources. In this way, it is more natural to link syntax
103 and semantics and to disentangle our investigation from the evaluation mechanism. Here we
104 focus on a particular monadic construction (the symmetric strict monoidal completion, see
105 Section 2) and we do not extend the more general and abstract method of [43] to the bang
106 calculus because in this way we can avoid introducing too much categorical background.

107 Our categorical approach allows the introduction of a suitable *category of types*, where
108 morphisms between types are a generalization of *subtyping*. Given a type morphism $a' \rightarrow a$,
109 the intuition is that the type a' somehow *refines* the type a . We prove that non-idempotent
110 intersection type distributors characterize normalization at depth 0 in the bang calculus.
111 Normalization at depth 0 in the bang calculus is a notion that encompasses both CbN
112 solvability [2, 37] and CbV potential valuability [45, 11]. The argument to prove this result is
113 combinatorial and standard (similar results for the bang calculus are proved in [24, 8] using
114 relational semantics), but thanks to the categorified setting we gain a much more fine-grained
115 understanding of the dynamics of type derivations under reduction. Indeed, in our setting,
116 subject reduction and expansion (Theorem 12) clearly open the possibility to define an
117 explicit *deterministic* reduction relation on (equivalent classes) of type derivations, but the
118 investigation of this line of thought is left to future work. We just notice that the substitution
119 operation on type derivations is strictly linked to morphism composition, respecting the basic
120 intuition of categorical semantics: substitution corresponds to composition.

121 **Outline.** Some preliminaries and notations are presented in Section 2. Section 3 shows how
122 the category of distributors Dist can be seen as a generalization of the categories Rel of sets
123 and relations and Polr of preorders. In Section 4 we define a proof-relevant denotational
124 model of the bang calculus in Dist as a generalization of non-idempotent intersection type
125 systems and we prove a semantic characterization of depth 0 normalization in the bang
126 calculus. Conclusions are in Section 5.

define an approximation presheaf in the sense of [41] that is also a denotational semantics, *i.e.*, a pseudomorphism in that context, which supports subtyping. This generalization is left to future work, but is clearly in the spirit of the general Grothendieck construction of [41].

Terms:	$S, T, U ::= x \mid \lambda x.S \mid ST \mid S^!$	(set: $!\Lambda$)
Contexts:	$\mathbf{C} ::= [\cdot] \mid \lambda x.\mathbf{C} \mid \mathbf{C}S \mid S\mathbf{C} \mid \mathbf{C}^!$	(set: $!\Lambda_{\mathbf{C}}$)
Ground Contexts:	$\mathbf{G} ::= [\cdot] \mid \lambda x.\mathbf{G} \mid \mathbf{G}S \mid S\mathbf{G}$	(set: $!\Lambda_{\mathbf{G}}$)
Root-step:	$(\lambda x.S)T^! \mapsto_{\mathbf{b}} S\{T/x\}$	
$\rightarrow_{\mathbf{b}}$ -reduction:	$S \rightarrow_{\mathbf{b}} T \Leftrightarrow \exists \mathbf{C} \in !\Lambda_{\mathbf{C}}, \exists S', T' \in !\Lambda : S = \mathbf{C}[S'], T = \mathbf{C}[T'], S' \mapsto_{\ell} T'$	
$\rightarrow_{\mathbf{b}_{\mathbf{g}}}$ -reduction:	$S \rightarrow_{\mathbf{b}_{\mathbf{g}}} T \Leftrightarrow \exists \mathbf{G} \in !\Lambda_{\mathbf{G}}, \exists S', T' \in !\Lambda : S = \mathbf{G}[S'], T = \mathbf{G}[T'], S' \mapsto_{\ell} T'$	

■ **Figure 1** The bang calculus: its syntax and reduction rules.

2 Preliminaries

127

128 **The bang calculus.** The syntax and operational semantics of the *bang calculus* [33] are
 129 defined in Figure 1. Terms are built up from a countably infinite set of *variables* (denoted
 130 by x, y, z, \dots). Terms of the form $S^!$ (resp. $\lambda x.S$; ST) are called *boxes* (resp. *abstractions*;
 131 *(linear) applications*). The set of boxes is denoted by $!\Lambda_!$. The set of free variables of a term
 132 S , denoted by $\text{fv}(S)$, is defined as expected, λ being the only binding construct. All terms
 133 are considered up to α -conversion. Given $S, T \in !\Lambda$ and a variable x , $S\{T/x\}$ denotes the
 134 term obtained by the *capture-avoiding substitution* of T for each free occurrence of x in S .

135 *Contexts* \mathbf{C} and (with exactly one hole $[\cdot]$) are defined in Figure 1. We write $\mathbf{C}[S]$ for the
 136 term obtained by capture-allowing substitution of the term S for the hole $[\cdot]$ in the context
 137 \mathbf{C} . *Ground contexts* \mathbf{G} are the restriction to contexts where the hole is not inside any $!$.

138 The *bang calculus* is the set $!\Lambda$ endowed with reduction $\rightarrow_{\mathbf{b}}$ (Figure 1), which is confluent
 139 [33]. Intuitively in the root-step $\mapsto_{\mathbf{b}}$ the box-construct $!$ marks the only terms that can be
 140 erased and duplicated: a β -like redex $(\lambda x.S)T$ can be fired only when its argument is a box,
 141 *i.e.* $T = U^!$: if it is so, the content U of the box T replaces any free occurrence of x in S .

142 Reduction $\rightarrow_{\mathbf{b}_{\mathbf{g}}} \subseteq \rightarrow_{\mathbf{b}}$ is said *at depth 0* and defined as the closure of $\mapsto_{\mathbf{b}}$ under ground
 143 contexts (see Figure 1): it does not reduce inside boxes. It has the diamond-property [33].

144 ► **Example 1.** Let $\Delta = \lambda x.xx^!$. Then $\Delta\Delta^! \rightarrow_{\mathbf{b}_{\mathbf{g}}} \Delta\Delta^! \rightarrow_{\mathbf{b}_{\mathbf{g}}} \dots$ (and so $\Delta\Delta^! \rightarrow_{\mathbf{b}} \Delta\Delta^! \rightarrow_{\mathbf{b}} \dots$).

145 ► **Definition 2 (Clash).** A clash is a term of the form $S^!T$ or $T(\lambda x.S)$.

146 Let $S \in !\Lambda$: S is clash-free if and only if it contains no clash; S is clash-free at depth 0
 147 if and only if each clash occurring in S is under the scope of a $!$.

148 For instance, $(\lambda z.x)(x^!y)^!$ is clash-free at depth 0 but not clash-free. Roughly, a clash is
 149 a “meaningless” term that cannot inherently be typed (see [24, 8]): boxes cannot be applied,
 150 abstractions cannot be the argument of an application.

151 The bang calculus can be extended (see [24]) with the reduction $\rightarrow_{\sigma} = \rightarrow_{\sigma_1} \cup \rightarrow_{\sigma_2} \cup \rightarrow_{\sigma_3}$
 152 where \rightarrow_{σ_1} , \rightarrow_{σ_2} and \rightarrow_{σ_3} are the contextual closure of the following rules, respectively:

$$153 \quad (\lambda x.S)TU \mapsto_{\sigma_1} (\lambda x.SU)T \quad (\lambda y.\lambda x.S)T \mapsto_{\sigma_2} \lambda x.(\lambda y.S)T \quad T((\lambda x.S)U) \mapsto_{\sigma_3} (\lambda x.TS)U$$

154 with $x \notin \text{fv}(U)$ in \mapsto_{σ_1} and \mapsto_{σ_3} , while $x \notin \text{fv}(T) \cup \{y\}$ in \mapsto_{σ_2} . We set $\rightarrow_{\mathbf{b}\sigma} = \rightarrow_{\mathbf{b}} \cup \rightarrow_{\sigma}$ and
 155 $\rightarrow_{\mathbf{b}_{\mathbf{g}}\sigma} = \rightarrow_{\mathbf{b}_{\mathbf{g}}} \cup \rightarrow_{\sigma}$, where $\rightarrow_{\sigma} = \rightarrow_{\sigma_1} \cup \rightarrow_{\sigma_2} \cup \rightarrow_{\sigma_3}$ and \rightarrow_{σ_i} is the closure under ground
 156 contexts of \mapsto_{σ_i} , for $i \in \{1, 2, 3\}$. Reductions \rightarrow_{σ} and $\rightarrow_{\sigma_{\mathbf{g}}}$ are strongly normalizing [24] and
 157 can “unveil” hidden b-redexes and hidden clashes. For instance,

$$158 \quad ((\lambda x.\Delta)x)\Delta^! \rightarrow_{\sigma_1} (\lambda x.\Delta\Delta^!)x \quad x((\lambda y.\lambda x.z)y) \rightarrow_{\sigma_2} x(\lambda x.(\lambda y.z)y)$$

160 where $((\lambda x.\Delta)x)\Delta^!$ is b-normal but $(\lambda x.\Delta\Delta^!)x$ is not ($\rightarrow_{\mathbf{b}_{\mathbf{g}}}$ can fire the b-redex $\Delta\Delta^!$), and
 161 $x((\lambda y.\lambda x.z)y)$ is clash-free but $x(\lambda x.(\lambda y.z)y)$ is not (not even at depth 0).

162 **Integers and Permutations.** For $n \in \mathbb{N}$, we set $[n] = \{1, \dots, n\}$, so $[0] = \emptyset$. The set of
 163 permutations over $[n]$ is denoted by S_n . We define the category \mathbb{P} of integers and permutations:

164 ■ the objects of \mathbb{P} are $\text{ob}(\mathbb{P}) = \{[n] \mid n \in \mathbb{N}\}$; the identity on $[n]$ is denoted by 1_n ;

165 ■ the homset from $[n]$ to $[m]$ is $\mathbb{P}[[n], [m]] = \begin{cases} S_n & \text{if } n = m \\ \emptyset & \text{otherwise;} \end{cases}$

166 ■ the category \mathbb{P} is symmetric strict monoidal, with tensor product given by addition:
 167 $[n] \oplus [m] = [n + m]$. Given $\sigma \in S_{k_1}$ and $\tau \in S_{k_2}$, we define $\sigma \oplus \tau \in S_{k_1+k_2}$ as

$$168 \quad (\sigma \oplus \tau)(i) = \begin{cases} \sigma(i) & \text{if } 1 \leq i \leq k_1 \\ \tau(i - k_1) + k_1 & \text{otherwise.} \end{cases}$$

169 Given $k_1, \dots, k_n \in \mathbb{N}$ and $\sigma \in S_n$, we define $\bar{\sigma}: [\sum_{i \in [n]} k_i] \rightarrow [\sum_{i \in [n]} k_{\sigma(i)}]$ as $\bar{\sigma}(\sum_{r=1}^{l-1} k_r +$
 170 $p) = \sum_{r=1}^{l-1} k_{\sigma(r)} + p$, where $l \in [n]$ and $1 \leq p \leq k_{\sigma(l)}$.

171 **Symmetric strict monoidal completion.** For a list $\vec{a} = \langle a_1, \dots, a_k \rangle$, we set $\text{len}(\vec{a}) = k$.
 172 Lists are denoted by $\vec{a}, \vec{b}, \vec{c}, \dots$, concatenation of two lists \vec{a} and \vec{b} is denoted by $\vec{a} \oplus \vec{b}$.

173 Let A be a small category. For each object $a \in \text{ob}(A)$, the identity morphism on a is
 174 denoted by 1_a . The *symmetric strict monoidal completion* $!A$ of A is the category:

175 ■ $\text{ob}(!A) = \{\langle a_1, \dots, a_n \rangle \mid a_i \in A \text{ and } n \in \mathbb{N}\}$;

176 ■ $!A[\langle a_1, \dots, a_n \rangle, \langle a'_1, \dots, a'_{n'} \rangle] = \begin{cases} \{\langle \sigma, f_1, \dots, f_n \rangle \mid f_i: a_i \rightarrow a'_{\sigma(i)}, \sigma \in S_n\} & \text{if } n = n'; \\ \emptyset & \text{otherwise;} \end{cases}$

177 ■ for $\vec{a} = \langle a_1, \dots, a_n \rangle \in \text{ob}(!A)$, the identity on \vec{a} is $1_{\vec{a}} = \langle 1_n, 1_{a_1}, \dots, 1_{a_n} \rangle$;

178 ■ for $f = \langle \sigma, f_1, \dots, f_n \rangle: \vec{a} \rightarrow \vec{b}$ and $g = \langle \tau, g_1, \dots, g_n \rangle: \vec{b} \rightarrow \vec{c}$, the composition is $g \circ f =$
 179 $\langle \tau\sigma, g_{\sigma(1)} \circ f_1, \dots, g_{\sigma(n)} \circ f_n \rangle$;

180 ■ the monoidal structure is given by list concatenation. The tensor product is symmetric,
 181 with symmetries given by the morphisms of the shape (where $\sigma: [n] \rightarrow [n]$ is a permutation)

$$182 \quad \langle \sigma, \vec{1} \rangle: \langle a_1, \dots, a_n \rangle \rightarrow \langle a_{\sigma(1)}, \dots, a_{\sigma(n)} \rangle$$

183 Given a permutation $\sigma: [n] \rightarrow [n]$ and $\vec{a}_1, \dots, \vec{a}_n \in \text{ob}(!A)$ with $\text{len}(\vec{a}_i) = k_i$ we define
 184 $\sigma^*: \bigoplus_{i=1}^n \vec{a}_i \rightarrow \bigoplus_{i=1}^n \vec{a}_{\sigma(i)}$ as $\langle \bar{\sigma}, 1_{a_1}, \dots, 1_{a_k} \rangle$, where $k = \sum_{i \in [n]} k_i$.

185 We use the following shortenings: $!A^n = (!A)^n$ and $!A^{\text{op}} = (!A)^{\text{op}}$.

186 **Bicategory.** We assume the reader to be familiar with bicategories [3, 6] and two-dimensional
 187 monads [5]. Some basic notions are briefly recalled in Appendix A. For a diagram $F: C \rightarrow D$,
 188 its colimit is denoted by $\varinjlim_{c \in C} F(c)$. Given a bicategory \mathcal{C} , \mathcal{C}^{op} is the bicategory obtained by

189 reversing the 1-cells of \mathcal{C} , but not the 2-cells.

190 **3 Rel, Polr, Dist**

191 We sketches the structure of some categories providing denotational models of linear logic.

192 We use linear logic notations for cartesian products, comonads modelling exponentials, etc.

193 **Rel.** A simple model of linear logic is the category Rel of sets and relations. It is a prototype
 194 of *quantitative* semantics: the interpretation of a program gives information about its resource
 195 consumption during computation. Intuitively, an element in a set represents a non-idempotent
 196 intersection type. For the bang calculus, this model has been studied in [24, 33].

38:6 Categorifying Non-Idempotent Intersection Types

197 Objects of Rel are sets, and morphisms of Rel are binary relations. Identities are diagonal
198 relations. Composition of morphisms in Rel is the usual composition of relations

$$199 \quad g \circ f = \{\langle x, z \rangle \mid \exists y \in Y : \langle x, y \rangle \in f, \langle y, z \rangle \in g\} \text{ for } f \subseteq X \times Y \text{ and } g \subseteq Y \times Z.$$

200 For $X_1, X_2 \in \text{ob}(\text{Rel})$, the cartesian product $X_1 \& X_2$ in Rel is the disjoint union of sets
201 $X_1 \sqcup X_2 = (\{1\} \times X_1) \cup (\{2\} \times X_2)$, where projections $\pi_i: X_1 \& X_2 \rightarrow X_i$ (for $i \in \{1, 2\}$) are
202 injections $\{\langle \langle i, x \rangle, x \rangle \mid x \in X_i\}$, and the terminal (and initial) object \top is the empty set \emptyset .

203 Rel is a *symmetric monoidal* category, where the tensor $X \otimes Y$ is the cartesian product
204 of sets $X \times Y$ and its unit $\mathbf{1}$ is an arbitrary singleton set. It is *closed*, with $X \multimap Y = X \times Y$
205 and evaluation $\text{ev}_{X,Y}: (X \multimap Y) \times X \rightarrow Y$ defined by $\{\langle \langle \langle x, y \rangle, x \rangle, y \rangle \mid x \in X, y \in Y\}$.

206 Rel comes with an *exponential comonad* $(!, \text{der}, \text{dig})$. The functor $!$ is given by $!X = \mathcal{M}_f(X)$
207 (finite multisets over X) and, for a morphism $f \in \text{Rel}[X, Y]$, $!f = \{\langle [x_1, \dots, x_n], [y_1, \dots, y_n] \rangle \mid$
208 $n \in \mathbb{N}, \langle x_1, y_1 \rangle, \dots, \langle x_n, y_n \rangle \in f\}$. Dereliction $\text{der}_X \in \text{Rel}[!X, X]$ is $\{\langle [x], x \rangle \mid x \in X\}$, and
209 digging $\text{dig}_X \in \text{Rel}[!X, !!X]$ is $\{\langle m_1 + \dots + m_k, [m_1, \dots, m_k] \rangle \mid m_1, \dots, m_k \in !X\}$ (for two
210 finite multisets $\bar{a} = [a_1, \dots, a_k]$ and $\bar{b} = [b_1, \dots, b_n]$, we set $\bar{a} + \bar{b} = [a_1, \dots, a_k, b_1, \dots, b_n]$).

211 **Polr.** To work within a more informative setting, providing not only *quantitative*, but also
212 *qualitative* information, consider the category Polr of preordered sets and monotonic relations
213 [21, 23]. Intuitively, given two types a and b , if $a \leq b$ then a is an approximant of b . All the
214 constructions in Polr are a refinement and generalization of the ones for Rel .

215 In Polr , objects are preordered sets; a morphism f from $\mathcal{X} = \langle |\mathcal{X}|, \leq_{\mathcal{X}} \rangle$ to $\mathcal{Y} = \langle |\mathcal{Y}|, \leq_{\mathcal{Y}} \rangle$
216 is a *monotonic* relation³ from $|\mathcal{X}|$ to $|\mathcal{Y}|$, *i.e.*, if $\langle x, y \rangle \in f$ with $x' \leq_{\mathcal{X}} x$ and $y \leq_{\mathcal{Y}} y'$ then
217 $\langle x', y' \rangle \in f$. The identity at \mathcal{X} is $\{\langle x, x' \rangle \mid x \leq_{\mathcal{X}} x'\}$. Composition preserves monotonicity.

218 In Polr the cartesian product $\mathcal{X}_1 \& \mathcal{X}_2$ is the disjoint union of sets $|\mathcal{X}_1| \sqcup |\mathcal{X}_2|$ with the
219 preorder $\leq_{\mathcal{X}_1} \sqcup \leq_{\mathcal{X}_2}$ defined as $\langle i, x \rangle \leq_{\mathcal{X}_1 \& \mathcal{X}_2} \langle j, y \rangle$ if $i = j$ and $x \leq_{\mathcal{X}_i} y$. The terminal object
220 \top is \emptyset with the empty order. Projections $\pi_i: \mathcal{X}_1 \& \mathcal{X}_2 \rightarrow \mathcal{X}_i$ are $\pi_i = \{\langle \langle i, x \rangle, x' \rangle \mid x \leq_{\mathcal{X}_i} x'\}$.

221 Polr has a symmetric monoidal structure. The tensor $\mathcal{X}_1 \otimes \mathcal{X}_2$ is the cartesian product of
222 sets with the product order. The endofunctor $\mathcal{X} \otimes _$ admits a right adjoint $_ \multimap \mathcal{Y}$ defined
223 as follows: $|\mathcal{X} \multimap \mathcal{Y}| = |\mathcal{X}| \times |\mathcal{Y}|$ and $\langle x, y \rangle \leq_{\mathcal{X} \multimap \mathcal{Y}} \langle x', y' \rangle$ if $x' \leq_{\mathcal{X}} x$ and $y \leq_{\mathcal{Y}} y'$. The
224 evaluation morphism $\text{ev}_{\mathcal{X}_1, \mathcal{X}_2}: (\mathcal{X}_1 \multimap \mathcal{X}_2) \& \mathcal{X}_1 \rightarrow \mathcal{X}_2$ is $\{\langle \langle \langle x, y \rangle, x' \rangle, y' \rangle \mid x \leq_{\mathcal{X}_1} x', y \leq_{\mathcal{X}_2} y' \rangle\}$.

225 Polr has exponential comonad $(!, \text{der}, \text{dig})$.⁴ The endofunctor $!: \text{Polr} \rightarrow \text{Polr}$ is given by
226 $!\mathcal{X} = \langle \mathcal{M}_f(|\mathcal{X}|), \leq_{\mathcal{X}} \rangle$ with $[x_1, \dots, x_n] \leq_{!\mathcal{X}} [x'_1, \dots, x'_n]$ if $n = n'$ and there is $\sigma \in S_n$ such
227 that $x_i \leq_{\mathcal{X}} x'_{\sigma(i)}$ for all $1 \leq i \leq n$; for $f \in \text{Polr}[\mathcal{X}, \mathcal{Y}]$, we set $!f = \{\langle [x_1, \dots, x_n], [y_1, \dots, y_k] \rangle \mid$
228 $\langle x_i, y_i \rangle \in f, k \in \mathbb{N}\}$. Dereliction $\text{der}_{\mathcal{X}}: !\mathcal{X} \rightarrow \mathcal{X}$ is $\{\langle [x], x' \rangle \mid x \leq_{\mathcal{X}} x'\}$, and digging
229 $\text{dig}_{\mathcal{X}}: !\mathcal{X} \rightarrow !!\mathcal{X}$ is $\{\langle m, [m_1, \dots, m_k] \rangle \mid m \leq_{!\mathcal{X}} m_1 + \dots + m_k\}$.

230 Rel is the full subcategory of Polr where objects are sets equipped with the discrete order.

231 **Polr as a model of the bang calculus.** A categorical model of the bang calculus [23, 24]
232 consists of a \star -autonomous category $(A, \otimes, I, \multimap, (-)^{\perp})$, cartesian with product $\&$ and
233 terminal object \top (and, by \star -autonomy, cocartesian with coproduct \oplus and initial object 0),
234 endowed with a comonad $(!, \text{der}, \text{dig})$ with suitable Seely isomorphisms [23, 33]. Also, we

³ In [21, 23], monotonicity is slightly different, so that the type system generated by the model is covariant on the left of \vdash and contravariant on the right of \vdash . With our definition, the type system generated by the model is contravariant on the left of \vdash and covariant on the right of \vdash , in accordance with [1].

⁴ Akin to [21] and unlike [23], our exponential comonad is based on finite multiset construction. But our preorder on $!\mathcal{X}$ is different from [21]: there $[a] \leq_{!\mathcal{X}} [a, a]$ (idempotency is a sort of approximation), here $[a]$ and $[a, a]$ are incomparable, so that approximation is completely independent from idempotency.

Types:

$$a := x \in \mathcal{X} \mid [a_1, \dots, a_k] \multimap a \mid [a_1, \dots, a_k]$$

Preorder \leq_U in U :

$$\frac{x \leq_{\mathcal{X}} x' \quad m' \leq_U m \quad a \leq_U a'}{x \leq_U x' \quad (m \multimap a) \leq_U (m' \multimap a')}$$

$$\frac{\sigma \in S_k \quad a_1 \leq_U a'_{\sigma(1)} \quad \dots \quad a_k \leq_U a'_{\sigma(k)}}{[a_1, \dots, a_k] \leq_U [a'_1, \dots, a'_k]}$$

Derivation rules:

$$\frac{a' \leq_U a}{x_1 : [], \dots, x_i : [a'], \dots, x_n : [] \vdash x_i : a}$$

$$\frac{\Gamma \vdash S : m \multimap a \quad \Gamma' \vdash T : m \quad \Delta \leq_{U^n} \Gamma \otimes \Gamma'}{\Delta \vdash ST : a}$$

$$\frac{\Gamma_1 \vdash S : a_1 \quad \dots \quad \Gamma_k \vdash S : a_k \quad \Delta \leq_{U^n} \bigotimes_{i=1}^k \Gamma_i}{\bigotimes_{i=1}^k \Gamma_i \vdash S^! : [a_1, \dots, a_k]}$$

$$\frac{\Delta, x : m \vdash S : a}{\Delta \vdash \lambda x. S : m \multimap a}$$

■ **Figure 2** Non-idempotent intersection type system \mathcal{R}_{\leq} associated with the preorder U in Polr.

235 require that $0 \cong \top$. An *extensional model* of the bang calculus is then an object $U \in \text{ob}(A)$
 236 such that $U \cong !U \& (!U \multimap U)$. To have a *non-extensional model* for the bang calculus a
 237 retraction $!U \& (!U \multimap U) \triangleleft U$ is enough.

238 We build a retraction in the category Polr. We define a family of preorders as follows:

$$239 \quad U_0 = \mathcal{X} \text{ (any preorder)} \quad U_{n+1} = !U_n \sqcup ((!U_n \multimap U_n) \sqcup \mathcal{X}) \quad (1)$$

240 We define a family of canonical inclusions $(\iota_n : U_n \hookrightarrow U_{n+1})_{n \in \mathbb{N}}$ as $\iota_0 = \iota_{\mathcal{X}}$ (the inclusion
 241 $\mathcal{X} \hookrightarrow !\mathcal{X} \sqcup ((!\mathcal{X} \multimap \mathcal{X}) \sqcup \mathcal{X})$) and $\iota_{n+1} = !\iota_n \sqcup ((!\iota_n \multimap \iota_n) \sqcup 1_{\mathcal{X}})$, so the preorder U_n is
 242 just the restriction to the elements of U_n of the preorder U_{n+1} . We set $U = \varinjlim_{n \in \mathbb{N}} U_n$, that

243 is a directed colimit of the directed diagram $\langle \iota_i \rangle_{i \in \mathbb{N}}$. It is easy to check that there exists a
 244 canonical inclusion $\iota : !U \sqcup (!U \multimap U) \hookrightarrow U$ and that we have a retraction $!U \& (!U \multimap U) \triangleleft U$.

245 We can define the interpretation of the terms of the bang calculus in Polr. Let $S \in !\Lambda$
 246 and $\text{fv}(S) \subseteq \vec{x} = \langle x_1, \dots, x_n \rangle$ with the x_i 's pairwise distinct. The *semantics* (or *denotation*)
 247 of S is a monotonic relation $\llbracket S \rrbracket_{\vec{x}} : !U^{\otimes n} \rightarrow U$ defined by induction as follows:

- 248 ■ $\llbracket x_i \rrbracket_{\vec{x}} = \{ \langle \langle [], \dots, [a'], \dots, [] \rangle, a \rangle \mid a' \leq a \}$ (a' is in the i^{th} position in $\langle [], \dots, [a'], \dots, [] \rangle$);
- 249 ■ $\llbracket \lambda y. T \rrbracket_{\vec{x}} = \{ \langle \Delta, \iota(\langle m, a \rangle) \rangle \mid \langle \Delta \oplus \langle m \rangle, a \rangle \in \llbracket T \rrbracket_{\vec{x} \oplus \langle y \rangle} \}$, where $y \notin \vec{x}$;
- 250 ■ $\llbracket ST \rrbracket_{\vec{x}} = \bigcup_{m \in !U} \bigcup_{\Gamma, \Gamma' \in U^n} \{ \langle \Delta, a \rangle \mid \langle \Gamma, \iota(\langle m, a \rangle) \rangle \in \llbracket S \rrbracket_{\vec{x}}, \langle \Gamma', \iota(m) \rangle \in \llbracket T \rrbracket_{\vec{x}} \text{ and } \Delta \leq_{U^n} \Gamma \otimes \Gamma' \}$;
- 252 ■ $\llbracket T^! \rrbracket_{\vec{x}} = \bigcup_{k \in \mathbb{N}} \bigcup_{\Gamma_1, \dots, \Gamma_k \in U^n} \{ \langle \Delta, [a_1, \dots, a_k] \rangle \mid \langle \Gamma_i, a_i \rangle \in \llbracket T \rrbracket_{\vec{x}} \text{ and } \Delta \leq_{U^n} \bigotimes_{i=1}^k \Gamma_i \}$
 253 where if $\Gamma = \langle m_1, \dots, m_n \rangle$ and $\Gamma' = \langle m'_1, \dots, m'_n \rangle$ then $\Gamma \otimes \Gamma' = \langle m_1 + m'_1, \dots, m_n + m'_n \rangle$.

254 Ehrhard [23] showed this is a denotational semantics. By setting $m \multimap a = \langle m, a \rangle \in !U \times U$,
 255 we can give a type-theoretic description of the preorder U as in Figure 2. Such a type system
 256 \mathcal{R}_{\leq} is similar to de Carvalho's non-idempotent intersection type system \mathcal{R} [16, 17]. The
 257 main difference is that in \mathcal{R}_{\leq} types are elements of a *preorder* U (an object of Polr), while in
 258 \mathcal{R} types are elements of a *set* U (an object of Rel). The additional information provided by
 259 the preorder accounts for *approximation*: if $a \leq_U b$ then the type a approximates the type b .
 260 This is evident in the rule for the variable in Figure 2: a' can be seen as a *subtype* of a .

261 By easy inspection of the definition, $\langle \Delta, a \rangle \in \llbracket S \rrbracket_{\vec{x}}$ if and only if $\Delta \vdash S : a$. In other
 262 words, the semantics of a term S is the set of conclusions of the type derivations for S . The
 263 semantics is then a *semantics of types* in the non-idempotent intersection type system \mathcal{R}_{\leq} .

264 We now try to shift our standpoint. In system \mathcal{R}_{\leq} , let us try to define a *semantics of*
 265 *proofs*. Given a term S , a context Δ and type a , we set $\llbracket S \rrbracket_{\vec{x}}(\Delta, a) = \left\{ \begin{array}{l} \pi \\ \vdots \\ \Delta \vdash M : a \end{array} \mid \pi \in \mathcal{R}_{\leq} \right\}$.

266 It is easy to see that this proof-relevant structure is not a denotational semantics. Indeed,

38:8 Categorifying Non-Idempotent Intersection Types

267 the reduction over type derivations in system \mathcal{R}_{\leq} is non-deterministic, since it deals with
 268 multisets. Take $(\lambda z.(yz^!)z^!)S^! \rightarrow_{\ell} (yS^!)S^!$ and the following type derivation:

$$\begin{array}{c}
 \frac{\frac{\frac{y: [[a] \multimap [a] \multimap c] \vdash y: [a] \multimap [a] \multimap c}{y: [[a] \multimap [a] \multimap c], z: [a] \vdash yz^!: [a] \multimap c} \quad \frac{z: [a] \vdash z^!: [a]}{z: [a] \vdash z^!: [a]}}{y: [[a] \multimap [a] \multimap c], z: [a, a] \vdash (yz^!)z^!: c} \quad \frac{\pi_1 \quad \pi_2}{\Gamma_1 \vdash S: a \quad \Gamma_2 \vdash S: a}}{\Gamma_1 \otimes \Gamma_2 \vdash S^!: [a, a]} \\
 \frac{\Gamma_1 \otimes \Gamma_2, y: [[a] \multimap [a] \multimap c] \vdash (\lambda z.(yz^!)z^!)S^!: c}{\Gamma_1 \otimes \Gamma_2, y: [[a] \multimap [a] \multimap c] \vdash (\lambda z.(yz^!)z^!)S^!: c}
 \end{array}$$

270 Suppose that x and z are not free in S and that $\pi_1 \neq \pi_2$. Then if we consider the reduct
 271 $(yS^!)S^!$ we have two possible choices for the typing, $\pi\{\pi_1/z_1, \pi_2/z_2\}$ or $\pi\{\pi_2/z_1, \pi_1/z_2\}$. This
 272 non-determinism stems from the multiset structure, but we shall see that simply passing to
 273 a list-oriented framework does not solve the problem. A natural way to make this kind of
 274 structure a denotational semantics is the lifting to Set enriched distributors.

275 **From Rel and Polr to Dist.** We recall a basic but pivotal fact: a relation $f \subseteq X \times Y$ can be
 276 identified with its characteristic function $\chi_f: X \times Y \rightarrow \mathbf{2}$ where $\mathbf{2} = \{0, 1\}$ is the two-element
 277 boolean algebra with sum (join) and product (meet). Composition is then defined as

$$278 \quad \chi_{g \circ f}(x, z) = \sum_{y \in Y} \chi_g(y, z) \cdot \chi_f(x, y) \quad \text{where } \chi_f: X \times Y \rightarrow \mathbf{2} \text{ and } \chi_g: Y \times Z \rightarrow \mathbf{2}. \quad (2)$$

279 All the constructions in Rel and Polr can be reformulated in this characteristic function
 280 perspective. For instance, in Rel, the identity at X becomes the characteristic function of X .

281 In Polr, a monotonic relation f from $\mathcal{X} = \langle |\mathcal{X}|, \leq_{\mathcal{X}} \rangle$ to $\mathcal{Y} = \langle |\mathcal{Y}|, \leq_{\mathcal{Y}} \rangle$ can be seen as a
 282 monotonic characteristic function $\chi_f: \mathcal{X}^{\text{op}} \times \mathcal{Y} \rightarrow \mathbf{2}$, where $\mathcal{X}^{\text{op}} = \langle X, \geq_{\mathcal{X}} \rangle$ and $\mathbf{2}$ is endowed
 283 with the boolean order. Any preorder $\mathcal{X} = \langle |\mathcal{X}|, \leq_{\mathcal{X}} \rangle$ forms a category where $\text{ob}(\mathcal{X}) = |\mathcal{X}|$
 284 and $\mathcal{X}[x, x']$ is a singleton (if $x \leq_{\mathcal{X}} x'$) or the empty set (otherwise), so \mathcal{X}^{op} is the opposite
 285 category of \mathcal{X} . Thus, $\chi_f: \mathcal{X}^{\text{op}} \times \mathcal{Y} \rightarrow \mathbf{2}$ is a bifunctor, contravariant in \mathcal{X} and covariant in
 286 \mathcal{Y} . The semantics of a term S is then a Polr morphism $\llbracket S \rrbracket_{\bar{x}}: (!U^{\otimes n})^{\text{op}} \times U \rightarrow \mathbf{2}$.

287 It is then natural to generalize the characteristic function viewpoint to generic categories,
 288 which gives rise to the notion of *distributor* (also known as *profunctors*).

289 **Dist.** For two small categories A, B , a *distributor* $F: A \multimap B$ is a functor $F: A^{\text{op}} \times B \rightarrow \text{Set}$.
 290 Composition of distributors relies on the notion of *coend*, a kind of colimit (a coequalizer).

291 **► Definition 3 (Coend, [40]).** Let $F: C^{\text{op}} \times C \rightarrow D$ be a functor. A *cowedge* for F is an object
 292 $T \in D$ together with a family of morphisms $w_c: F(c, c) \rightarrow T$ such that diagram (3) below
 293 commutes, for $f: c \rightarrow c'$. A *coend* for F , denoted by $\int^{c \in C} F(c, c)$, is a universal cowedge.

$$\begin{array}{ccc}
 F(c', c) & \xrightarrow{F(f, 1)} & F(c, c) \\
 \downarrow F(1, f) & & \downarrow w_c \\
 F(c', c') & \xrightarrow{w_{c'}} & T
 \end{array} \quad (3)$$

295 We now define the bicategory *Dist* of *distributors*. For a proper presentation of the
 296 structure of this bicategory we refer to [10, 12, 27, 30].

297 **■** *0-cells* are small categories $A, B, C \dots$; *1-cells* $F: A \multimap B$ are distributors, *i.e.* functors
 298 $F: A^{\text{op}} \times B \rightarrow \text{Set}$; *2-cells* $\alpha: F \Rightarrow G$ are natural transformations.

- 299 ■ Given any 0-cells A and B , 1-cells and 2-cells are organized as a category $\text{Dist}(A, B)$.
 300 Composition $\alpha \star \beta$ in $\text{Dist}(A, B)$ is called *vertical composition*. We define the *zero*
 301 *distributor* $\emptyset_{A,B} \in \text{ob}(\text{Dist}(A, B))$ as $\emptyset_{A,B}(a, b) = \emptyset$ for all $a \in \text{ob}(A)$ and $b \in \text{ob}(B)$.
 302 ■ For $A \in \text{Dist}$, the identity $1_A: A \rightarrow A$ is Yoneda's embedding $1_A(a', a) = A(a', a)$.
 303 ■ For 1-cells $F: A \rightarrow B$ and $G: B \rightarrow C$, their composition is given by

$$304 \quad (G \circ F)(a, c) = \int^{b \in B} G(b, c) \times F(a, b)$$

305 Note the analogy with (2). Composition is only associative up to canonical isomorphisms.
 306 For this reason Dist is a bicategory [6].

- 307 ■ The cartesian product $A \& B$ is the disjoint union $A \sqcup B$ of categories. The terminal
 308 object \top is given by the empty category. The bicategory Dist admits also coproducts,
 309 with $A \oplus B = A \sqcup B$ (the canonical inclusions are denoted by ι_A and ι_B) and $0 = \top$.
 310 ■ There is a symmetric monoidal structure on Dist given by the cartesian product of
 311 categories: $A \otimes B = A \times B$, with any one-object category as a unit. The bicategory of
 312 distributors is monoidal closed, with linear exponential object $A \multimap B = A^{\text{op}} \times B$.

313 The symmetric strict monoidal completion of a small category A (Section 2) lifts to
 314 an endofunctor in Cat , by setting $!F(\langle a_1, \dots, a_n \rangle) = \langle F(a_1), \dots, F(a_n) \rangle$ for any functor
 315 $F: A \rightarrow B$. The endofunctor $!$ can be extended to Dist , determining a pseudocomonad
 316 $(!, \text{dig}_A, \text{der}_A)$ on Dist [27, 30]. The two components of the pseudocomonad are defined
 317 as follows: $\text{dig}_A(\vec{a}, \langle \vec{a}_1, \dots, \vec{a}_n \rangle) = !A[\vec{a}, \bigoplus_{i=1}^n \vec{a}_i]$ and $\text{der}_A(\vec{a}, a) = !A[\vec{a}, \langle a \rangle]$. The Kleisli
 318 bicategory $Kl(!)(\text{Dist})$ is the bicategory of *categorical symmetric sequences* [30], biequivalent
 319 to the bicategory of *generalized species of structure* [27, 28]. There are Seelye equivalences
 320 $!(A \& B) \simeq !A \times !B$ and $!\top \simeq 1$, pseudonatural in both A and B [27].

321 **4 A Type-Theoretic Non-Extensional Model for the Bang Calculus**

322 **Distributors-Induced Model for the Bang Calculus.** The bicategory of distributors fulfills
 323 a bicategorical generalization of the categorical model of the bang calculus shown in Section
 324 3.⁵ However, we leave the proper development of a general notion of bicategorical model for
 325 the bang calculus to future work, since the notion of symmetric monoidal bicategory is highly
 326 non-trivial. For our purpose, it is enough to present a denotational model inside a particular
 327 bicategory, *i.e.*, the bicategory of distributors. A denotational model in this setting will be
 328 an interpretation of bang terms as suitable 1-cells, such that $\llbracket S \rrbracket_{\vec{x}} \cong \llbracket T \rrbracket_{\vec{x}}$ if $S \rightarrow_{\ell} T$. In
 329 particular, we want $\llbracket S \rrbracket_{\vec{x}}: (!U^{\text{op}})^{\text{op}} \times U \rightarrow \text{Set}$ (for $\text{len}(\vec{x}) = n$), with $!U \& (!U \multimap U) \triangleleft U$.
 330 The intuition is that, in Dist , 0-cells represent types (and in our untyped setting, they satisfy
 331 a retraction), 1-cells represent type derivations and 2-cells represent reduction on derivations.

332 We build the retraction in Dist , in analogy with the construction (1) in Polr . Indeed, they
 333 are both special cases of the free-algebra construction for an (unpointed) endofunctor [36]. We
 334 recall that, in Dist , $A \& B = A \sqcup B$, $A \otimes B = A \times B$ (so $A^{\otimes n} = A^n$) and $A \multimap B = A^{\text{op}} \times B$.

335 ► **Definition 4.** Let A be a small category. We define a family of small categories $(U_n)_{n \in \mathbb{N}}$ by:

$$336 \quad U_0 = A \quad U_{n+1} = !U_n \sqcup ((!U_n^{\text{op}} \times U_n) \sqcup A)$$

⁵ The only delicate point is the \star -autonomy of the bicategory, since it does not exist in the literature a notion of \star -autonomous bicategory. However it is possible to equip distributors with a dualizing pseudo-endofunctor, as shown for example in [12, 27].

38:10 Categorifying Non-Idempotent Intersection Types

Types:

$$a := x \in A \mid \langle a_1, \dots, a_k \rangle \Rightarrow a \mid \langle a_1, \dots, a_k \rangle$$

Morphisms in U :

$$\frac{f \in A[x, x']}{f \in U[x, x']} \quad \frac{\langle \sigma, \vec{f} \rangle : \vec{a}' \rightarrow \vec{a} \quad f : a \rightarrow a'}{\langle \sigma, \vec{f} \rangle \Rightarrow f : (\vec{a} \Rightarrow a) \rightarrow (\vec{a}' \Rightarrow a')}$$

$$\frac{\sigma \in S_n \quad f_1 : a_1 \rightarrow a'_{\sigma(1)} \quad \dots \quad f_n : a_n \rightarrow a'_{\sigma(n)}}{\langle \sigma, f_1, \dots, f_n \rangle : \langle a_1, \dots, a_n \rangle \rightarrow \langle a'_1, \dots, a'_n \rangle}$$

Derivation rules:

$$\frac{f : a' \rightarrow a}{x_1 : \langle \rangle, \dots, x_i : \langle a' \rangle, \dots, x_n : \langle \rangle \vdash x_i : a}$$

$$\frac{\Gamma \vdash S : \vec{a} \Rightarrow a \quad \Gamma' \vdash T : \vec{a} \quad \eta : \Delta \rightarrow \Gamma \otimes \Gamma'}{\Delta \vdash ST : a}$$

$$\frac{\Gamma_1 \vdash S : a_1 \quad \dots \quad \Gamma_k \vdash S : a_k \quad \eta : \Delta \rightarrow \bigotimes_{i=1}^k \Gamma_i}{\Delta \vdash S^! : \langle a_1, \dots, a_k \rangle}$$

$$\frac{\Delta, x : \vec{a} \vdash S : a}{\Delta \vdash \lambda x. S : \vec{a} \Rightarrow a}$$

■ **Figure 3** Non-idempotent intersection type system $\mathcal{R}_{\rightarrow}$ associated with the 0-cell U in Dist .

337 We define a family of inclusions $(\iota_n : U_n \hookrightarrow U_{n+1})_{n \in \mathbb{N}}$ in the canonical way:

$$338 \quad \iota_0 = \iota_A \quad \iota_{n+1} = !(\iota_n) \sqcup ((!(\iota_n)^{\text{op}} \times \iota_n) \sqcup 1_A)$$

339 Then we set $U_A = \varinjlim_{n \in \mathbb{N}} U_n$. From now on, the 0-cell U_A will be simply denoted by U , keeping
340 the parameter A implicit. We denote by $\xi_n : !U_n \sqcup (!U_n^{\text{op}} \times U_n) \hookrightarrow U_n$ the canonical inclusions.

341

342 ► **Lemma 5** (Inclusion). *There exists a canonical inclusion $\iota : !U \sqcup (!U^{\text{op}} \times U) \hookrightarrow U$.*

343 **Proof.** Since U is a filtered colimit, we have $!U \sqcup (!U^{\text{op}} \times U) \cong \varinjlim_{n \in \mathbb{N}} !U_n \sqcup (\varinjlim_{n \in \mathbb{N}} !U_n^{\text{op}} \times \varinjlim_{n \in \mathbb{N}} U_n)$,
344 and so we can explicitly define the inclusion functor as $\iota(a) = y_{j+1}(\xi_j(a))$ where $j = \min\{n \in \mathbb{N} \mid a \in U_n \sqcup (!U_n^{\text{op}} \times U_n)\}$ and $y_{j+1} : U_{j+1} \rightarrow U$ is the canonical injection of U_{j+1} . ◀

346 ► **Theorem 6** (Retraction). *We have that $!U \& (!U \multimap U) \triangleleft U$ in Dist .*

347 So, the 0-cell U is a (non-extensional) denotational model of the bang calculus. By seeing
348 the objects of A (resp. U) as the *atomic types* (resp. *types*) and setting $\vec{a} \Rightarrow a = \langle \vec{a}, a \rangle \in !U \times U$,
349 we give in Figure 3 a type-theoretic description of the 0-cell U . This *non-idempotent*
350 *intersection type system*, called $\mathcal{R}_{\rightarrow}$, is the generalization in Dist of the system \mathcal{R}_{\leq} in
351 Figure 2 associated with Polr . A morphism $f : a \rightarrow b$ in Figure 3 can be seen as a witness in
352 Dist of the subtyping relation between a and b , generalizing $a \leq_U b$ of Polr .

353 **Semantics of Bang Terms.** We now present the *semantics* (or denotation) of bang terms
354 as distributors in the bicategory Dist . We recall that $\iota : !U \& (!U^{\text{op}} \times U) \hookrightarrow U$. Let
355 $\Gamma = \langle \vec{b}_1, \dots, \vec{b}_n \rangle$, $\Delta = \langle \vec{b}'_1, \dots, \vec{b}'_n \rangle \in !U^n$. A morphism $\eta : \Gamma \rightarrow \Delta$ is a list of morphisms $\eta =$
356 $\langle \langle \sigma_1, \vec{f}_1 \rangle, \dots, \langle \sigma_n, \vec{f}_n \rangle \rangle : \Gamma \rightarrow \Delta$ where $\langle \sigma_i, \vec{f}_i \rangle : \vec{b}_i \rightarrow \vec{b}'_i$. We set $\Gamma \otimes \Delta = \langle \vec{b}_1 \oplus \vec{b}'_1, \dots, \vec{b}_n \oplus \vec{b}'_n \rangle$.
357 This tensor product inherits the relevant structure from \oplus . In particular, the symmetries
358 $\vec{\sigma} : \bigotimes_{i=1}^k \Gamma_i \rightarrow \bigotimes_{i=1}^k \Gamma_{\sigma(i)}$ are built from the σ^* construction presented in Section 2.

359 ► **Definition 7** (Semantics). *Let $S \in !\Lambda$ and $\text{fv}(S) \subseteq \vec{x} = \langle x_1, \dots, x_n \rangle$, with the x_i 's pairwise
360 distinct. The semantics $\llbracket S \rrbracket_{\vec{x}} : !U^{\otimes n} \multimap U$ of S with respect to \vec{x} is defined by induction on S :*

$$361 \quad \llbracket x_i \rrbracket_{\vec{x}}(\Delta, a) = !U^n[\Delta, \langle \langle \rangle, \dots, \langle a \rangle, \dots, \langle \rangle \rangle] \quad (\langle a \rangle \text{ is in the } i^{\text{th}} \text{ position in } \langle \langle \rangle, \dots, \langle a \rangle, \dots, \langle \rangle \rangle);$$

$$362 \quad \llbracket \lambda y. S \rrbracket_{\vec{x}}(\Delta, a) = \begin{cases} \llbracket S \rrbracket_{\vec{x} \oplus \langle y \rangle}(\Delta \oplus \langle \vec{a} \rangle, a') & \text{if } a = \iota(\langle \vec{a}, a' \rangle) \\ \emptyset & \text{otherwise.} \end{cases}, \text{ where } y \notin \vec{x};$$

$$363 \quad \llbracket ST \rrbracket_{\vec{x}}(\Delta, a) = \int^{\vec{a} \in !U} \int^{\Gamma_1, \Gamma_2 \in !U^n} \llbracket S \rrbracket_{\vec{x}}(\Gamma_1, \iota(\langle \vec{a}, a \rangle)) \times \llbracket T \rrbracket_{\vec{x}}(\Gamma_2, \iota(\vec{a})) \times (!U^n)(\Delta, \Gamma_1 \otimes \Gamma_2);$$

$$\begin{aligned}
[g: a \rightarrow b] \left(\frac{f: a' \rightarrow a}{x_1: \langle \rangle, \dots, x_i: \langle a' \rangle, \dots, x_n: \langle \rangle \vdash x_i: a} \right) &= \frac{g \circ f: a' \rightarrow b}{x_1: \langle \rangle, \dots, x_i: \langle a' \rangle, \dots, x_n: \langle \rangle \vdash x_i: b} \\
[(\sigma, \vec{g}) \Rightarrow g: (\vec{a} \Rightarrow a) \rightarrow (\vec{b} \Rightarrow b)] \left(\frac{\begin{array}{c} \vdots \pi \\ \Delta, x: \vec{a} \vdash S: a \\ \Delta \vdash \lambda x. S: \vec{a} \Rightarrow a \end{array}}{\Delta \vdash \lambda x. S: \vec{a} \Rightarrow a} \right) &= \frac{\begin{array}{c} \vdots [g]\pi\{1, \langle \sigma, \vec{g} \rangle\} \\ \Delta, x: \vec{b} \vdash S: b \\ \Delta \vdash \lambda x. S: \vec{b} \Rightarrow b \end{array}}{\Delta \vdash \lambda x. S: \vec{b} \Rightarrow b} \\
[g: a \rightarrow b] \left(\frac{\begin{array}{c} \vdots \pi_1 \quad \vdots \pi_2 \\ \Gamma_1 \vdash S: \vec{a} \Rightarrow a \quad \Gamma_2 \vdash T: \vec{a} \quad \eta: \Delta \rightarrow \Gamma_1 \otimes \Gamma_2 \\ \Delta \vdash ST: a \end{array}}{\Delta \vdash ST: a} \right) &= \frac{\begin{array}{c} \vdots [1 \Rightarrow g]\pi_1 \quad \vdots \pi_2 \\ \Gamma_1 \vdash S: \vec{a} \Rightarrow b \quad \Gamma_2 \vdash T: \vec{a} \quad \eta: \Delta \rightarrow \Gamma_1 \otimes \Gamma_2 \\ \Delta \vdash ST: b \end{array}}{\Delta \vdash ST: b} \\
[(\sigma, \vec{g}): \vec{a} \rightarrow \vec{b}] \left(\frac{\begin{array}{c} \vdots \pi_i \\ \Gamma_i \vdash S: a_i \\ \Delta \vdash S^! : \vec{a} = \langle a_1, \dots, a_k \rangle \end{array}}{\Delta \vdash S^! : \vec{a} = \langle a_1, \dots, a_k \rangle} \right) &= \frac{\begin{array}{c} \vdots \pi'_i \\ \Gamma_{\sigma^{-1}(i)} \vdash S: a_{\sigma^{-1}(i)} \\ \Delta \vdash S^! : \vec{b} = \langle b_1, \dots, b_k \rangle \end{array}}{\Delta \vdash S^! : \vec{b} = \langle b_1, \dots, b_k \rangle}
\end{aligned}$$

■ **Figure 4** Left action on derivations. In the last identity, on the right, $\pi'_i = [g_{\sigma^{-1}(i)}]\pi_{\sigma^{-1}(i)}$.

$$\begin{aligned}
364 \quad \dashv \llbracket S^! \rrbracket_{\vec{x}}(\Delta, a) &= \begin{cases} \int^{\Gamma_1, \dots, \Gamma_k \in !U^n} \prod_{i=1}^k \llbracket S \rrbracket_{\vec{x}}(\Gamma_i, a_i) \times (!U^n)(\Delta, \bigotimes_{i=1}^k \Gamma_i) & \text{if } a = \iota(\langle a_1, \dots, a_k \rangle) \\ \emptyset & \text{otherwise.} \end{cases}
\end{aligned}$$

365 Given $\langle \Delta, a \rangle \in !U^n \times U$ we call *points* the elements of $\llbracket S \rrbracket_{\vec{x}}(\Delta, a)$. From now on, when we
366 write $\llbracket S \rrbracket_{\vec{x}}$ we always assume that $\text{fv}(S) \subseteq \vec{x} = \langle x_1, \dots, x_n \rangle$ and the x_i 's are pairwise distinct.

367 The semantics of a term S is a functor $\llbracket S \rrbracket_{\vec{x}}: (!U^n)^{\text{op}} \times U \rightarrow \text{Set}$. As such, it must be
368 defined on the objects of the category $(!U^n)^{\text{op}} \times U$ (as done in Definition 7) and on the
369 morphisms of the category $(!U^n)^{\text{op}} \times U$. The action on morphisms (omitted in Definition 7) is
370 given by induction on S and, in the application and bang cases, also by the universal property
371 of the coend construction. The variable case is just the hom-functor. An explicit definition
372 of the application and bang cases can be given by considering coends as coequalizers [44].

373 **Non-idempotent Intersection Type Distributors.** We aim to define the *non-idempotent*
374 *intersection type distributor* $\mathbb{T}_U(S)_{\vec{x}}$ for any term S . Let π be a type derivation in system
375 $\mathcal{R}_{\rightarrow}$, as defined in Figure 3. The *left* and *right actions* of morphisms on π are defined in
376 Figures 4 and 5, respectively (by induction on π). Given $f: a \rightarrow a'$ and $\theta: \Delta' \rightarrow \Delta$, the left
377 and right actions may change the conclusion of a type derivation:

$$\begin{aligned}
378 \quad \text{left: } [f] \left(\frac{\begin{array}{c} \pi \\ \vdots \\ \Delta \vdash S: a \end{array}}{\Delta \vdash S: a} \right) &\rightsquigarrow \frac{[f]\pi}{\Delta \vdash S: a'} \quad \text{right: } \left(\frac{\begin{array}{c} \pi \\ \vdots \\ \Delta \vdash S: a \end{array}}{\Delta \vdash S: a} \right) \{\theta\} &\rightsquigarrow \frac{\pi\{\theta\}}{\Delta' \vdash S: a}
\end{aligned}$$

379 Notice the contravariance of the right action, and that $[f](\pi\{\theta\}) = ([f]\pi)\{\theta\}$.

380 We define \sim as the smallest congruence on type derivations generated by the rules in
381 Figure 6. We denote by $\tilde{\pi}$ the equivalence class of π modulo \sim . Note that $[f]\tilde{\pi}\{\theta\} = \widetilde{[f]\pi\{\theta\}}$.

382 ► **Example 8.** We give a couple of examples of the equivalence \sim between type derivations
383 in system $\mathcal{R}_{\rightarrow}$. The intuition is that \sim equalizes type derivations for the same term and with
384 the same conclusion, where the “same” permutations are performed at different moments.
385 Let $f: a' \rightarrow a$ be a morphism between types a' and a . One can think of them as, e.g.
386 $a = \langle *, \langle * \rangle \Rightarrow * \rangle$ and $a' = \langle \langle * \rangle \Rightarrow *, * \rangle$ with $f = \sigma \Rightarrow 1$ being the obvious permutation.

$$\begin{aligned}
 & \frac{f : a' \rightarrow a}{x_1 : \langle \rangle, \dots, x_i : \langle a' \rangle, \dots, x_n : \langle \rangle \vdash x_i : a} \{(g : b \rightarrow a')\} = \frac{f \circ g : b \rightarrow a}{x_1 : \langle \rangle, \dots, x_i : \langle b \rangle, \dots, x_n : \langle \rangle \vdash x_i : a} \\
 & \left(\frac{\frac{\pi}{\Delta, x : \vec{a} \vdash S : a}}{\Delta \vdash \lambda x. S : \vec{a} \Rightarrow a'} \right) \{\theta\} = \frac{\frac{\pi \{\theta \oplus \langle 1 \rangle\}}{\Delta', x : \vec{a} \vdash S : a}}{\Delta' \vdash \lambda x. S : \vec{a} \Rightarrow a} \\
 & \left(\frac{\frac{\frac{\pi_1}{\Gamma_1 \vdash S : \vec{a} \Rightarrow a} \quad \frac{\pi_2}{\Gamma_2 \vdash T : \vec{a}} \quad \eta : \Delta \rightarrow \Gamma_1 \otimes \Gamma_2}{\Delta \vdash ST : a}}{\Delta \vdash S^l : \langle a_1, \dots, a_k \rangle} \right) \{\theta\} = \frac{\frac{\frac{\pi_1}{\Gamma_1 \vdash S : \vec{a} \Rightarrow a} \quad \frac{\pi_2}{\Gamma_2 \vdash T : \vec{a}} \quad \eta \circ \theta : \Delta' \rightarrow \Gamma_1 \otimes \Gamma_2}{\Delta' \vdash ST : a}}{\Delta' \vdash S^l : \langle a_1, \dots, a_k \rangle} \\
 & \left(\frac{\left(\frac{\pi_i}{\Gamma_i \vdash S : a_i} \right)_{i=1}^k \quad \eta : \Delta \rightarrow \bigotimes_{i=1}^k \Gamma_i}{\Delta \vdash S^l : \langle a_1, \dots, a_k \rangle} \right) \{\theta\} = \left(\frac{\left(\frac{\pi_i}{\Gamma_i \vdash S : a_i} \right)_{i=1}^k \quad \eta \circ \theta : \Delta' \rightarrow \bigotimes_{i=1}^k \Gamma_i}{\Delta' \vdash S^l : \langle a_1, \dots, a_k \rangle} \right)
 \end{aligned}$$

■ **Figure 5** Right action on derivations, where $\theta : \Delta' \rightarrow \Delta$.

- 387 1. Let us type the term $xx^!$ with the following type derivation π (where $A = \langle \langle a \rangle \Rightarrow a, a \rangle$,
 388 $A' = \langle a', \langle a \rangle \Rightarrow a \rangle$ and $(12) \in S_2$ is the swap permutation on $\{1, 2\}$):

$$\begin{aligned}
 & \frac{1_{\langle a \rangle \Rightarrow a} : (\langle a \rangle \Rightarrow a) \rightarrow (\langle a \rangle \Rightarrow a) \quad \frac{1_a : a \rightarrow a}{x : \langle a \rangle \vdash x : a} \quad 1_{\langle a \rangle} : \langle a \rangle \rightarrow \langle a \rangle}{x : \langle \langle a \rangle \Rightarrow a \rangle \vdash x : \langle a \rangle \Rightarrow a} \quad \frac{x : \langle a \rangle \vdash x : a \quad 1_{\langle a \rangle} : \langle a \rangle \rightarrow \langle a \rangle}{x : \langle a \rangle \vdash x^! : \langle a \rangle} \quad \langle (12), f, 1_{\langle a \rangle \Rightarrow a} \rangle : A' \rightarrow A}{x : \langle a', \langle a \rangle \Rightarrow a \rangle \vdash xx^! : a}
 \end{aligned}$$

390 Now consider the following type derivation π' (with $A'' = \langle \langle a \rangle \Rightarrow a, a' \rangle$)

$$\begin{aligned}
 & \frac{1_{\langle a \rangle \Rightarrow a} : (\langle a \rangle \Rightarrow a) \rightarrow (\langle a \rangle \Rightarrow a) \quad \frac{f : a' \rightarrow a}{x : \langle a' \rangle \vdash x : a} \quad 1_{\langle a' \rangle} : \langle a' \rangle \rightarrow \langle a' \rangle}{x : \langle \langle a \rangle \Rightarrow a \rangle \vdash x : \langle a \rangle \Rightarrow a} \quad \frac{x : \langle a' \rangle \vdash x : a \quad 1_{\langle a' \rangle} : \langle a' \rangle \rightarrow \langle a' \rangle}{x : \langle a' \rangle \vdash x^! : \langle a \rangle} \quad \langle (12), 1_{a'}, 1_{\langle a \rangle \Rightarrow a} \rangle : A' \rightarrow A''}{x : \langle a', \langle a \rangle \Rightarrow a \rangle \vdash xx^! : a}
 \end{aligned}$$

392 Compared to π , π' brings forward the morphism f . By the second rule in Figure 6, $\pi \sim \pi'$.

- 393 2. Let us type the term $(\lambda x.x)z^!$ (we omit the index on the identity morphisms 1):

$$\begin{aligned}
 \pi = & \frac{\frac{f : a' \rightarrow a}{x : \langle a' \rangle \vdash x : a} \quad \frac{1 : a' \rightarrow a'}{z : \langle a' \rangle \vdash z : a'} \quad 1 : \langle a' \rangle \rightarrow \langle a' \rangle}{\vdash \lambda x.x : \langle a' \rangle \Rightarrow a} \quad \frac{z : \langle a' \rangle \vdash z^! : \langle a' \rangle}{z : \langle a' \rangle \vdash (\lambda x.x)z^! : a} \quad 1 : \langle a' \rangle \rightarrow \langle a' \rangle
 \end{aligned}$$

395 Now consider the following derivation (note the different position of f with respect to π)

$$\begin{aligned}
 \pi' = & \frac{\frac{1 : a \rightarrow a}{x : \langle a \rangle \vdash x : a} \quad \frac{f : a' \rightarrow a}{z : \langle a' \rangle \vdash z : a} \quad 1 : \langle a' \rangle \rightarrow \langle a' \rangle}{\vdash \lambda x.x : \langle a \rangle \Rightarrow a} \quad \frac{z : \langle a' \rangle \vdash z^! : \langle a \rangle}{z : \langle a' \rangle \vdash (\lambda x.x)z^! : a} \quad 1 : \langle a' \rangle \rightarrow \langle a' \rangle
 \end{aligned}$$

397 According to the first rule in Figure 6, $\pi \sim \pi'$.

398 Let S be a term and $\text{fv}(S) \subseteq \vec{x} = \{x_1, \dots, x_n\}$ with the x_i 's pairwise distinct. With any
 399 $\langle \Delta, a \rangle \in \text{ob}((!U^n)^{\text{op}} \times U)$, the distributor $\mathbb{T}_U(S)_{\vec{x}} : !U^n \rightarrow U$ associates the set of (equivalence
 400 classes of) type derivations for S with conclusion $\Delta \vdash S : a$. Formally, $\mathbb{T}_U(S)_{\vec{x}}$ is defined by:

$$\begin{array}{c}
\begin{array}{c} \pi_1 \\ \vdots \\ \Gamma_1 \vdash S : \vec{b} \Rightarrow a \end{array} \quad \begin{array}{c} [(\sigma, \vec{f})]\pi_2 \\ \vdots \\ \Gamma_2 \vdash T : \vec{b} \end{array} \quad \eta : \Delta \rightarrow \Gamma_1 \otimes \Gamma_2 \\
\hline
\Delta \vdash ST : a \\
\begin{array}{c} \pi_1\{\theta_1\} \\ \vdots \\ \Gamma_1 \vdash S : \vec{a} \Rightarrow a \end{array} \quad \begin{array}{c} \pi_2\{\theta_2\} \\ \vdots \\ \Gamma_2 \vdash T : \vec{a} \end{array} \quad \eta : \Delta \rightarrow \Gamma_1 \otimes \Gamma_2 \\
\hline
\Delta \vdash ST : a \\
\left(\begin{array}{c} \pi_i\{\theta_i\} \\ \vdots \\ \Gamma_i \vdash S : a_i \end{array} \right)_{i=1}^k \quad \eta : \Delta \rightarrow \bigotimes_{i=1}^k \Gamma_i \\
\hline
\Delta \vdash S^! : \langle a_1, \dots, a_k \rangle
\end{array}
\sim
\begin{array}{c}
\begin{array}{c} [(\sigma, \vec{f}) \Rightarrow 1]\pi_1 \\ \vdots \\ \Gamma_1 \vdash S : \vec{a} \Rightarrow a \end{array} \quad \begin{array}{c} \pi_2 \\ \vdots \\ \Gamma_2 \vdash T : \vec{a} \end{array} \quad \eta : \Delta \rightarrow \Gamma_1 \otimes \Gamma_2 \\
\hline
\Delta \vdash ST : a \\
\begin{array}{c} \pi_1 \\ \vdots \\ \Gamma'_1 \vdash S : \vec{a} \Rightarrow a \end{array} \quad \begin{array}{c} \pi_2 \\ \vdots \\ \Gamma'_2 \vdash T : \vec{a} \end{array} \quad \theta \circ \eta : \Delta \rightarrow \Gamma'_1 \otimes \Gamma'_2 \\
\hline
\Delta \vdash ST : a \\
\left(\begin{array}{c} \pi_i \\ \vdots \\ \Gamma'_i \vdash S : a_i \end{array} \right)_{i=1}^k \quad \bigotimes_{i=1}^k \theta_i \circ \eta : \Delta \rightarrow \bigotimes_{i=1}^k \Gamma'_i \\
\hline
\Delta \vdash S^! : \langle a_1, \dots, a_k \rangle
\end{array}
\end{array}$$

■ **Figure 6** Congruence on type derivations, where $\langle \sigma, \vec{f} \rangle : \vec{a} \rightarrow \vec{b}$ and $\theta = \theta_1 \otimes \theta_2$ with $\theta_i : \Gamma_i \rightarrow \Gamma'_i$.

- 401 1. for $\langle \Delta, a \rangle \in \text{ob}((!U^n)^{\text{op}} \times U)$, $\mathbb{T}_U(S)_{\vec{x}}(\Delta, a) = \left\{ \begin{array}{c} \tilde{\pi} \\ \vdots \\ \Delta \vdash S : a \end{array} \mid \pi \text{ is a type derivation for } S \right\}$;
- 402 2. for $f : a \rightarrow a'$ and $\eta : \Delta' \rightarrow \Delta$, $\mathbb{T}_U(S)_{\vec{x}}(\eta, f) : \mathbb{T}_U(S)_{\vec{x}}(\Delta, a) \rightarrow \mathbb{T}_U(S)_{\vec{x}}(\Delta', a')$ such that
- 403 $\mathbb{T}_U(S)_{\vec{x}}(\eta, f)(\tilde{\pi}) = [f]\pi\{\eta\} \in \mathbb{T}_U(S)_{\vec{x}}(\Delta', a')$ for any $\tilde{\pi} \in \mathbb{T}_U(S)_{\vec{x}}(\Delta, a)$.

404 ► **Lemma 9** (Functoriality). *For any $S \in !\Lambda$, $\mathbb{T}_U(S)_{\vec{x}}$ is a functor from $(!U^n)^{\text{op}} \times U$ to Set .*

405 The following theorem states that the distributor semantics induced by our category
406 of types U can be seen in a completely type-theoretic way. The semantics $\llbracket S \rrbracket_{\vec{x}}(\Delta, a)$ of a
407 term S is equal to the set of (equivalence classes of) type derivations whose conclusion is the
408 sequent $\Delta \vdash S : a$. For this reason we have a bicategorical *proof relevant* semantics. This is a
409 major improvement over relational semantics, where the elements of the denotation of a term
410 are only witnesses of typability. Said differently, the relational semantics of S is just the set
411 of conclusions of the type derivations for S , while the distributor semantics of S provides,
412 for any conclusion, the set of type derivations for S with such a conclusion.

413 ► **Theorem 10** (Proof-relevance). *Let $S \in !\Lambda$. There is an isomorphism of functors*

414
$$\psi : \llbracket S \rrbracket_{\vec{x}} \cong \mathbb{T}_U(S)_{\vec{x}} \quad \text{which is natural in } \langle \Delta, a \rangle \in \text{ob}((!U^n)^{\text{op}} \times U).$$

415 **Proof.** By induction on the structure of S . The core of the proof is the remark that we can
416 write the equivalence relation induced by the coend in the application and box cases with
417 the rules in Figure 6. ◀

418 For any type derivation π in system $\mathcal{R}_{\rightarrow}$ we define its *size* $s(\pi)$ in Figure 7 (by induction on
419 π). It counts the number of rules for application in π . Note that if $\pi \sim \pi'$ then $s(\pi) = s(\pi')$.
420 We also have that size is invariant under morphisms action: $s([f]\pi) = s(\pi\{\eta\}) = s(\pi)$.

421 Let $\psi : \llbracket S \rrbracket_{\vec{x}} \cong \mathbb{T}_U(S)_{\vec{x}}$ be the natural isomorphism of Theorem 10. For $\alpha \in \llbracket S \rrbracket_{\vec{x}}(\Delta, a)$
422 we set $s(\alpha) = s(\psi_{\Delta, a}(\alpha))$, *i.e.* the size of a point α is the size of its derivation $\psi_{\Delta, a}(\alpha)$.

423 **Substitution and Reduction.** We prove both subject reduction and expansion for non-
424 idempotent intersection type distributors. We enrich this result with a quantitative flavor,
425 accounting for how the size of points is affected by a reduction step. In this way, we can give
426 a combinatorial proof for the characterization of terms that are normalizable at depth 0.

38:14 Categorifying Non-Idempotent Intersection Types

$$\begin{aligned}
s\left(\frac{f : a' \rightarrow a}{x_1 : \langle \rangle, \dots, x_i : \langle a' \rangle, \dots, x_n : \langle \rangle \vdash x_i : a}\right) &= 0 & s\left(\frac{\left(\frac{\pi_i}{\Gamma_i \vdash S : a_i}\right)_{i=1}^k}{\Delta \vdash !S : \langle a_1, \dots, a_k \rangle}\right) &= \sum_{i \in [k]} s(\pi_i) \\
s\left(\frac{\pi'}{\Delta, x : \vec{a} \vdash S : a}\right) &= s(\pi') & s\left(\frac{\frac{\pi_1}{\Gamma_1 \vdash S : \vec{a} \Rightarrow a} \quad \frac{\pi_2}{\Gamma_2 \vdash T : \vec{a}} \quad \theta : \Delta \rightarrow \Gamma_1 \otimes \Gamma_2}{\Delta \vdash ST : a}\right) &= s(\pi_1) + s(\pi_2) + 1
\end{aligned}$$

■ **Figure 7** Size of type derivations in system $\mathcal{R}_{\rightarrow}$.

427 The key ingredient is the substitution lemma below. We set:

$$428 \quad Sub_{S,x,T}(\Delta, a) = \int^{\vec{a} \in !U} \int^{\Gamma_0, \Gamma_1 \in !U^n} \llbracket S \rrbracket_{\vec{x} \oplus \langle x \rangle}(\Gamma_0 \oplus \langle \vec{a} \rangle, a) \times \llbracket T \rrbracket_{\vec{x}}(\Gamma_1, \vec{a}) \times !U^n(\Delta, \Gamma_0 \otimes \Gamma_1).$$

429 ► **Lemma 11** (Substitution). *Let S and T be terms. There is an isomorphism of functors*

$$430 \quad \varphi : Sub_{S,x,T} \cong \llbracket S\{T/x\} \rrbracket_{\vec{x}}$$

431 *natural in $\langle \Delta, a \rangle \in \text{ob}(!U^n)^{\text{op}} \times U$ and such that $s(\varphi_{\Delta,a}(\langle \widetilde{\alpha_1}, \alpha_2, \eta \rangle)) = s(\alpha_1) + s(\alpha_2)$.*

432 **Proof.** By induction on the structure of S , via lengthy coend manipulations. The proof of
433 the application and list cases strongly relies on the fact that the tensor product of $!U$
434 is symmetric. The proof of the preservation of sizes relies on the fact that size is invariant
435 under morphism actions and equivalence. Details are in Appendix A. ◀

436 ► **Theorem 12** (Subject reduction and expansion). *Let S, T be two terms.*

- 437 1. *If $S \rightarrow_b T$ then there is a natural isomorphism $\llbracket S \rrbracket_{\vec{x}}(\Delta, a) \cong \llbracket T \rrbracket_{\vec{x}}(\Delta, a)$.*
- 438 2. *If $S \rightarrow_{b_g} T$ then $\llbracket S \rrbracket_{\vec{x}}(\Delta, a) \cong \llbracket T \rrbracket_{\vec{x}}(\Delta, a)$ via a natural isomorphism $\varphi_{\Delta,a}$ such that*
439 *$s(\varphi_{\Delta,a}(\alpha)) = s(\alpha) - 1$ for any $\alpha \in \llbracket S \rrbracket_{\vec{x}}(\Delta, a)$.*
- 440 3. *If $S \rightarrow_{\sigma} T$ then $\llbracket S \rrbracket_{\vec{x}}(\Delta, a) \cong \llbracket T \rrbracket_{\vec{x}}(\Delta, a)$ via a natural isomorphism $\varphi_{\Delta,a}$ such that*
441 *$s(\varphi_{\Delta,a}(\alpha)) = s(\alpha)$ for any $\alpha \in \llbracket S \rrbracket_{\vec{x}}(\Delta, a)$.*

442 **Proof.** We prove the base case of Item 2, which follows from the substitution lemma
443 (Lemma 11). Let $S = (\lambda x.S_1)S_2 \mapsto_b S_1\{S_2/x\} = T$. By definition, we have

$$444 \quad \llbracket S \rrbracket_{\vec{x}}(\Delta, a) = \int^{\vec{a} \in !U} \int^{\Gamma_1, \Gamma_2 \in !U^n} \llbracket \lambda x.S_1 \rrbracket_{\vec{x}}(\Gamma_1, \iota(\langle \vec{a}, a \rangle)) \times \llbracket S_2 \rrbracket_{\vec{x}}(\Gamma_2, \vec{a}) \times !U^n(\Delta, \Gamma_1 \otimes \Gamma_2).$$

445 By definition of an abstraction's denotation we have $\llbracket \lambda x.S_1 \rrbracket_{\vec{x}}(\Gamma_1, \iota(\langle \vec{a}, a \rangle)) = \llbracket S_1 \rrbracket_{\vec{x} \oplus \langle x \rangle}(\Gamma_1 \oplus$
446 $\langle \vec{a} \rangle, a)$. Then, $\llbracket S \rrbracket_{\vec{x}}(\Delta, a) = Sub_{S_1,x,S_2}(\Delta, a)$. By Lemma 11, $\varphi_{\Delta,a} : \llbracket (\lambda x.S_1)S_2 \rrbracket_{\vec{x}}(\Delta, a) \cong$
447 $\llbracket S_1\{S_2/x\} \rrbracket_{\vec{x}}(\Delta, a)$. Again by Lemma 11, $s(\varphi_{\Delta,a}(\beta)) = s(\alpha_1) + s(\alpha_2)$ for $\beta = \langle \widetilde{\alpha_1}, \alpha_2, \eta \rangle \in$
448 $\llbracket S \rrbracket_{\vec{x}}(\Delta, a)$. By definition, we have that $s(\beta) = s(\alpha_1) + s(\alpha_2) + 1$. So, we can conclude.

449 For Item 3, a step \rightarrow_{σ} just requires to rearrange the rule order in a type derivation. ◀

450 Roughly, Theorem 12.2 states that if $S \rightarrow_{b_g} T$ then for every type derivation for S there is
451 a type derivation for T , with the same conclusion, whose size decreases by 1. In Theorem 12.1
452 such a quantitative account does not hold. Indeed, consider $((\lambda x.x)y^!) \mapsto_b y^!$: each of the
453 two terms can be typed with a derivation of size 0 (take the rule for boxes with 0 premises).

454 ► **Example 13.** We provide a simple example of reduction of type derivations to ease the
 455 understanding of the congruence's role in establishing the natural isomorphisms. Consider
 456 $S = (\lambda x.x)y^!$. We type it with the following type derivations:

$$457 \quad \pi_1 = \frac{\frac{h \circ f : a \rightarrow b}{x : \langle a \rangle \vdash x : b} \quad \frac{g : c \rightarrow a}{y : \langle c \rangle \vdash y : a} \quad 1}{\vdash \lambda x.x : \langle a \rangle \Rightarrow b} \quad \frac{y : \langle c \rangle \vdash y^! : \langle a \rangle}{1}}{y : \langle c \rangle \vdash (\lambda x.x)y^! : b} \quad \pi_2 = \frac{\frac{h \circ f' : a' \rightarrow b}{x : \langle a \rangle \vdash x : b} \quad \frac{g' : c \rightarrow a'}{y : \langle c \rangle \vdash y : a'} \quad 1}{\vdash \lambda x.x : \langle d \rangle \Rightarrow b} \quad \frac{y : \langle c \rangle \vdash y^! : \langle a' \rangle}{1}}{y : \langle c \rangle \vdash (\lambda x.x)y^! : b}$$

458 Suppose that $f \circ g = f' \circ g'$ and $h : b \rightarrow b$, $f : a \rightarrow b$, $f' : a' \rightarrow b$. We have that $\pi_1 \sim \pi_2$.
 459 Indeed, by the first rule of Figure 6:

$$460 \quad \pi_1 \sim \frac{\frac{h : b \rightarrow b}{x : \langle b \rangle \vdash x : b} \quad \frac{f \circ g : c \rightarrow b}{y : \langle c \rangle \vdash y : b} \quad 1}{\vdash \lambda x.x : \langle b \rangle \Rightarrow b} \quad \frac{y : \langle c \rangle \vdash y^! : \langle b \rangle}{1}}{y : \langle c \rangle \vdash (\lambda x.x)y^! : b} \quad \pi_2 \sim \frac{\frac{h : b \rightarrow b}{x : \langle b \rangle \vdash x : b} \quad \frac{f' \circ g' : c \rightarrow b}{y : \langle c \rangle \vdash y : b} \quad 1}{\vdash \lambda x.x : \langle b \rangle \Rightarrow b} \quad \frac{y : \langle c \rangle \vdash y^! : \langle b \rangle}{1}}{y : \langle c \rangle \vdash (\lambda x.x)y^! : b}$$

461 and, by the hypothesis $f \circ g = f' \circ g'$, we conclude that $\pi_1 \sim \pi_2$ by transitivity. In particular,
 462 this means that the quotient identify all couple of morphisms leading to the same composition.

463 Now, we have that $S \rightarrow_{\text{b}_g} y$. Consider the following type derivation of y :

$$464 \quad \pi_3 = \frac{h \circ (f \circ g) : c \rightarrow b}{y : \langle c \rangle \vdash y : b} \quad (\text{note that } s(\pi_1) = s(\pi_2) = 1 \text{ and } s(\pi_3) = 0).$$

465 By an easy inspection of the definitions we have that for $\varphi_{\langle c \rangle, b} : \llbracket S \rrbracket_{\langle y \rangle}(\langle c \rangle, b) \cong \llbracket y \rrbracket_{\langle y \rangle}(\langle c \rangle, b)$,
 466 $\varphi_{\langle c \rangle, b}(\pi_1) = \pi_3$, where we keep implicit the isomorphism given by Theorem 10. There is
 467 then a nice correspondence between *substitution* on the term side and *composition* on the
 468 morphism side, that validates the basic intuition of categorical semantics⁶.

469 We prove that non-idempotent intersection type distributors characterize normalization
 470 at depth 0, when normal forms are clash-free at depth 0. First, we characterize syntactically
 471 the normal forms for $\rightarrow_{\text{b}_{\sigma_g}}$ that are clash-free at depth 0. Consider the subsets $!\Lambda_d$, $!\Lambda_n$, $!\Lambda_\ell$
 472 (whose elements are denoted by D , N , L , respectively) of $!\Lambda$:

$$473 \quad (!\Lambda_d) \quad D ::= x \mid DS^! \mid DD' \quad (!\Lambda_n) \quad N ::= S^! \mid D \mid (\lambda x.N)D \quad (!\Lambda_\ell) \quad L ::= N \mid \lambda x.L$$

475 All terms in $!\Lambda_d$ are not closed (they have a free ‘‘head variable’’) and are neither a box nor
 476 a β -like redex nor an abstraction. Clearly, $!\Lambda_d \subsetneq !\Lambda_n$ and $!\Lambda_n \subsetneq !\Lambda_\ell$ with $!\Lambda_d \cap !\Lambda_n = \emptyset$.

- 477 ► **Proposition 14** (Syntactic characterization of clash-free at depth 0 normal forms for $\rightarrow_{\text{b}_{\sigma_g}}$).
- 478 1. A term S is normal for $\rightarrow_{\text{b}_{\sigma_g}}$, clash-free at depth 0 and is neither a box nor a β -like redex
 479 (i.e. nor of the form $(\lambda x.S)T$) nor an abstraction iff $S \in !\Lambda_d$.
 - 480 2. A term S is normal for $\rightarrow_{\text{b}_{\sigma_g}}$, clash-free at depth 0 and is not an abstraction iff $S \in !\Lambda_n$.
 - 481 3. A term S is normal for $\rightarrow_{\text{b}_{\sigma_g}}$ and clash-free at depth 0 iff $S \in !\Lambda_\ell$.

482 ► **Lemma 15** (Semantics vs. clash-free at depth 0). Let S be a term.

- 483 1. If $\llbracket S \rrbracket_{\vec{x}} \neq \emptyset_{!U^n, U}$ then S is clash-free at depth 0.
- 484 2. If S is normal for $\rightarrow_{\text{b}_{\sigma_g}}$ and clash-free at depth 0, then $\llbracket S \rrbracket_{\vec{x}} \neq \emptyset_{!U^n, U}$.

⁶ The natural isomorphism $\varphi_{\langle c \rangle, b} : \llbracket S \rrbracket_{\langle y \rangle}(\langle c \rangle, b) \cong \llbracket y \rrbracket_{\langle y \rangle}(\langle c \rangle, b)$ is a particular instance of Yoneda's lemma for coends (see Lemma 20 in Appendix A), also known as the density formula for coends [40].

485 **Proof.** 1. By induction on $S \in !\Lambda$.

486 2. According to Proposition 14, we can proceed by induction on $S \in !\Lambda_\ell$. ◀

487 ▶ **Theorem 16** (Normalization at depth 0). *Let S be a term. The following are equivalent:*

488 1. S is typable in system $\mathcal{R}_{\rightarrow}$;

489 2. $\llbracket S \rrbracket_{\vec{x}} \neq \emptyset_{!U^n, U}$;

490 3. S is strongly $\rightarrow_{\text{b}\sigma_{\mathfrak{g}}}$ -normalizable with a normal form for $\rightarrow_{\text{b}\sigma_{\mathfrak{g}}}$ that is clash-free at depth 0;

491 4. S is weakly $\rightarrow_{\text{b}\sigma_{\mathfrak{g}}}$ -normalizable with a normal form for $\rightarrow_{\text{b}\sigma_{\mathfrak{g}}}$ that is clash-free at depth 0;

492 5. $S \rightarrow_{\text{b}\sigma}^* T$ for some term T that is normal for $\rightarrow_{\text{b}\sigma_{\mathfrak{g}}}$ and clash free at depth 0.

493 **Proof.** The equivalence (1) \Leftrightarrow (2) is given by Theorem 10. The implication (5) \Rightarrow (2) follows
494 from Lemma 15.2 and Theorem 12. The implication (4) \Rightarrow (5) holds because $\rightarrow_{\text{b}\sigma_{\mathfrak{g}}} \subseteq \rightarrow_{\text{b}\sigma}$.
495 The implication (3) \Rightarrow (4) is trivial.

496 For the implication (2) \Rightarrow (3), as $\llbracket S \rrbracket_{\vec{x}} \neq \emptyset_{!U^n, U}$, there is a point $\alpha \in \llbracket S \rrbracket_{\vec{x}}(\Delta, a)$ for some
497 $\langle \Delta, a \rangle \in \text{ob}(!U^n \times U)$. Let k_S be the sum of the lengths of all $\rightarrow_{\sigma_{\mathfrak{g}}}$ -reduction sequences from
498 S to a normal form for $\rightarrow_{\sigma_{\mathfrak{g}}}$ (such a k_S exists because $\rightarrow_{\sigma_{\mathfrak{g}}}$ is strongly normalizing [24]). We
499 prove (3) by induction on $(s(\alpha), k_S)$ ordered lexicographically. If S is normal for $\rightarrow_{\text{b}\sigma_{\mathfrak{g}}}$, we
500 are done by Lemma 15.1, as $\alpha \in \llbracket S \rrbracket_{\vec{x}}(\Delta, a)$ implies $\llbracket S \rrbracket_{\vec{x}} \neq \emptyset_{!U^n, U}$. Suppose $S \rightarrow_{\text{b}\sigma_{\mathfrak{g}}} S'$.

501 1. If $S \rightarrow_{\sigma_{\mathfrak{g}}} S'$, let $\varphi: \llbracket S \rrbracket_{\vec{x}} \cong \llbracket S' \rrbracket_{\vec{x}}$ be the natural isomorphism of Theorem 12.3. Thus,
502 $\varphi_{\Delta, a}(\alpha) \in \llbracket S' \rrbracket_{\vec{x}}(\Delta, a)$ and $s(\alpha) = s(\varphi_{\Delta, a}(\alpha))$ but $k_{S'} = k_S - 1$.

503 2. If $S \rightarrow_{\text{b}\sigma_{\mathfrak{g}}} S'$, let $\varphi: \llbracket S \rrbracket_{\vec{x}} \cong \llbracket S' \rrbracket_{\vec{x}}$ be the natural isomorphism of Theorem 12.2. Thus,
504 $\varphi_{\Delta, a}(\alpha) \in \llbracket S' \rrbracket_{\vec{x}}(\Delta, a)$ and $s(\varphi_{\Delta, a}(\alpha)) = s(\alpha) - 1$.

505 In both cases, by *i.h.*, (3) holds for S' . Therefore, (3) holds for S . ◀

506 5 Conclusions

507 In this paper, we recalled some well-known and linear-logic based categorical semantics
508 with an intersection type presentation. We showed that they can be generalized in the
509 bicategory of distributors. We defined non-idempotent intersection type distributors in
510 the bang calculus and provided a syntactic presentation of them as a non-idempotent
511 intersection type system generalizing De Carvalho's system \mathcal{R} [16, 17]. We proved that non-
512 idempotent intersection type distributors determine a proof-relevant denotational semantics,
513 and characterize normalization at depth 0 in the bang calculus via a combinatorial proof.

514 **Perspectives.** Reconciling the different techniques used in [43, 41, 47] to categorify—non-
515 idempotent or possibly idempotent—intersection types is the first and natural open question.
516 The (non-trivial) answer should rely on a *subtyping-aware polyadic calculus* to be defined.

517 Another line of research is the study of the extensional collapse [22] in the bicategorical
518 setting of distributors, which should shed new light on the link between non-idempotent and
519 idempotent intersection types. Relating the methods of [29, 43] should be a first step.

520 A relevant question immediately arises also for what concerns typed call-by-push-value
521 [23, 39]. The extension of our work to that framework is tricky, since the semantics of types
522 adds technical machinery. Moreover, we believe that difficulties similar to the ones found in
523 [13] in order to define the Taylor expansion could arise also in our perspective.

524 Other interesting perspectives are the investigation of the relationship between our
525 categorified rigid framework and rigid intersection types [49], and an extension of our
526 approach to probabilistic computation. This extension is far from trivial, but the results
527 of Tsukada, Asada and Ong [48] are encouraging, and the study of probabilistic Taylor
528 expansion [38] and probabilistic intersection types [7] might be a starting point.

529 — References —

- 530 1 Fabio Alessi, Franco Barbanera, and Mariangiola Dezani-Ciancaglini. Intersection types and
531 lambda models. *Theoretical Computer Science*, 355(2):108 – 126, 2006. Logic, Language,
532 Information and Computation. doi:10.1016/j.tcs.2006.01.004.
- 533 2 Hendrik Pieter Barendregt. *The lambda calculus - its syntax and semantics*, volume 103 of
534 *Studies in logic and the foundations of mathematics*. North-Holland, 1984.
- 535 3 Jean Bénabou. Introduction to bicategories. In *Reports of the Midwest Category Seminar*,
536 pages 1–77, Berlin, Heidelberg, 1967. Springer Berlin Heidelberg.
- 537 4 Alexis Bernadet and Stéphane Jean Lengrand. Non-idempotent intersection types and strong
538 normalisation. *Logical Methods in Computer Science*, Volume 9, Issue 4, 2013. doi:10.2168/
539 LMCS-9(4:3)2013.
- 540 5 Robert Blackwell, Gregory Maxwell Kelly, and John Power. Two-dimensional monad theory.
541 *Journal of Pure and Applied Algebra*, 59(1):1 – 41, 1989. doi:https://doi.org/10.1016/
542 0022-4049(89)90160-6.
- 543 6 Francis Borceux. *Handbook of Categorical Algebra*, volume 1 of *Encyclopedia of Mathematics*
544 *and its Applications*. Cambridge University Press, 1994. doi:10.1017/CB09780511525858.
- 545 7 Flavien Breuvar and Ugo Dal Lago. On Intersection Types and Probabilistic Lambda Calculi.
546 In *Proceedings of the 20th International Symposium on Principles and Practice of Declarative*
547 *Programming, PPDP 2018, Frankfurt am Main, Germany, September 03-05, 2018*, pages
548 8:1–8:13, 2018. doi:10.1145/3236950.3236968.
- 549 8 Antonio Bucciarelli, Delia Kesner, Alejandro Ríos, and Andrés Viso. The bang calculus
550 revisited. In *Functional and Logic Programming - 15th International Symposium, FLOPS*
551 *2020*, volume 12073 of *Lecture Notes in Computer Science*, pages 13–32. Springer, 2020.
552 doi:10.1007/978-3-030-59025-3_2.
- 553 9 Antonio Bucciarelli, Delia Kesner, and Daniel Ventura. Non-idempotent intersection types for
554 the lambda-calculus. *Logic Journal of the IGPL*, 25(4):431–464, 2017. doi:10.1093/jigpal/
555 jzx018.
- 556 10 Jean Bénabou. Distributors at work. Lecture notes of a course given at TU Darmstadt, 2000.
557 URL: <http://www2.mathematik.tu-darmstadt.de/~streicher/FIBR/DiWo.pdf>.
- 558 11 Alberto Carraro and Giulio Guerrieri. A Semantical and Operational Account of Call-by-Value
559 Solvability. In *Foundations of Software Science and Computation Structures, FOSSACS 2014*,
560 volume 8412 of *Lecture Notes in Computer Science*, pages 103–118, Berlin, Heidelberg, 2014.
561 Springer. doi:10.1007/978-3-642-54830-7_7.
- 562 12 Gian Luca Cattani and Glynn Winskel. Profunctors, open maps and bisimulation. *Mathematical*
563 *Structures in Computer Science*, 15(3):553–614, 2005. doi:10.1017/S0960129505004718.
- 564 13 Jules Chouquet and Christine Tasson. Taylor expansion for Call-by-Push-Value. In *28th*
565 *EACSL Annual Conference on Computer Science Logic, CSL 2020*, volume 152 of *LIPICs*,
566 pages 16:1–16:16. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020. doi:10.4230/
567 LIPICs.CSL.2020.16.
- 568 14 Mario Coppo and Mariangiola Dezani-Ciancaglini. A new type-assignment for lambda terms.
569 *Arch. Math. Log.*, 19(1):139–156, 1978. doi:10.1007/BF02011875.
- 570 15 Mario Coppo and Mariangiola Dezani-Ciancaglini. An extension of the basic functionality
571 theory for the λ -calculus. *Notre Dame Journal of Formal Logic*, 21(4):685–693, 1980. doi:
572 10.1305/ndjfl/1093883253.
- 573 16 Daniel de Carvalho. *Semantique de la logique lineaire et temps de calcul*. PhD thesis, Aix-
574 Marseille Université, 2007.
- 575 17 Daniel de Carvalho. Execution time of λ -terms via denotational semantics and intersection
576 types. *Math. Struct. Comput. Sci.*, 28(7):1169–1203, 2018. doi:10.1017/S0960129516000396.
- 577 18 Daniel de Carvalho. Taylor expansion in linear logic is invertible. *Log. Methods Comput. Sci.*,
578 14(4), 2018. doi:10.23638/LMCS-14(4:21)2018.

- 579 19 Daniel de Carvalho, Michele Pagani, and Lorenzo Tortora de Falco. A semantic measure of
580 the execution time in linear logic. *Theoretical Computer Science*, 412(20):1884 – 1902, 2011.
581 doi:10.1016/j.tcs.2010.12.017.
- 582 20 Daniel de Carvalho and Lorenzo Tortora de Falco. A semantic account of strong normalization
583 in linear logic. *Inf. Comput.*, 248:104–129, 2016. doi:10.1016/j.ic.2015.12.010.
- 584 21 Thomas Ehrhard. Collapsing non-idempotent intersection types. In *Computer Science Logic*
585 *(CSL'12) - 26th International Workshop/21st Annual Conference of the EACSL, CSL 2012*,
586 volume 16 of *LIPICs*, pages 259–273. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2012.
587 doi:10.4230/LIPICs.CSL.2012.259.
- 588 22 Thomas Ehrhard. The Scott model of linear logic is the extensional collapse of its relational
589 model. *Theoretical Computer Science*, 424:20 – 45, 2012. doi:https://doi.org/10.1016/j.
590 tcs.2011.11.027.
- 591 23 Thomas Ehrhard. Call-by-push-value from a linear logic point of view. In *Programming*
592 *Languages and Systems - 25th European Symposium on Programming, ESOP 2016*, volume
593 9632 of *Lecture Notes in Computer Science*, pages 202–228. Springer, 2016. doi:10.1007/
594 978-3-662-49498-1_9.
- 595 24 Thomas Ehrhard and Giulio Guerrieri. The bang calculus: An untyped lambda-calculus gener-
596 alizing call-by-name and call-by-value. In *Proceedings of the 18th International Symposium on*
597 *Principles and Practice of Declarative Programming, PPDP 2016*, pages 174–187. Association
598 for Computing Machinery, 2016. doi:10.1145/2967973.2968608.
- 599 25 Thomas Ehrhard and Laurent Regnier. Böhm trees, Krivine’s machine and the Taylor expansion
600 of lambda-terms. In *Logical Approaches to Computational Barriers, Second Conference on*
601 *Computability in Europe, CiE 2006*, volume 3988 of *Lecture Notes in Computer Science*, pages
602 186–197. Springer, 2006. doi:10.1007/11780342_20.
- 603 26 Thomas Ehrhard and Laurent Regnier. Uniformity and the Taylor expansion of ordinary
604 λ -terms. *Theoretical Computer Science*, 403(2-3), 2008. doi:10.1016/j.tcs.2008.06.001.
- 605 27 Marcelo Fiore, Nicola Gambino, Martin Hyland, and Glynn Winskel. The cartesian closed
606 bicategory of generalised species of structures. *J. of the London Mathematical Society*, 77(1):203–
607 220, 2008. doi:10.1112/jlms/jdm096.
- 608 28 Marcelo Fiore, Nicola Gambino, Martin Hyland, and Glynn Winskel. Relative pseudo-
609 monads, Kleisli bicategories, and substitution monoidal structures. *Selecta Mathematica*,
610 24(3):2791–2830, Nov 2017. doi:10.1007/s00029-017-0361-3.
- 611 29 Zeinab Galal. A Profunctorial Scott Semantics. In *5th International Conference on Formal*
612 *Structures for Computation and Deduction (FSCD 2020)*, volume 167 of *Leibniz International*
613 *Proceedings in Informatics (LIPICs)*, pages 16:1–16:18, Dagstuhl, Germany, 2020. Schloss
614 Dagstuhl–Leibniz-Zentrum für Informatik. doi:10.4230/LIPICs.FSCD.2020.16.
- 615 30 Nicola Gambino and André Joyal. On operads, bimodules and analytic functors. *Memoirs of*
616 *the American Mathematical Society*, 249(1184):0–0, Sep 2017. doi:10.1090/memo/1184.
- 617 31 Philippa Gardner. Discovering needed reductions using type theory. In *Theoretical Aspects*
618 *of Computer Software, International Conference TACS '94*, volume 789 of *Lecture Notes in*
619 *Computer Science*, pages 555–574. Springer, 1994. doi:10.1007/3-540-57887-0_115.
- 620 32 Jean-Yves Girard. Linear logic. *Theoretical Computer Science*, 50(1):1 – 101, 1987. doi:
621 https://doi.org/10.1016/0304-3975(87)90045-4.
- 622 33 Giulio Guerrieri and Giulio Manzonetto. The bang calculus and the two Girard’s translations.
623 In *Proceedings Joint International Workshop on Linearity & Trends in Linear Logic and*
624 *Applications, Linearity-TLLA@FLoC 2018*, volume 292 of *EPTCS*, pages 15–30, 2018. doi:
625 10.4204/EPTCS.292.2.
- 626 34 Martin Hyland. Classical lambda calculus in modern dress. *Mathematical Structures in*
627 *Computer Science*, 27(5):762–781, 2017. doi:10.1017/S0960129515000377.
- 628 35 André Joyal. Foncteurs analytiques et espèces de structures. In *Combinatoire énumérative*,
629 pages 126–159, Berlin, Heidelberg, 1986. Springer Berlin Heidelberg.

- 630 36 Gregory Maxwell Kelly. A unified treatment of transfinite constructions for free algebras, free
631 monoids, colimits, associated sheaves, and so on. *Bulletin of the Australian Mathematical*
632 *Society*, 22(1):1–83, 1980. doi:10.1017/S0004972700006353.
- 633 37 Jean-Louis Krivine. Lambda-calculus, types and models. In *Ellis Horwood series in computers*
634 *and their applications*, 1993.
- 635 38 Ugo Dal Lago and Thomas Leventis. On the Taylor expansion of probabilistic lambda-
636 terms. In Herman Geuvers, editor, *4th International Conference on Formal Structures for*
637 *Computation and Deduction, FSCD 2019, June 24-30, 2019, Dortmund, Germany*, volume
638 131 of *LIPICs*, pages 13:1–13:16. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2019.
639 doi:10.4230/LIPICs.FSCD.2019.13.
- 640 39 Paul Blain Levy. Call-by-Push-Value: A Subsuming Paradigm. In *Typed Lambda Calculi*
641 *and Applications, 4th International Conference, TLCA'99*, volume 1581 of *Lecture Notes in*
642 *Computer Science*, page 228–242. Springer, 1999. doi:10.1007/3-540-48959-2_17.
- 643 40 Fosco Loregian. This is the (co)end, my only (co)friend, 2015. arXiv:1501.02503.
- 644 41 Damiano Mazza, Luc Pellissier, and Pierre Vial. Polyadic approximations, fibrations and
645 intersection types. *Proc. ACM Program. Lang.*, 2(POPL):6:1–6:28, 2018. doi:10.1145/
646 3158094.
- 647 42 Paul-André Melliès and Noam Zeilberger. Functors are type refinement systems. *SIGPLAN*
648 *Not.*, 50(1):3–16, January 2015. doi:10.1145/2775051.2676970.
- 649 43 Federico Olimpieri. Intersection Type Distributors, 2020. URL: [https://arxiv.org/abs/](https://arxiv.org/abs/2002.01287)
650 [2002.01287](https://arxiv.org/abs/2002.01287), arXiv:2002.01287.
- 651 44 Federico Olimpieri. *Intersection Types and Resource Calculi in the Denotational Semantics of*
652 *Lambda-Calculus*. PhD thesis, Aix-Marseille Université, 2020.
- 653 45 Luca Paolini and Simona Ronchi Della Rocca. Call-by-value solvability. *RAIRO Theor.*
654 *Informatics Appl.*, 33(6):507–534, 1999. doi:10.1051/ita:1999130.
- 655 46 Alex K. Simpson. Reduction in a linear lambda-calculus with applications to operational
656 semantics. In *Term Rewriting and Applications, 16th International Conference, RTA 2005*,
657 volume 3467 of *Lecture Notes in Computer Science*, pages 219–234. Springer, 2005. doi:
658 10.1007/978-3-540-32033-3_17.
- 659 47 Takeshi Tsukada, Kazuyuki Asada, and C.-H. Luke Ong. Generalised Species of Rigid Resource
660 Terms. In *Proceedings of the 32nd Annual ACM/IEEE Symposium on Logic in Computer*
661 *Science, LICS 2017, 2017*. doi:10.1109/LICS.2017.8005093.
- 662 48 Takeshi Tsukada, Kazuyuki Asada, and C.-H. Luke Ong. Species, profunctors and Taylor ex-
663 pansion weighted by SMCC: A unified framework for modelling nondeterministic, probabilistic
664 and quantum programs. In *Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic*
665 *in Computer Science, LICS '18*, pages 889–898, 2018. doi:10.1145/3209108.3209157.
- 666 49 Pierre Vial. Infinitary intersection types as sequences: A new answer to Klop’s problem. In
667 *32nd Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2017*, pages 1–12.
668 IEEE Computer Society, 2017. doi:10.1109/LICS.2017.8005103.

669 **A Appendix**

670 **Bicategories in a Nutshell [3, 6].** Intuitively, a bicategory is a category with “morphisms
671 between morphisms”, that is, where each hom-set itself carries the structure of a category,
672 but the composition of morphisms is only associative up to an isomorphism, and similarly
673 for the identities laws. Formally, a *bicategory* \mathcal{C} consists of:

- 674 ■ a set $\text{ob}(\mathcal{C})$ of *objects*, also called *0-cells* and denoted by A, B, C, \dots ;
- 675 ■ for all $A, B \in \text{ob}(\mathcal{C})$, a category $\mathcal{C}(A, B)$; objects in $\mathcal{C}(A, B)$ are called *1-cells* or *morphisms*
676 from A to B ; while arrows in $\mathcal{C}(A, B)$ (between 1-cells from A to B) are called *2-cells* or
677 *2-morphisms*; composition of 2-cells is generally called *vertical composition*;
- 678 ■ for every $A, B, C \in \text{ob}(\mathcal{C})$, a bifunctor

$$679 \quad \circ_{A,B,C}: \mathcal{C}(B, C) \times \mathcal{C}(A, B) \rightarrow \mathcal{C}(A, C)$$

680 called *horizontal composition* (often the indices A, B, C in $\circ_{A,B,C}$ are omitted); hence,
681 for all 1-cells $F: A \rightarrow B$, $F': A \rightarrow B$ and $G: B \rightarrow C$, $G': B \rightarrow C$, and for all 2-cells
682 $\alpha: F \Rightarrow F'$ and $\beta: G \Rightarrow G'$, we have

$$683 \quad \text{a 1-cell } G \circ_{A,B,C} F: A \rightarrow C \quad \text{a 2-cell } \beta \circ_{A,B,C} \alpha: (G \circ_{A,B,C} F) \Rightarrow (G' \circ_{A,B,C} F');$$

- 685 ■ for every $A \in \text{ob}(\mathcal{C})$ a functor $1_A: 1 \rightarrow \mathcal{C}(A, A)$; with an abuse of notation we identify
686 $1_A(\star)$ with 1_A and we call it the identity of A ;
- 687 ■ for all 1-cells $F: A \rightarrow B$, $G: B \rightarrow C$ and $H: C \rightarrow D$, a family of invertible 2-cells

$$688 \quad \alpha_{H,G,F}: H \circ (G \circ F) \cong (H \circ G) \circ F$$

689 expressing the associativity laws;


- 690 ■ for every 1-cell $F: A \rightarrow B$, two families of invertible 2-cells

$$691 \quad \lambda_F: 1_B \circ F \cong F \quad \rho_F: F \cong F \circ 1_A$$

692 expressing the identity laws.

693 This data is subject to additional coherence axioms. A *2-category* is a bicategory where the
694 associativity and identities are strict equalities, not only isomorphisms.

695 **► Definition 17 (Retraction).** Let D, E be 0-cells in a bicategory \mathcal{C} . A retraction of D to E
696 is a couple of 1-cells $i: E \rightarrow D$, $j: D \rightarrow E$ together with an invertible 2-cell β such that the
697 diagram below commute. We write $E \triangleleft D$ if there is a retraction of D to E .



699 **Coends.** Given a functor $F: C^{op} \times C \rightarrow \text{Set}$ we recall that the coend is the coequalizer of
700 the following diagram

$$701 \quad \sum_{c,c' \in C} C(c', c) \times F(c, c') \rightrightarrows \sum_{c \in C} F(c, c) \rightarrow \int^{c \in C} F(c, c)$$

702 Where the parallel arrows are given by left and right actions of F on morphisms $f \in C(c', c)$.
703 Since we work with coends in the category of set, we have that this coequalizer is actually
704 given by the quotient $\sum_{c \in C} F(c, c) / \sim$ where the equivalence relation is generated by the
705 rule $x \sim y$ iff $F(f, c')(x) = y, F(c, f)(y) = x$, for $f: c' \rightarrow c$.

706 We list the three fundamental lemmas of coend calculus [40].

707 ▶ **Lemma 18.** *Every cocontinuous functor preserves coends.*

708 ▶ **Lemma 19** (Fubini [40]). *Let $F : C^{op} \times C \times D^{op} \times D \rightarrow Set$ be a functor. We have*

$$709 \quad \int^{(c,d) \in C \times D} F(c, c, d, d) \cong \int^{c \in C} \int^d F(c, c, d, d) \cong \int^{d \in D} \int^{c \in C} F(c, c, d, d).$$

710 ▶ **Lemma 20** (Yoneda Ninja [40]). *Let $K, H : C \rightarrow Set$ be, respectively, a contravariant and*
711 *a covariant functor. We have the following natural isomorphisms*

$$712 \quad K(-) \cong \int^{c \in C} K(c) \times C(-, c) \quad H(-) \cong \int^{c \in C} H(c) \times C(c, -).$$

713 **Denotation under Reduction.** In what follows we do not explicitly state, for readability
714 reasons, when we apply Lemmas 18 and 19. For $\vec{\Gamma} = \langle \Gamma_1, \dots, \Gamma_n \rangle$ we set $\otimes \vec{\Gamma} = \otimes_{i=1}^n \Gamma_i$.

715 ▶ **Lemma 11** (Substitution). *Let S and T be terms. There is an isomorphism of functors*

$$716 \quad \varphi : Sub_{S,x,T} \cong \llbracket S\{T/x\} \rrbracket_{\vec{x}}$$

717 *natural in $\langle \Delta, a \rangle \in \text{ob}((!U^n)^{op} \times U)$ and such that $s(\varphi_{\Delta,a}(\langle \alpha_1, \alpha_2, \eta \rangle)) = s(\alpha_1) + s(\alpha_2)$.*

718 **Proof.** By induction on the structure of $S \in !\Lambda$, via lengthy coend manipulations.

719 If $S = x$ then

$$720 \quad Sub_{S,x,T}(\Delta, a) = \int^{\Gamma_0, \Gamma_1 \in !U^n} \int^{\vec{a} \in !U} \llbracket x \rrbracket_{\vec{x}}(\Gamma_0 \oplus \langle \vec{a} \rangle, a) \times \llbracket T^! \rrbracket_{\vec{x}}(\Gamma_1, \vec{a}) \times !U(\Delta, \Gamma_0 \otimes \Gamma_1).$$

721 By definition we have

$$722 \quad \cong \int^{\Gamma_0, \Gamma_1 \in !U^n} \int^{\vec{a} \in !U} !U^n(\Gamma_0 \oplus \langle \vec{a} \rangle, \langle \langle \rangle, \dots, \langle \rangle, \langle a \rangle \rangle) \times \llbracket T^! \rrbracket_{\vec{x}}(\Gamma_1, \vec{a}) \times !U(\Delta, \Gamma_0 \otimes \Gamma_1).$$

723 Then, by the structure of the product category

$$724 \quad \cong \int^{\Gamma_0, \Gamma_1 \in !U^n} \int^{\vec{a} \in !U} !U^n(\Gamma_0, \langle \langle \rangle, \dots, \langle \rangle \rangle) \times !U(\vec{a}, \langle a \rangle) \times \llbracket T^! \rrbracket_{\vec{x}}(\Gamma_1, \vec{a}) \times !U(\Delta, \Gamma_0 \otimes \Gamma_1).$$

725 Then, by Yoneda (Lemma 20) we have

$$726 \quad \cong \int^{\Gamma_1 \in !U^n} \int^{\vec{a} \in !U} !U(\vec{a}, \langle a \rangle) \times \llbracket T^! \rrbracket_{\vec{x}}(\Gamma_1, \vec{a}) \times !U(\Delta, \Gamma_1).$$

727 Again, by Yoneda (Lemma 20),

$$728 \quad \cong \int^{\Gamma_1 \in !U^n} \llbracket T^! \rrbracket_{\vec{x}}(\Gamma_1, \langle a \rangle) \times !U(\Delta, \Gamma_1).$$

729 Then, by applying Yoneda one more time on the context Γ and by definition of the denotation
730 of a box we can conclude. For what concerns the size, simply notice that $s(\tilde{\pi}) = s(\llbracket f \rrbracket \tilde{\pi})$.

731 The abstraction case follows from the *i.h.* immediately.

732 We do the application, the box case being similar to it. If $S = QR$ then

$$733 \quad Sub_{S,x,T}(\Delta, a) = \int^{\Gamma_1, \Gamma_2} \int^{\vec{a}} \llbracket QR \rrbracket_{\vec{x} \oplus (x)}(\Gamma_1 \oplus \langle \vec{a} \rangle, a) \times \llbracket T^! \rrbracket_{\vec{x}}(\Gamma_2, \vec{a}) \times !U^n(\Delta, \Gamma_1 \otimes \Gamma_2).$$

38:22 Categorifying Non-Idempotent Intersection Types

734 We develop $\llbracket QR \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma_1 \oplus \langle \vec{a} \rangle, a)$:

$$\begin{aligned} \llbracket QR \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma_1 \oplus \langle \vec{a} \rangle, a) &= \int^{\Gamma'_1 \oplus \langle \vec{a}_1 \rangle, \Gamma'_2 \oplus \langle \vec{a}_2 \rangle} \int^{\vec{b}} \llbracket Q \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_1 \oplus \langle \vec{a}_1 \rangle, \iota(\vec{b}, a)) \\ &\times \llbracket R \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_2 \oplus \langle \vec{a}_2 \rangle, \vec{b}) \times !U^n(\Gamma_1 \oplus \langle \vec{a} \rangle, \Gamma'_1 \oplus \langle \vec{a}_1 \rangle \otimes \Gamma'_2 \oplus \langle \vec{a}_2 \rangle). \end{aligned}$$

736 By the structure of the product category, we have

$$\begin{aligned} \llbracket QR \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma_1 \oplus \langle \vec{a} \rangle, a) &= \int^{\Gamma'_1 \Gamma'_2} \int^{\vec{a}_1, \vec{a}_2} \int^{\vec{b}} \llbracket Q \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_1 \oplus \langle \vec{a}_1 \rangle, \iota(\vec{b}, a)) \\ &\times \llbracket R \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_2 \oplus \langle \vec{a}_2 \rangle, \vec{b}) \times !U^n(\Gamma_1, \Gamma'_1 \otimes \Gamma'_2) \times !U(\vec{a}, \vec{a}_1 \oplus \vec{a}_2). \end{aligned}$$

738 We apply Yoneda (Lemma 20) on Γ_1 and on \vec{a} and we get

$$\begin{aligned} Sub_{S,x,T}(\Delta, a) &\cong \int^{\Gamma'_1 \Gamma'_2, \Gamma_2} \int^{\vec{b}, \vec{a}_1, \vec{a}_2} \llbracket Q \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_1 \oplus \langle \vec{a}_1 \rangle, \iota(\vec{b}, a)) \times \llbracket R \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_2 \oplus \langle \vec{a}_2 \rangle, \vec{b}) \\ &\times \llbracket T^! \rrbracket_{\bar{x}}(\Gamma_2, \vec{a}_1 \oplus \vec{a}_2) \times !U^n(\Delta, (\Gamma'_1 \otimes \Gamma'_2) \otimes \Gamma_2). \end{aligned}$$

740 By a simple inspection of the definition of the denotation of a box, we can rewrite it as

$$\begin{aligned} &\cong \int^{\Gamma'_i, \Gamma_{\vec{a}_i}, \Gamma_2} \int^{\vec{b}, \vec{a}_1, \vec{a}_2} \llbracket Q \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_1 \oplus \langle \vec{a}_1 \rangle, \iota(\vec{b}, a)) \times \llbracket R \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_2 \oplus \langle \vec{a}_2 \rangle, \vec{b}) \times \llbracket T^! \rrbracket_{\bar{x}}(\Gamma_{\vec{a}_1}, \vec{a}_1) \\ &\times \llbracket T^! \rrbracket_{\bar{x}}(\Gamma_{\vec{a}_2}, \vec{a}_2) \times !U^n(\Gamma_2, \bigotimes_{i \in \{1,2\}} \Gamma_{\vec{a}_i}) \times !U^n(\Delta, (\Gamma'_1 \otimes \Gamma'_2) \otimes \Gamma_2). \end{aligned}$$

742 Where, if we set $\vec{a}_i = \langle a_{i,1}, \dots, a_{i,k_i} \rangle$, $\llbracket T^! \rrbracket_{\bar{x}}(\Gamma_{\vec{a}_i}, \vec{a}_i) = \prod_{j \in k_i} \llbracket T^! \rrbracket_{\bar{x}}(\Gamma_{i,j}, a_{i,j})$ and $\bigotimes_{i \in \{1,2\}} \Gamma_{\vec{a}_i} =$
743 $\bigotimes_{j \in k_i} \Gamma_{i,j}$ with $i \in \{1,2\}$. We apply Yoneda on Γ_2

$$\begin{aligned} &\cong \int^{\Gamma'_i, \Gamma_{\vec{a}_i}} \int^{\vec{b}, \vec{a}_1, \vec{a}_2} \llbracket Q \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_1 \oplus \langle \vec{a}_1 \rangle, \iota(\vec{b}, a)) \times \llbracket R \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_2 \oplus \langle \vec{a}_2 \rangle, \vec{b}) \times \llbracket T^! \rrbracket_{\bar{x}}(\Gamma_{\vec{a}_1}, \vec{a}_1) \\ &\times \llbracket T^! \rrbracket_{\bar{x}}(\Gamma_{\vec{a}_2}, \vec{a}_2) \times !U^n(\Delta, (\Gamma'_1 \otimes \Gamma'_2) \otimes (\bigotimes_{i \in \{1,2\}} \Gamma_{\vec{a}_i})). \end{aligned}$$

745 Now, by the *symmetry* of the tensor product \otimes and by the fact that functors preserves
746 isomorphisms, we get

$$\begin{aligned} &\cong \int^{\Gamma'_i, \Gamma_{\vec{a}_i}} \int^{\vec{b}, \vec{a}_1, \vec{a}_2} \llbracket Q \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_1 \oplus \langle \vec{a}_1 \rangle, \iota(\vec{b}, a)) \times \llbracket R \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_2 \oplus \langle \vec{a}_2 \rangle, \vec{b}) \times \llbracket T^! \rrbracket_{\bar{x}}(\Gamma_{\vec{a}_1}, \vec{a}_1) \\ &\times \llbracket T^! \rrbracket_{\bar{x}}(\Gamma_{\vec{a}_2}, \vec{a}_2) \times !U^n(\Gamma_2, \bigotimes_{i \in \{1,2\}} \Gamma_{\vec{a}_i}) \times !U^n(\Delta, ((\Gamma'_1 \otimes \Gamma_{\vec{a}_1}) \otimes (\Gamma'_2 \otimes \Gamma_{\vec{a}_2}))). \end{aligned}$$

748 Now, if we apply Yoneda twice to $\Gamma'_i \otimes \Gamma_{\vec{a}_i}$, we get

$$\begin{aligned} &\cong \int^{\Gamma'_i, \Gamma_{\vec{a}_i}, \Delta_i} \int^{\vec{b}, \vec{a}_1, \vec{a}_2} \llbracket Q \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma_1 \oplus \langle \vec{a}_1 \rangle, \iota(\vec{b}, a)) \times \llbracket R \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_2 \oplus \langle \vec{a}_2 \rangle, \vec{b}) \times \llbracket T^! \rrbracket_{\bar{x}}(\Gamma'_{\vec{a}_1}, \vec{a}_1) \\ &\times \llbracket T^! \rrbracket_{\bar{x}}(\Gamma_{\vec{a}_2}, \vec{a}_2) \times !U^n(\Delta, \Delta_1 \otimes \Delta_2) \otimes !U^n(\Delta_1, \Gamma'_1 \otimes \bigotimes_{i \in \{1,2\}} \Gamma_{\vec{a}_i}) \otimes !U^n(\Delta_2, \Gamma'_2 \otimes \bigotimes_{i \in \{1,2\}} \Gamma_{\vec{a}_2}). \end{aligned}$$

750 By co-continuity and commutativity, and by applying Yoneda (Lemma 20) twice, we have

$$\begin{aligned} &\cong \int^{\vec{b}} \int^{\Gamma'_1, \Gamma_{\vec{a}_1}, \Delta_1, \Phi_1} \int^{\vec{a}_1} \llbracket Q \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_1 \oplus \langle \vec{a}_1 \rangle, \iota(\vec{b}, a)) \times \llbracket T^! \rrbracket_{\bar{x}}(\Phi_1, \vec{a}_1) \\ &\times !U^n(\Phi_1, \bigotimes_{i \in \{1,2\}} \Gamma'_{\vec{a}_i}) \times U^n(\Delta_1, \Gamma'_1 \otimes \Phi_1) \\ &\times \int^{\Gamma'_2, \Gamma_{\vec{a}_2}, \Delta_2, \Phi_2} \int^{\vec{a}_2} \llbracket R \rrbracket_{\bar{x} \oplus \langle x \rangle}(\Gamma'_2 \oplus \langle \vec{a}_2 \rangle, \vec{b}) \times \llbracket T^! \rrbracket_{\bar{x}}(\Phi_2, \vec{a}_2) \\ &\times !U^n(\Phi_2, \bigotimes_{i \in \{1,2\}} \Gamma'_{\vec{a}_2}) \times U^n(\Delta_2, \Gamma'_2 \otimes \Phi_2) \times !U^n(\Delta, \Delta_1 \otimes \Delta_2). \end{aligned}$$

751

752 By definition, the former coend is just

$$753 \int^{\vec{b}} \int^{\Delta_1, \Delta_2} \text{Sub}_{Q,x,T}(\Delta_1, \iota(\vec{b}, a)) \times \text{Sub}_{R,x,T}(\Delta_2, \vec{b}) \times !U^n(\Delta, \Delta_1 \otimes \Delta_2).$$

754 We remark that, forgetting the equivalence relation, the built isomorphism

$$755 \text{Sub}_{S,x,T}(\Delta, a) \cong \int^{\vec{b}} \text{Sub}_{Q,x,T}(\Delta_1, \iota(\vec{b}, a)) \times \text{Sub}_{R,x,T}(\Delta_2, \vec{b}) \times !U^n(\Delta, \Delta_1 \otimes \Delta_2)$$

756 consists of the following map

$$757 \langle \vec{a}, \langle \vec{b}, \langle \Gamma_1, \Gamma_2, \langle \langle \Gamma'_1 \oplus \langle \vec{a}_1 \rangle, \Gamma'_2 \oplus \langle \vec{a}_2 \rangle, \langle \alpha_1, \alpha_2, \eta_1 \rangle, \langle \langle \vec{\Gamma} = \langle \Gamma_{2,1}, \dots, \Gamma_{2, \text{len}(\vec{a})} \rangle, \vec{\beta} = \langle \beta_1, \dots, \beta_{\text{len}(\vec{a})} \rangle, \eta_2 \rangle \rangle, \theta \rangle \rangle \mapsto$$

$$758 \langle \vec{b}, \langle \Gamma'_1 \otimes \bigotimes \Gamma_{\vec{a}_1}, \alpha_1, \langle \vec{\beta}_{\vec{a}_1}, 1_{\bigotimes \Gamma_{\vec{a}_1}} \rangle, 1_{\Gamma'_{1 \otimes \bigotimes \Gamma_{\vec{a}_1}}} \rangle, \langle \Gamma'_2 \otimes \bigotimes \Gamma_{\vec{a}_2}, \alpha_2, \langle \vec{\beta}_{\vec{a}_2}, 1_{\bigotimes \Gamma_{\vec{a}_2}} \rangle, 1_{\Gamma'_{2 \otimes \bigotimes \Gamma_{\vec{a}_2}}} \rangle,$$

$$759 ((\eta_1 \otimes (\sigma^* \circ \eta_2)) \circ \theta) \circ \tau \rangle$$

$$760$$

$$761$$

762 where

- 763 ■ $\theta : \Delta \rightarrow \Gamma_1 \otimes \Gamma_2, \alpha_1 \in \llbracket Q \rrbracket_{\vec{x}}(\Gamma'_1 \oplus \langle \vec{a}_1 \rangle, \iota(\vec{b}, a))$ and $\alpha_2 \in \llbracket R \rrbracket_{\vec{x}}(\Gamma'_2 \oplus \langle \vec{a}_2 \rangle, \vec{b})$;
- 764 ■ $\langle \eta_1, f = \langle \sigma, \vec{f} \rangle \rangle : \Gamma_1 \oplus \langle \vec{a} \rangle \rightarrow \Gamma'_1 \oplus \langle \vec{a}_1 \rangle \otimes \Gamma'_2 \oplus \langle \vec{a}_2 \rangle$ and $\eta_2 : \Gamma_2 \rightarrow \bigotimes \vec{\Gamma}, \vec{\beta} \in \llbracket T \rrbracket_{\vec{x}}(\Gamma_2, \vec{a})$;
- 765 ■ $[f]\vec{\Gamma} = \Gamma_{\vec{a}_1} \otimes \Gamma_{\vec{a}_2}$ and $[f]\vec{\beta} = \vec{\beta}_{\vec{a}_1} \oplus \vec{\beta}_{\vec{a}_2}$;
- 766 ■ $\tau : (\Gamma'_1 \otimes \Gamma'_2) \otimes (\bigotimes \Gamma_{\vec{a}_1} \otimes \bigotimes \Gamma_{\vec{a}_2}) \rightarrow (\Gamma'_1 \otimes \bigotimes \Gamma_{\vec{a}_1}) \otimes (\Gamma'_2 \otimes \bigotimes \Gamma_{\vec{a}_2})$ is the obvious symmetry.

767 By definition, we have (for $S = QR$)

$$768 \llbracket S\{T/x\} \rrbracket_{\vec{x}}(\Delta, a) = \int^{\vec{b}} \int^{\Delta_1, \Delta_2} \llbracket Q\{T/x\} \rrbracket_{\vec{x}}(\Delta_1, \iota(\vec{b}, a)) \times \llbracket R\{T/x\} \rrbracket_{\vec{x}}(\Delta_2, \vec{b}) \times !U^n(\Delta, \Delta_1 \otimes \Delta_2).$$

769 By *i.h.*, we get two isomorphisms $\llbracket Q\{T/x\} \rrbracket_{\vec{x}}(\Delta_1, \iota(\vec{b}, a)) \cong \text{Sub}_{Q,x,T}(\Delta_1, \iota(\vec{b}, a))$ and
 770 $\llbracket R\{T/x\} \rrbracket_{\vec{x}}(\Delta_2, \vec{b}) \cong \text{Sub}_{R,x,T}(\Delta_2, \vec{b})$. We have our isomorphism, since isomorphisms are
 771 preserved by products and coends. Then we can conclude, since morphism actions do not
 772 change size of points and we have $s(\vec{\beta}) = s(\vec{\beta}_{\vec{a}_1}) + s(\vec{\beta}_{\vec{a}_2})$. ◀

773 **Failure of Subject Reduction with Subtyping for Polyadic Terms.** We recall the definition
 774 of linear polyadic calculus in the framework of bang calculus.

$$775 p, q ::= x \mid \lambda \langle x_1, \dots, x_k \rangle . p \mid pq \mid \langle p_1, \dots, p_k \rangle \mid \perp$$

776 Terms are taken up to α -equivalence and up to linearity with respect to \perp (*i.e.*, $\lambda \vec{x} . \perp =$
 777 $p \langle \perp \rangle = \perp$, etc.⁷). The reduction \rightarrow_p is the contextual closure of the following base case:

$$778 (\lambda \vec{x} . p) \vec{q} \mapsto_p \begin{cases} p\{\vec{q}/\vec{x}\} & \text{if } \text{len}(\vec{q}) = \text{len}(\vec{x}) \\ \perp & \text{otherwise.} \end{cases}$$

779 Since we want to link a calculus of approximants to intersection type distributors, the first
 780 thing to check is that the calculus satisfies subject reduction and expansion within our system

⁷ This is slightly different from the original definition of [41], but being up to linearity simplify calculations.

38:24 Categorifying Non-Idempotent Intersection Types

781 $\mathcal{R}_{\rightarrow}$. Let $\zeta = \langle \vec{x}_1, \dots, \vec{x}_n \rangle$ and $\Delta = \langle \vec{a}_1, \dots, \vec{a}_n \rangle$. We write $\zeta : \Delta$ for $\vec{x}_1 : \vec{a}_1, \dots, \vec{x}_n : \vec{a}_n$. We
 782 give the following naive type assignment:

$$\begin{array}{c}
 783 \quad \frac{f : a' \rightarrow a}{\langle \rangle : \langle \rangle, \dots, \langle x \rangle : \langle a' \rangle, \dots, \langle \rangle : \langle \rangle \vdash x : a} \quad \frac{(\zeta_i : \Gamma_i \vdash q_i)_{i=1}^k \quad \eta : \Delta \rightarrow \bigotimes_{i=1}^k \Gamma_i}{[\eta](\bigotimes_{i=1}^k \zeta_i) : \Delta \vdash \langle q_1, \dots, q_k \rangle : \langle a_1, \dots, a_k \rangle} \\
 784 \quad \frac{\zeta \oplus \langle \vec{x} \rangle : \Delta \oplus \langle \vec{a} \rangle \vdash p : a}{\zeta : \Delta \vdash \lambda \vec{x}. p : \vec{a} \Rightarrow a} \quad \frac{\zeta_0 : \Gamma_0 \vdash p : \vec{a} \Rightarrow a \quad \zeta_1 : \Gamma_1 \vdash q \quad \eta : \Delta \rightarrow \Gamma_0 \otimes \Gamma_1}{[\eta](\zeta_0 \otimes \zeta_1) : \Delta \vdash pq : a} \\
 785 \quad \zeta : \Delta \vdash \lambda \vec{x}. p : \vec{a} \Rightarrow a
 \end{array}$$

786 where in the application case the left action $[\eta]\zeta$ means only that the positions of variables
 787 in ζ are rearranged in accordance with the permutation induced by the morphism η . This
 788 is reasonable and necessary, since the morphism η can in general rearrange the position
 789 of types. This means that if $\zeta = \langle \vec{x}_1, \dots, \vec{x}_n \rangle$ and $\eta = \langle \langle \sigma_1, \vec{f}_1 \rangle, \dots, \langle \sigma_n, \vec{f}_n \rangle \rangle$ then $[\eta]\zeta =$
 790 $\langle [\sigma_1]\vec{x}_1, \dots, [\sigma_n]\vec{x}_n \rangle$ where $[\sigma]\langle x_1, \dots, x_k \rangle = \langle x_{\sigma(1)}, \dots, x_{\sigma(k)} \rangle$ is just the left action of the
 791 symmetry group. It is easy to see that \perp is not typable in the type system above.

792 **► Example 21.** We present a counter-example for the subject reduction of the former system.
 793 Take the polyadic term $p = (\lambda x.x(\lambda \langle \rangle.y_1 \langle \rangle, \lambda \langle f \rangle.y_2 \langle f \rangle))(\lambda \langle z_1, z_2 \rangle.z_1 \langle z_2 \rangle)$. This term clearly
 794 reduces to \perp , but it is typable in the former type system. Let $\pi =$

$$\begin{array}{c}
 795 \quad \frac{g : b' \rightarrow b}{\langle x \rangle : \langle b' \rangle, \langle \rangle \vdash x : b} \quad \frac{\langle \rangle : \langle \rangle, \langle y_1 \rangle : \langle \langle \rangle \Rightarrow a \rangle \vdash \lambda \langle \rangle.y_1 \langle \rangle : \langle \rangle \Rightarrow a \quad \langle \rangle : \langle \rangle, \langle y_1 \rangle : \langle \langle c \rangle \Rightarrow a \rangle \vdash \lambda \langle f \rangle.y_1 \langle f \rangle : \langle c \rangle \Rightarrow a}{\langle x \rangle : \langle b' \rangle, \langle y_1, y_2 \rangle : \langle \langle \rangle \Rightarrow a, \langle c \rangle \Rightarrow a \rangle \vdash x(\lambda \langle \rangle.y_1 \langle \rangle, \lambda \langle f \rangle.y_2 \langle f \rangle) : a} \\
 \frac{\langle y_1, y_2 \rangle : \langle \langle \rangle \Rightarrow a, \langle c \rangle \Rightarrow a \rangle \vdash \lambda \langle x \rangle.x(\lambda \langle \rangle.y_1 \langle \rangle, \lambda \langle f \rangle.y_2 \langle f \rangle) : \langle b' \rangle \Rightarrow a}{}
 \end{array}$$

796 Where $c = \langle \rangle \Rightarrow a$ and $b' = \langle \langle c \rangle \Rightarrow a, \langle \rangle \Rightarrow a \rangle \Rightarrow a$ and $b = \langle \langle \rangle \Rightarrow a, \langle c \rangle \Rightarrow a \rangle \Rightarrow a$
 797 the morphism g being of the shape $\langle \sigma, 1_{\langle \rangle \Rightarrow a}, 1_{\langle c \rangle \Rightarrow a} \rangle \Rightarrow 1$ with sigma being the obvious
 798 permutation. Consider $\rho =$

$$\begin{array}{c}
 799 \quad \frac{\langle z_1 \rangle : \langle \langle c \rangle \Rightarrow a \rangle \vdash z_1 : \langle c \rangle \Rightarrow a \quad \langle z_2 \rangle : \langle c \rangle \vdash z_2 : c}{\langle z_1, z_2 \rangle : \langle \langle c \rangle \Rightarrow a, c \rangle \vdash z_1 \langle z_2 \rangle : a} \\
 \vdash \lambda \langle z_1, z_2 \rangle.z_1 \langle z_2 \rangle : \langle \langle c \rangle \Rightarrow a, c \rangle \Rightarrow a
 \end{array}$$

800 Now take $\pi' =$

$$\begin{array}{c}
 801 \quad \frac{\langle y_1, y_2 \rangle : \langle \langle \rangle \Rightarrow a, \langle a \rangle \Rightarrow a \rangle \vdash \lambda \langle x \rangle.x(\lambda \langle \rangle.y_1 \langle \rangle, \lambda \langle f \rangle.y_2 \langle f \rangle) : \langle b' \rangle \Rightarrow a \quad \vdash \lambda \langle z_1, z_2 \rangle.z_1 \langle z_2 \rangle : b'}{\langle y_1, y_2 \rangle : \langle \langle \rangle \Rightarrow a, \langle a \rangle \Rightarrow a \rangle \vdash p : a}
 \end{array}$$

802 The term p reduces to \perp . Indeed,

$$\begin{array}{c}
 803 \quad p = (\lambda x.x(\lambda \langle \rangle.y_1 \langle \rangle, \lambda \langle f \rangle.y_2 \langle f \rangle))(\lambda \langle z_1, z_2 \rangle.z_1 \langle z_2 \rangle) \\
 804 \quad \rightarrow_p (\lambda \langle z_1, z_2 \rangle.z_1 \langle z_2 \rangle)(\lambda \langle \rangle.y_1 \langle \rangle, \lambda \langle f \rangle.y_2 \langle f \rangle) \\
 805 \quad \rightarrow_p (\lambda \langle \rangle.y_1 \langle \rangle)(\lambda \langle f \rangle.y_2 \langle f \rangle) \rightarrow_p \perp
 \end{array}$$

807 Therefore, $p \rightarrow_p^* \perp$ and p is typable, while \perp it is not. The problem relies completely in
 808 the variable rule: the subtyping feature of the system is not detected by the syntax of the
 809 standard polyadic calculus. If we want to find an appropriate term language for our system,
 810 whose elements are also approximants of ordinary bang terms, we need to take seriously the
 811 qualitative information produced by the subtyping.