

Lambda theories of effective lambda models

Chantal Berline¹, Giulio Manzonetto^{1,2,*}, and Antonino Salibra²
chantal.berline@pps.jussieu.fr, {gmanzone, salibra}@dsi.unive.it

¹ Laboratoire PPS, CNRS-Université Paris 7,
2, place Jussieu (case 7014), 75251 Paris Cedex 05, France

² Università Ca'Foscari di Venezia,
Dipartimento di Informatica Via Torino 155, 30172 Venezia, Italy

Abstract. A longstanding open problem is whether there exists a non-syntactical model of the untyped λ -calculus whose theory is exactly the least λ -theory $\lambda\beta$. In this paper we investigate the more general question of whether the equational/order theory of a model of the untyped λ -calculus can be recursively enumerable (r.e. for brevity). We introduce a notion of *effective model* of λ -calculus, which covers in particular all the models individually introduced in the literature, and prove that the equational theory of an effective model cannot be $\lambda\beta$, $\lambda\beta\eta$. In other results of the paper we show that the order theory of an effective model cannot be r.e. and that no effective model living in the stable or strongly stable semantics has an r.e. equational theory. Concerning Scott's semantics, we investigate the class of graph models and prove that no order theory of a graph model can be r.e., and that there exists an effective graph model whose equational/order theory is the minimum one. Finally, we show that the class of graph models enjoys a kind of downwards Löwenheim-Skolem theorem.

Keywords: Lambda calculus, Effective lambda models, Recursively enumerable lambda theories, Graph models, Löwenheim-Skolem theorem.

1 Introduction

Lambda theories are equational extensions of the untyped λ -calculus closed under derivation. They arise by syntactical or semantic considerations. Indeed, a λ -theory may correspond to a possible operational (observational) semantics of λ -calculus, as well as it may be induced by a model of λ -calculus through the kernel congruence relation of the interpretation function. Although researchers have mainly focused their interest on a limited number of them, the class of λ -theories constitutes a very rich and complex structure (see [1, 4, 5]).

Topology is at the center of the known approaches to giving models of the untyped λ -calculus. The first model, found by Scott in 1969 in the category of complete lattices and Scott continuous functions, was successfully used to show that all the unsolvable λ -terms can be consistently equated. After Scott, a large

* Corresponding author

number of mathematical models for λ -calculus, arising from syntax-free constructions, have been introduced in various categories of domains and were classified into semantics according to the nature of their representable functions, see e.g. [1, 4, 19]. Scott continuous semantics [22] is given in the category whose objects are complete partial orders and morphisms are Scott continuous functions. The stable semantics (Berry [7]) and the strongly stable semantics (Bucciarelli-Ehrhard [8]) are refinements of the continuous semantics, introduced to capture the notion of “sequential” Scott continuous function. In each of these semantics it is possible to build up 2^{\aleph_0} models inducing pairwise distinct λ -theories [16, 17]. Nevertheless, all these semantics are equationally *incomplete* (see [15, 2, 20, 21]) in the sense that they do not represent all the possible consistent λ -theories. It is interesting to note that there are very few known equational theories of λ -models living in these semantics that can be described syntactically: namely, the theory of Böhm trees and variants of it. None of these theories is recursively enumerable (r.e. for brevity).

Berline has raised in [4] the natural question of whether, given a class of models of λ -calculus, there is a minimum λ -theory represented by it. This question relates to the longstanding open problem proposed by Barendregt about the existence of a continuous model or, more generally, of a non-syntactical model of $\lambda\beta$ ($\lambda\beta\eta$). Di Gianantonio, Honsell and Plotkin [12] have shown that Scott continuous semantics admits a minimum theory, at least if we restrict to extensional models. Another result of [12], in the same spirit, is the construction of an extensional model whose theory is $\lambda\beta\eta$, a fortiori minimal, in a weakly-continuous semantics. However, the construction of this model starts from the term model of $\lambda\beta\eta$, and hence it cannot be seen as having a purely non syntactical presentation. More recently, Bucciarelli and Salibra [9, 10] have shown that the class of graph models admits a minimum λ -theory different from $\lambda\beta$. Graph models, isolated in the seventies by Plotkin, Scott and Engeler (see e.g. [1]) within the continuous semantics, have proved useful for showing the consistency of extensions of λ -calculus and for studying operational features of λ -calculus (see [4]).

In this paper we investigate the related question of whether the equational theory of a model can be r.e. As far as we know, this problem was first raised in [5], where it is conjectured that no graph model can have an r.e. theory. But we expect that this could indeed be true for all models living in the continuous semantics, and its refinements.

We find natural to concentrate on models with built-in effectivity properties. It seems indeed reasonable to think that, if effective models do not even succeed to have an r.e. theory, then it is unlikely that the other ones may succeed. Another justification for considering effective models comes from a previous result obtained for typed λ -calculus. Indeed, it was proved in [3] that there exists a non-syntactical model of Girard’s system F whose theory is $\lambda\beta\eta$. This model lives in Scott’s continuous semantics, and can easily be checked to be “effective” in the same spirit as in the present paper (see [3, Appendix C] for a sketchy presentation of the model).

Starting from the known notion of an effective domain, we introduce a general notion of an *effective model* of λ -calculus and we study the main properties of these models³. Effective models are omni-present in the continuous, stable and strongly stable semantics. In particular, all the models which have been introduced individually in the literature can easily be proved effective. The following is the first result of the paper:

- (i) The equational theory of an effective model of untyped λ -calculus cannot be $\lambda\beta$, $\lambda\beta\eta$.

Concerning the existence of a non-syntactical effective model with an r.e. theory, we are able to give a definite negative answer for all order theories and for the equational theories of all stable and strongly stable models:

- (ii) No effective model can have an r.e. order theory.
- (iii) No effective model living in the stable or strongly stable semantics has an r.e. equational theory.

Concerning Scott continuous semantics, the problem looks much more difficult. We concentrate here on the class of graph models (see [5, 6, 9–11] for earlier investigation of this class) and show that:

- (iv) No effective graph model, freely generated by a “partial model” which is finite modulo its group of automorphisms, has an r.e. equational theory.
- (v) There exists a minimum order graph theory (where “graph theory” means “theory of a graph model”), which happens to be the theory of an effective graph model.
- (vi) No order graph theory can be r.e.
- (vii) (Löwenheim-Skolem theorem for graph models) Every equational/order graph theory is the theory of a graph model having a countable carrier set.

The last result positively answers Question 3 in [4, Section 6.3] for the class of graph models and has the consequence that every graph theory (we know from Kerth [16] that there exists a continuum of them) is the theory of a graph model whose carrier set is the set of natural numbers.

The central technical device used in this paper is Visser’s result [25] stating that the complements of β -closed r.e. sets of λ -terms enjoy the finite intersection property (see Theorem 2).

2 Preliminaries

We denote by \mathbb{N} the set of natural numbers. The complement of a recursively enumerable set (r.e. set for short) is called a *co-r.e.* set. If both A and its complement are r.e., A is called *decidable*. We will denote by \mathcal{RE} the collection of all r.e. subsets of \mathbb{N} .

³ As far as we know, only Giannini and Longo [13] have introduced a notion of an effective model; but their definition is *ad hoc* for two particular models (Scott’s P_ω and Plotkin’s T_ω) and their results depend on the fact that these models have a very special (and well known) common theory.

A numeration of a set A is a map from \mathbb{N} onto A . $\mathcal{W} : \mathbb{N} \rightarrow \mathcal{RE}$ denotes the usual numeration of r.e. sets (i.e., \mathcal{W}_n is the domain of the n -th computable function ϕ_n).

2.1 Lambda calculus and lambda models

Λ and Λ^o are, respectively, the set of λ -terms and of closed λ -terms. Concerning specific λ -terms we set: $\mathbf{i} \equiv \lambda x.x$; $\mathbf{T} \equiv \lambda xy.x$; $\mathbf{F} \equiv \lambda xy.y$; $\Omega \equiv (\lambda x.xx)(\lambda x.xx)$.

A set X of λ -terms is *trivial* if either $X = \emptyset$ or $X = \Lambda$.

We denote $\alpha\beta$ -conversion by $\lambda\beta$. A λ -theory \mathcal{T} is a congruence on Λ (with respect to the operators of abstraction and application) which contains $\lambda\beta$. We write $M =_{\mathcal{T}} N$ for $(M, N) \in \mathcal{T}$. If \mathcal{T} is a λ -theory, then $[M]_{\mathcal{T}}$ denotes the set $\{N : N =_{\mathcal{T}} M\}$. A λ -theory \mathcal{T} is: *consistent* if $\mathcal{T} \neq \Lambda \times \Lambda$; *extensional* if it contains the equation $\mathbf{i} = \lambda xy.xy$; *recursively enumerable* if the set of Gödel numbers of all pairs of \mathcal{T} -equivalent λ -terms is r.e. Finally, $\lambda\beta\eta$ is the least extensional λ -theory.

The λ -theory \mathcal{H} , generated by equating all the unsolvable λ -terms, is consistent by [1, Theorem 16.1.3]. A λ -theory \mathcal{T} is *sensible* if $\mathcal{H} \subseteq \mathcal{T}$, while it is *semi-sensible* if it contains no equations of the form $U = S$ where S is solvable and U unsolvable. Consistent sensible theories are semi-sensible (see [1, Cor. 4.1.9]) and are never r.e. (see [1, Section 17.1]).

It is well known [1, Ch. 5] that a model of λ -calculus (λ -model, for short) can be defined as a reflexive object of a ccc (Cartesian closed category) \mathbf{C} , that is to say a triple $(D, \mathcal{F}, \lambda)$ such that D is an object of \mathbf{C} and $\mathcal{F} : D \rightarrow [D \rightarrow D]$, $\lambda : [D \rightarrow D] \rightarrow D$ are morphisms such that $\mathcal{F} \circ \lambda = id_{[D \rightarrow D]}$. In the following we will be interested in Scott's ccc of cpos (complete partial orderings) and Scott continuous functions (*continuous semantics*), but we will also draw conclusions for Berry's ccc of DI -domains and stable functions (*stable semantics*), and for Ehrhard's ccc of DI -domains with coherence and strongly stable functions between them (*strongly stable semantics*). We recall that DI -domains are special Scott domains, and that Scott domains are special cpos (see, e.g., [24]).

Let D be a cpo. The partial order of D will be denoted by \sqsubseteq_D . We let Env_D be the set of environments ρ mapping the set Var of variables of λ -calculus into D . For every $x \in Var$ and $d \in D$ we denote by $\rho[x := d]$ the environment ρ' which coincides with ρ , except on x , where ρ' takes the value d . A reflexive cpo D generates a λ -model $\mathcal{D} = (D, \mathcal{F}, \lambda)$ of λ -calculus with the interpretation of a λ -term defined as follows:

$$x_{\rho}^{\mathcal{D}} = \rho(x); (MN)_{\rho}^{\mathcal{D}} = \mathcal{F}(M_{\rho}^{\mathcal{D}})(N_{\rho}^{\mathcal{D}}); (\lambda x.M)_{\rho}^{\mathcal{D}} = \lambda(f),$$

where f is defined by $f(d) = M_{\rho[x:=d]}^{\mathcal{D}}$ for all $d \in D$. In the following $\mathcal{F}(d)(e)$ will also be written $d \cdot e$ or de .

Each λ -model \mathcal{D} induces a λ -theory, denoted here by $\text{Eq}(\mathcal{D})$, and called *the equational theory of \mathcal{D}* . Thus, $M = N \in \text{Eq}(\mathcal{D})$ if, and only if, M and N have the same interpretation in \mathcal{D} . A reflexive cpo D induces also an *order theory* $\text{Ord}(\mathcal{D}) = \{M \sqsubseteq N : M_{\rho}^{\mathcal{D}} \sqsubseteq_D N_{\rho}^{\mathcal{D}} \text{ for all environments } \rho\}$.

2.2 Effective domains

A triple $\mathcal{D} = (D, \sqsubseteq_D, d)$ is called an *effective domain* if (D, \sqsubseteq_D) is a Scott domain and d is a numeration of the set $K(\mathcal{D})$ of its compact elements such that the relations “ d_m and d_n have an upper bound” and “ $d_n = d_m \sqcup d_k$ ” are both decidable (see, e.g., [24, Ch. 10]).

We recall that an element v of an effective domain \mathcal{D} is said *r.e.* (*decidable*) if the set $\{n : d_n \sqsubseteq_D v\}$ is r.e. (decidable); we will write $\mathcal{D}^{r.e.}$ (\mathcal{D}^{dec}) for the set of r.e. (decidable) elements of \mathcal{D} . The set $K(\mathcal{D})$ of compact elements is included within \mathcal{D}^{dec} . Using standard techniques of recursion theory it is possible to get in a uniform way a numeration $\xi : \mathbb{N} \rightarrow \mathcal{D}^{r.e.}$ which is *adequate* in the sense that the relation $d_k \sqsubseteq_D \xi_n$ is r.e. in (k, n) and the inclusion mapping $\iota : K(\mathcal{D}) \rightarrow \mathcal{D}^{r.e.}$ is computable w.r.t. d, ξ .

The full subcategory **ED** of the category of Scott-domains with effective domains as objects and continuous functions as morphisms is a ccc.

A continuous function $f : D \rightarrow D'$ is an r.e. element in the effective domain of Scott continuous functions (i.e., $f \in [D \rightarrow D']^{r.e.}$) if, and only if, its restriction $f| : \mathcal{D}^{r.e.} \rightarrow \mathcal{D}'^{r.e.}$ is *computable w.r.t.* ξ, ξ' , i.e., there is a computable map $g : \mathbb{N} \rightarrow \mathbb{N}$ such that $f(\xi_n) = \xi'_{g(n)}$. We say that g *tracks* f .

2.3 Graph models

The class of graph models belongs to Scott continuous semantics (see [5] for a complete survey on this class of models). Historically, the first graph model was Scott’s P_ω , which is also known in the literature as “the graph model”. “Graph” referred to the fact that the continuous functions were encoded in the model via (a sufficient fragment of) their graph.

As a matter of notation, for every set G , G^* is the set of all finite subsets of G , while $\mathcal{P}(G)$ is the powerset of G .

Definition 1. A graph model \mathcal{G} is a pair $(G, c_{\mathcal{G}})$, where G is an infinite set, called the carrier set of \mathcal{G} , and $c_{\mathcal{G}} : G^* \times G \rightarrow G$ is an injective total function.

Such a pair \mathcal{G} generates the reflexive cpo $((\mathcal{P}(G), \subseteq), \lambda, \mathcal{F})$, where λ and \mathcal{F} are defined as follows, for all $f \in [\mathcal{P}(G) \rightarrow \mathcal{P}(G)]$ and $X, Y \subseteq G$: $\lambda(f) = \{c_{\mathcal{G}}(a, \alpha) : \alpha \in f(a) \text{ and } a \in G^*\}$ and $\mathcal{F}(X)(Y) = \{\alpha \in G : (\exists a \subseteq Y) c_{\mathcal{G}}(a, \alpha) \in X\}$. For more details we refer the reader to Berline [4].

The interpretation of a λ -term M into a λ -model has been defined in Section 2.1. However, in this context we can make explicit the interpretation $M_\rho^{\mathcal{G}}$ of a λ -term M as follows:

$$(MN)_\rho^{\mathcal{G}} = \{\alpha : (\exists a \subseteq N_\rho^{\mathcal{G}}) c_{\mathcal{G}}(a, \alpha) \in M_\rho^{\mathcal{G}}\}; (\lambda x.M)_\rho^{\mathcal{G}} = \{c_{\mathcal{G}}(a, \alpha) : \alpha \in M_{\rho[x:=a]}^{\mathcal{G}}\}.$$

We turn now to the interpretation of Ω in graph models (the details of the proof are, for example, worked out in [6, Lemma 4]).

Lemma 1. $\alpha \in \Omega^{\mathcal{G}}$ if, and only if, there is $a \subseteq (\lambda x.xx)^{\mathcal{G}}$ such that $c_{\mathcal{G}}(a, \alpha) \in a$.

In the following we use the terminology “*graph theory*” as a shorthand for “theory of a graph model”. It is well known that the equational graph theories are never extensional and that there exists a continuum of them (see [16]). In [9, 10] the existence of a minimum equational graph theory was proved and it was also shown that this minimum theory is different from $\lambda\beta$.

The completion method for building graph models from “partial pairs” was initiated by Longo in [18] and developed on a wide scale by Kerth in [16, 17].

Definition 2. A partial pair \mathcal{A} is given by a set A and by a partial, injective function $c_{\mathcal{A}} : A^* \times A \rightarrow A$.

A partial pair is *finite* if A is finite, and is a graph model if $c_{\mathcal{A}}$ is total.

The interpretation of a λ -term in a partial pair \mathcal{A} is defined in the obvious way: $(MN)_{\rho}^{\mathcal{A}} = \{ \alpha \in A : (\exists a \subseteq N_{\rho}^{\mathcal{A}}) [(a, \alpha) \in \text{dom}(c_{\mathcal{A}}) \wedge c_{\mathcal{A}}(a, \alpha) \in M_{\rho}^{\mathcal{A}}] \}$; $(\lambda x.M)_{\rho}^{\mathcal{A}} = \{ c_{\mathcal{A}}(a, \alpha) \in \mathcal{A} : (a, \alpha) \in \text{dom}(c_{\mathcal{A}}) \wedge \alpha \in M_{\rho[x:=a]}^{\mathcal{A}} \}$.

Definition 3. Let \mathcal{A} be a partial pair. The completion of \mathcal{A} is the graph model $\mathcal{E}_{\mathcal{A}} = (E_{\mathcal{A}}, c_{\mathcal{E}_{\mathcal{A}}})$ defined as follows:

- $E_{\mathcal{A}} = \bigcup_{n \in \mathbb{N}} E_n$, where $E_0 = A$ and $E_{n+1} = E_n \cup ((E_n^* \times E_n) - \text{dom}(c_{\mathcal{A}}))$.
- Given $a \in E_{\mathcal{A}}^*$, $\alpha \in E_{\mathcal{A}}$,

$$c_{\mathcal{E}_{\mathcal{A}}}(a, \alpha) = \begin{cases} c_{\mathcal{A}}(a, \alpha) & \text{if } c_{\mathcal{A}}(a, \alpha) \text{ is defined} \\ (a, \alpha) & \text{otherwise} \end{cases}$$

A notion of *rank* can be naturally defined on the completion $\mathcal{E}_{\mathcal{A}}$ of a partial pair \mathcal{A} . The elements of A are the elements of rank 0, while an element $\alpha \in E_{\mathcal{A}} - A$ has rank n if $\alpha \in E_n$ and $\alpha \notin E_{n-1}$.

Let \mathcal{A} and \mathcal{B} be two partial pairs. A *morphism* from \mathcal{A} into \mathcal{B} is a map $f : A \rightarrow B$ such that $(a, \alpha) \in \text{dom}(c_{\mathcal{A}})$ implies $(fa, f\alpha) \in \text{dom}(c_{\mathcal{B}})$ and, in such a case $f(c_{\mathcal{A}}(a, \alpha)) = c_{\mathcal{B}}(fa, f\alpha)$. Isomorphisms and automorphisms can be defined in the obvious way. $\text{Aut}(\mathcal{A})$ denotes the group of automorphisms of the partial pair \mathcal{A} .

Lemma 2. Let $\mathcal{G}, \mathcal{G}'$ be graph models and $f : \mathcal{G} \rightarrow \mathcal{G}'$ be a morphism. If $M \in \mathcal{A}$ and $\alpha \in M_{\rho}^{\mathcal{G}}$, then $f\alpha \in M_{f\rho}^{\mathcal{G}'}$.

3 Co-r.e. sets of lambda terms

In this section we recall the main properties of recursion theory concerning λ -calculus that will be applied in the following sections.

An r.e. (co-r.e.) set of λ -terms closed under β -conversion will be called a β -r.e. (β -co-r.e.) set.

The following theorem is due to Scott (see [1, Thm. 6.6.2]).

Theorem 1. A set of λ -terms which is both β -r.e. and β -co-r.e. is trivial.

Definition 4. A family $X = (X_i : i \in I)$ of sets has the FIP (finite intersection property) if $X_{i_1} \cap \dots \cap X_{i_n} \neq \emptyset$ for all $i_1, \dots, i_n \in I$.

Visser (see [1, Ch. 17] and [25, Thm. 2.5]) has shown that the topology on Λ generated by the β -co-r.e. sets of λ -terms is hyperconnected (i.e., the intersection of two non-empty open sets is non-empty). In other words:

Theorem 2. The family of all non-empty β -co-r.e. subsets of Λ has the FIP.

Remark 1. The set of all unsolvable λ -terms is β -co-r.e. It follows from Theorem 2 that every non-empty β -co-r.e. set of terms contains unsolvable λ -terms.

We conclude this section by providing a new proof, based on Theorem 2, of the genericity lemma of λ -calculus. We recall that a proof of the genericity lemma, due to Barendregt (see [1, Thm. 14.3.24]), is obtained by using the tree topology on Λ which is induced by the Scott topology on the set of Böhm trees.

Lemma 3. (*Genericity Lemma*) Let $U, N \in \Lambda$, where U is unsolvable and N is β -normal. Then for all contexts $C[-]$

$$C[U] =_{\lambda\beta} N \Rightarrow \forall M \in \Lambda \quad C[M] =_{\lambda\beta} N.$$

Proof. The set $\{M : M =_{\lambda\beta} N\}$ is β -r.e. As the map defined by $M \mapsto C[M]$ is computable, the set $O = \{M : C[M] =_{\lambda\beta} N\}$ is a β -r.e. set, containing the unsolvable U . Since N is normal $[N]_{\mathcal{H}} = [N]_{\lambda\beta}$ [1, Theorem 16.1.9], where \mathcal{H} is the least sensible λ -theory. Hence the β -r.e. set O is a union of \mathcal{H} -equivalence classes, containing the set \mathcal{U} of all the unsolvable λ -terms. Since $\Lambda - O$ and \mathcal{U} are both β -co-r.e. and $(\Lambda - O) \cap \mathcal{U} = \emptyset$, then the FIP implies that $O = \Lambda$.

4 Effective lambda models

In this section we introduce the notion of an effective λ -model and we study the main properties of these models. We show that the order theory of an effective λ -model is not r.e. and that its equational theory is different from $\lambda\beta, \lambda\beta\eta$. Effective λ -models are omni-present in the continuous, stable and strongly stable semantics (see Section 4.1). In particular, all the λ -models which have been introduced individually in the literature, to begin with Scott's \mathcal{D}_∞ , can easily be proved effective.

The following natural definition is enough to force the interpretation function of λ -terms to be computable from A^o into $\mathcal{D}^{r.e.}$. However, other results of this paper will need a more powerful notion. That is the reason why we only speak of “weak effectivity” here.

Definition 5. A λ -model is called weakly effective if it is a reflexive object $(\mathcal{D}, \mathcal{F}, \lambda)$ in the category **ED** and, $\mathcal{F} \in [\mathcal{D} \rightarrow [\mathcal{D} \rightarrow \mathcal{D}]]$ and $\lambda \in [[\mathcal{D} \rightarrow \mathcal{D}] \rightarrow \mathcal{D}]$ are r.e. elements.

In the following a weakly effective λ -model $(\mathcal{D}, \mathcal{F}, \lambda)$ will be denoted by \mathcal{D} .

We fix bijective effective numerations $\nu_A : \mathbb{N} \rightarrow A$ of the set of λ -terms and $\nu_{var} : \mathbb{N} \rightarrow Var$ of the set of variables of λ -calculus. In particular this gives to the set Env_D of all environments a structure of effective domain. $A_\perp = A \cup \{\perp\}$ is the usual flat domain of λ -terms. The element \perp is always interpreted as \perp_D in a cpo (D, \sqsubseteq_D) .

Proposition 1. *Let \mathcal{D} be a weakly effective λ -model. Then the function f mapping $(\rho, M) \mapsto M_\rho^{\mathcal{D}}$ is an element of $[Env_D \times A_\perp \rightarrow \mathcal{D}]^{r.e.}$.*

Proof. (Sketch) By structural induction on M it is possible to show the existence of a partial computable map tracking f . The only difficult case is $M \equiv \lambda x.N$. Since λ is r.e. it is sufficient to prove that the function $g : e \mapsto N_{\rho[x:=e]}^{\mathcal{D}}$ is also r.e. Once shown that $h : (\rho, x, e) \mapsto \rho[x := e]$ is r.e., from the induction hypothesis it follows that the function $g'(\rho, x, e) = f(h(\rho, x, e), N)$ is r.e. Then by applying the s-m-n theorem of recursion theory to the computable function tracking g' we obtain a computable function tracking g , which is then r.e.

Notation 1. *We define for any $e \in D$ and $M \in A^\circ$:*

- (i) $e^- \equiv \{P \in A^\circ : P^{\mathcal{D}} \sqsubseteq_D e\}$;
- (ii) $M^- \equiv \{P \in A^\circ : P^{\mathcal{D}} \sqsubseteq_D M^{\mathcal{D}}\}$.

Corollary 1. *If $e \in \mathcal{D}^{dec}$, then e^- is a β -co-r.e. set of λ -terms.*

Proof. Let $\rho \in (Env_D)^{r.e.}$ be an environment. By Proposition 1 there is a computable map ϕ tracking the interpretation function $M \mapsto M_\rho^{\mathcal{D}}$ of λ -terms from A into $\mathcal{D}^{r.e.}$ with respect to the effective numeration ν_A of A and an adequate numeration ξ of $\mathcal{D}^{r.e.}$. From $e \in \mathcal{D}^{dec}$ it follows that the set $X = \{n : \xi_n \sqsubseteq_{\mathcal{D}} e\}$ is co-r.e. This implies that the set $\phi^{-1}(X)$, which is the set of the codes of the elements of $\{M \in A : M_\rho^{\mathcal{D}} \sqsubseteq_{\mathcal{D}} e\}$, is also co-r.e. We get the conclusion because A° is a decidable subset of A .

Definition 6. *A weakly effective λ -model \mathcal{D} is called effective if satisfies the following two further conditions:*

- (i) *If $d \in K(\mathcal{D})$ and $e_i \in \mathcal{D}^{dec}$, then $de_1 \dots e_n \in \mathcal{D}^{dec}$.*
- (ii) *If $f \in [\mathcal{D} \rightarrow \mathcal{D}]^{r.e.}$ and $f(e) \in \mathcal{D}^{dec}$ for all $e \in K(\mathcal{D})$, then $\lambda(f) \in \mathcal{D}^{dec}$.*

An environment ρ is *compact* in the effective domain Env_D (i.e., $\rho \in K(Env_D)$) if $\rho(x) \in K(\mathcal{D})$ for all variables x and $\{x : \rho(x) \neq \perp_D\}$ is finite.

Notation 2. *We define: $\Lambda_{\mathcal{D}}^{dec} \equiv \{M \in A : M_\rho^{\mathcal{D}} \in \mathcal{D}^{dec} \text{ for all } \rho \in K(Env_D)\}$.*

Theorem 3. *Suppose \mathcal{D} is an effective λ -model. Then the set $\Lambda_{\mathcal{D}}^{dec}$ is closed under the following rules:*

1. $x \in \Lambda_{\mathcal{D}}^{dec}$ for every variable x .
2. $M_1, \dots, M_k \in \Lambda_{\mathcal{D}}^{dec} \Rightarrow yM_1 \dots M_k \in \Lambda_{\mathcal{D}}^{dec}$.
3. $M \in \Lambda_{\mathcal{D}}^{dec} \Rightarrow \lambda x.M \in \Lambda_{\mathcal{D}}^{dec}$.

In particular, $\Lambda_{\mathcal{D}}^{dec}$ contains all the β -normal forms.

Proof. Let $\rho \in K(Env_{\mathcal{D}})$. We have three cases.

- (1) $x_{\rho}^{\mathcal{D}} = \rho(x)$ is compact, hence it is decidable.
- (2) By definition $(yM_1 \dots M_k)_{\rho}^{\mathcal{D}} = \rho(y)(M_1)_{\rho}^{\mathcal{D}} \dots (M_k)_{\rho}^{\mathcal{D}}$. Hence the result follows from Definition 6(i), $\rho(y) \in K(\mathcal{D})$ and $(M_i)_{\rho}^{\mathcal{D}} \in \mathcal{D}^{dec}$.
- (3) By definition we have that $(\lambda x.M)_{\rho}^{\mathcal{D}} = \lambda(f)$, where $f(e) = M_{\rho[x:=e]}^{\mathcal{D}}$ for all $e \in D$. Note that $\rho[x:=e]$ is also compact for all $e \in K(D)$. Hence the conclusion follows from $M_{\rho[x:=e]}^{\mathcal{D}} \in \mathcal{D}^{dec}$ ($e \in K(\mathcal{D})$), Definition 6(ii) and $f \in [\mathcal{D} \rightarrow \mathcal{D}]^{r.e.}$.

Recall that $Eq(\mathcal{D})$ and $Ord(\mathcal{D})$ are respectively the equational theory and the order theory of \mathcal{D} .

Theorem 4. *Let \mathcal{D} be an effective λ -model, and let $M_1, \dots, M_k \in \Lambda_{\mathcal{D}}^{dec}$ ($k \geq 1$) be closed terms. Then we have:*

- (i) $M_1^- \cap \dots \cap M_k^-$ is a β -co-r.e. set, which contains a non-empty β -co-r.e. set of unsolvable terms.
- (ii) If $e \in \mathcal{D}^{dec}$ and e^- is non-empty and finite modulo $Eq(\mathcal{D})$, then $Eq(\mathcal{D})$ is not r.e. (in particular, if $\perp_{\mathcal{D}}^- \neq \emptyset$ then $Eq(\mathcal{D})$ is not r.e.).
- (iii) $Ord(\mathcal{D})$ is not r.e.
- (iv) $Eq(\mathcal{D}) \neq \lambda\beta, \lambda\beta\eta$.

Proof. (i) By Theorem 3, Corollary 1, and the FIP.

(ii) By Corollary 1.

(iii) Let $M \in \Lambda_{\mathcal{D}}^{dec}$ be a closed term. If $Ord(\mathcal{D})$ were r.e., then we could enumerate the set M^- . However, by (i) this set is non-empty and β -co-r.e. By Theorem 1 it follows that $M^- = A^{\circ}$. By the arbitrariness of M , it follows that $T^- = F^-$. Since $F \in T^-$ and conversely we get $F = T$ in \mathcal{D} , contradiction.

(iv) Because of (iii), if $Eq(\mathcal{D})$ is r.e. then $Ord(\mathcal{D})$ strictly contains $Eq(\mathcal{D})$. Hence the conclusion follows from Selinger's result stating that in any partially ordered λ -model, whose theory is $\lambda\beta$, the interpretations of distinct closed terms are incomparable [23, Corollary 4]. Similarly for $\lambda\beta\eta$.

4.1 Can effective λ -models have an r.e. theory?

In this section we give a sufficient condition for a wide class of graph models to be effective and show that no effective graph model generated freely by a partial pair, which is finite modulo its group of automorphisms, can have an r.e. equational theory. Finally, we show that no effective λ -model living in the stable or strongly stable semantics can have an r.e. equational theory.

In Section 5 we will show that every equational/order graph theory is the theory of a graph model \mathcal{G} whose carrier set is the set \mathbb{N} of natural numbers. In the next theorem we characterize the effectivity of these models.

Theorem 5. *Let \mathcal{G} be a graph model such that, after encoding, $G = \mathbb{N}$ and $c_{\mathcal{G}}$ is a computable map. Then \mathcal{G} is weakly effective. Moreover, \mathcal{G} is effective under the further hypothesis that $c_{\mathcal{G}}$ has a decidable range.*

Proof. It is easy to check, using the definitions given in Section 2.3, that \mathcal{F}, λ are r.e. in their respective domains and that condition (i) of Definition 6 is satisfied. Then \mathcal{G} is weakly effective. Moreover, Definition 6(ii) holds under the hypothesis that the range of $c_{\mathcal{G}}$ is decidable.

Completions of partial pairs have been extensively studied in literature. They are useful for solving equational and inequational constraints (see [4, 5, 10, 11]). In [11] Bucciarelli and Salibra have recently proved that the theory of the completion of a partial pair which is not a graph model is semi-sensible. The following theorem shows, in particular, that the theory of the completion of a finite partial pair is not r.e.

Theorem 6. *Let \mathcal{A} be a partial pair such that A is finite or equal to \mathbb{N} after encoding, and $c_{\mathcal{A}}$ is a computable map with a decidable domain. Then we have:*

- (i) *The completion $\mathcal{E}_{\mathcal{A}}$ of \mathcal{A} is weakly effective;*
- (ii) *If the range of $c_{\mathcal{A}}$ is decidable, then $\mathcal{E}_{\mathcal{A}}$ is effective;*
- (iii) *If A is finite modulo its group of automorphisms (in particular, if A is finite), then $\text{Eq}(\mathcal{E}_{\mathcal{A}})$ is not r.e.*

Proof. Since A is finite or equal to \mathbb{N} we have that $E_{\mathcal{A}}$ is also decidable (see Definition 3). Moreover, the map $c_{\mathcal{E}_{\mathcal{A}}} : E_{\mathcal{A}}^* \times E_{\mathcal{A}} \rightarrow E_{\mathcal{A}}$ is computable, because it is an extension of a computable function $c_{\mathcal{A}}$ with decidable domain, and it is the identity on the decidable set $(E_{\mathcal{A}}^* \times E_{\mathcal{A}}) - \text{dom}(c_{\mathcal{A}})$. Then (i)-(ii) follow from Theorem 5.

Clearly A is a decidable subset of $E_{\mathcal{A}}$; then by Corollary 1 the set A^- is a β -co-r.e. set of λ -terms. We now show that this set is non-empty because $\Omega^{\mathcal{E}_{\mathcal{A}}} \subseteq A$. By Lemma 1 we have that $\alpha \in \Omega^{\mathcal{E}_{\mathcal{A}}}$ implies that $c_{\mathcal{E}_{\mathcal{A}}}(a, \alpha) \in a$ for some $a \in E_{\mathcal{A}}^*$. Immediate considerations on the rank show that this is only possible if $(a, \alpha) \in \text{dom}(c_{\mathcal{A}})$, which forces $\alpha \in A$.

The orbit of $\alpha \in A$ modulo $\text{Aut}(\mathcal{A})$ is defined by $O(\alpha) = \{\theta(\alpha) : \theta \in \text{Aut}(\mathcal{A})\}$.

We now show that, if the set of orbits of \mathcal{A} has cardinality k for some $k \in \mathbb{N}$, then the cardinality of A^- modulo $\text{Eq}(\mathcal{E}_{\mathcal{A}})$ is less than or equal to 2^k . Assume $\alpha \in M^{\mathcal{E}_{\mathcal{A}}} \subseteq A$. Then by Lemma 2 the orbit of α modulo $\text{Aut}(\mathcal{A})$ is included within $M^{\mathcal{E}_{\mathcal{A}}}$. By hypothesis the number of the orbits is k ; hence, the number of all possible values for $M^{\mathcal{E}_{\mathcal{A}}}$ cannot overcome 2^k .

In conclusion, A^- is non-empty, β -co-r.e. and modulo $\text{Eq}(\mathcal{E}_{\mathcal{A}})$ is finite. Then (iii) follows from Theorem 4.

All the material developed in Section 4 could be adapted to the stable semantics (Berry's ccc of *DI*-domains and stable functions) and strongly stable semantics (Ehrhard's ccc of *DI*-domains with coherence and strongly stable functions). We recall that the notion of an effectively given *DI*-domain has been introduced by Gruchalski in [14], where it is shown that the category having effective *DI*-domains as objects and stable functions as morphisms is a ccc. There are also many effective models in the stable and strongly stable semantics. Indeed, the stable semantics contains a class which is analogous to the class of

graph models (see [4]), namely Girard's class of *reflexive coherent spaces* called *G-models* in [4]. The results shown in Theorem 5 and in Theorem 6 for graph models could also be adapted for *G-models*, even if it is more delicate to complete partial pairs in this case (see [17]). It could also be developed for Ehrhard's class of strongly stable *H-models* (see [4]) even though working in the strongly stable semantics certainly adds technical difficulties.

Theorem 7. *Let \mathcal{D} be an effective λ -model in the stable or strongly stable semantics. Then $\text{Eq}(\mathcal{D})$ is not r.e.*

Proof. By Theorem 4(ii) it is enough to prove that $\perp_{\mathcal{D}} \neq \emptyset$. Since \mathcal{D} is effective, then by Theorem 4(i) $F^- \cap T^-$ is a non-empty and co-r.e. set of λ -terms. Let $N \in F^- \cap T^-$ and let $f, g, h : \mathcal{D} \rightarrow \mathcal{D}$ be three (strongly) stable functions such that $f(x) = T^{\mathcal{D}} \cdot x$, $g(x) = F^{\mathcal{D}} \cdot x$ and $h(x) = N^{\mathcal{D}} \cdot x$ for all $x \in \mathcal{D}$. By monotonicity we have $h \leq f, g$ in the stable ordering. Now, g is the constant function taking value $\mathbf{i}^{\mathcal{D}}$, and $f(\perp_{\mathcal{D}}) = T^{\mathcal{D}} \cdot \perp_{\mathcal{D}}$. The first assertion forces h to be a constant function, because in the stable ordering all functions under a constant map are also constant, while the second assertion together with the fact that h is pointwise smaller than f forces the constant function h to satisfy $h(x) = T^{\mathcal{D}} \cdot \perp_{\mathcal{D}}$ for all x . Then an easy computation gives $(NPP)^{\mathcal{D}} = \perp_{\mathcal{D}}$ for every closed term P . In conclusion, we have $\{M \in \Lambda^{\circ} : M^{\mathcal{D}} = \perp_{\mathcal{D}}\} \neq \emptyset$ and $\text{Eq}(\mathcal{D})$ is not r.e.

5 The Löwenheim-Skolem theorem

In this section we show that for each graph model \mathcal{G} there is a countable graph model \mathcal{P} with the same equational/order theory. This result is a kind of downwards Löwenheim-Skolem theorem for graph models which positively answers Question 3 in [4, Section 6.3]. Note that we cannot apply directly the classical Löwenheim-Skolem theorem since graph models are not first-order structures.

Let \mathcal{A}, \mathcal{B} be partial pairs. We say that \mathcal{A} is a *subpair* of \mathcal{B} , and we write $\mathcal{A} \leq \mathcal{B}$, if $A \subseteq B$ and $c_{\mathcal{B}}(a, \alpha) = c_{\mathcal{A}}(a, \alpha)$ for all $(a, \alpha) \in \text{dom}(c_{\mathcal{A}})$.

As a matter of notation, if ρ, σ are environments and C is a set, we let $\sigma = \rho \cap C$ mean $\sigma(x) = \rho(x) \cap C$ for every variable x , and $\rho \subseteq \sigma$ mean $\rho(x) \subseteq \sigma(x)$ for every variable x .

The proof of the following lemma is straightforward. Recall that the definition of interpretation with respect to a partial pair is defined in Section 2.3.

Lemma 4. *Suppose $\mathcal{A} \leq \mathcal{B}$, then $M_{\rho}^{\mathcal{A}} \subseteq M_{\sigma}^{\mathcal{B}}$ for all environments $\rho : \text{Var} \rightarrow \mathcal{P}(A)$ and $\sigma : \text{Var} \rightarrow \mathcal{P}(B)$ such that $\rho \subseteq \sigma$.*

Lemma 5. *Let M be a λ -term, \mathcal{G} be a graph model and $\alpha \in M_{\rho}^{\mathcal{G}}$ for some environment ρ . Then there exists a finite subpair \mathcal{A} of \mathcal{G} such that $\alpha \in M_{\rho \cap A}^{\mathcal{A}}$.*

Proof. The proof is by induction on M .

If $M \equiv x$, then $\alpha \in \rho(x)$, so that we define $A = \{\alpha\}$ and $\text{dom}(c_{\mathcal{A}}) = \emptyset$.

If $M \equiv \lambda x.P$, then $\alpha \equiv c_{\mathcal{G}}(b, \beta)$ for some b and β such that $\beta \in P_{\rho[x:=b]}^{\mathcal{G}}$. By induction hypothesis there exists a finite subpair \mathcal{B} of \mathcal{G} such that $\beta \in P_{\rho[x:=b] \cap B}^{\mathcal{B}}$. We define another finite subpair \mathcal{A} of \mathcal{G} as follows: $A = B \cup b \cup \{\beta, \alpha\}$; $\text{dom}(c_{\mathcal{A}}) = \text{dom}(c_{\mathcal{B}}) \cup \{(b, \beta)\}$. Then we have that $\mathcal{B} \leq \mathcal{A}$ and $\rho[x := b] \cap B \subseteq \rho[x := b] \cap A$. From $\beta \in P_{\rho[x:=b] \cap B}^{\mathcal{B}}$ and from Lemma 4 it follows that $\beta \in P_{\rho[x:=b] \cap A}^{\mathcal{A}} = P_{(\rho \cap A)[x:=b]}^{\mathcal{A}}$. Then we have that $\alpha \equiv c_{\mathcal{A}}(b, \beta) \in (\lambda x.P)_{\rho \cap A}^{\mathcal{A}}$.

If $M \equiv PQ$, then there is $a = \{\alpha_1, \dots, \alpha_n\}$ such that $c_{\mathcal{G}}(a, \alpha) \in P_{\rho}^{\mathcal{G}}$ and $a \subseteq Q_{\rho}^{\mathcal{G}}$. By induction hypothesis there exist finite subpairs $\mathcal{A}_0, \mathcal{A}_1, \dots, \mathcal{A}_n$ of \mathcal{G} such that $c_{\mathcal{G}}(a, \alpha) \in P_{\rho \cap A_0}^{\mathcal{A}_0}$ and $\alpha_k \in Q_{\rho \cap A_k}^{\mathcal{A}_k}$ for $k = 1, \dots, n$. We define another finite subpair \mathcal{A} of \mathcal{G} as follows: $A = \bigcup_{0 \leq k \leq n} A_k \cup a \cup \{\alpha\}$ and $\text{dom}(c_{\mathcal{A}}) = (\bigcup_{0 \leq k \leq n} \text{dom}(c_{\mathcal{A}_k})) \cup \{(a, \alpha)\}$. From Lemma 4 it follows the conclusion.

Proposition 2. *Let \mathcal{G} be a graph model, and suppose $\alpha \in M^{\mathcal{G}} - N^{\mathcal{G}}$ for some $M, N \in \Lambda^{\circ}$. Then there exists a finite $\mathcal{A} \leq \mathcal{G}$ such that: for all pairs $\mathcal{C} \geq \mathcal{A}$, if there is a morphism $f : \mathcal{C} \rightarrow \mathcal{G}$ such that $f(\alpha) = \alpha$, then $\alpha \in M^{\mathcal{C}} - N^{\mathcal{C}}$.*

Proof. By Lemma 5 there is a finite pair \mathcal{A} such that $\alpha \in M^{\mathcal{A}}$. By Lemma 4 we have $\alpha \in M^{\mathcal{C}}$. Now, if $\alpha \in N^{\mathcal{C}}$ then, by Lemma 2 $\alpha = f(\alpha) \in N^{\mathcal{G}}$, which is a contradiction.

Corollary 2. *Let \mathcal{G} be a graph model, and suppose $\alpha \in M^{\mathcal{G}} - N^{\mathcal{G}}$ for some $M, N \in \Lambda^{\circ}$. Then there exists a finite $\mathcal{A} \leq \mathcal{G}$ such that: for all pairs \mathcal{B} satisfying $\mathcal{A} \leq \mathcal{B} \leq \mathcal{G}$ we have $\alpha \in M^{\mathcal{B}} - N^{\mathcal{B}}$.*

Let \mathcal{G} be a graph model. A graph model \mathcal{P} is called a *sub graph model* of \mathcal{G} if $\mathcal{P} \leq \mathcal{G}$. It is easy to check that the class of sub graph models of \mathcal{G} is closed under (finite and infinite) intersection. If $\mathcal{A} \leq \mathcal{G}$ is a partial pair, then the *sub graph model generated by \mathcal{A}* is defined as the intersection of all graph models \mathcal{P} such that $\mathcal{A} \leq \mathcal{P} \leq \mathcal{G}$.

Theorem 8. (Löwenheim-Skolem Theorem for graph models) *For every graph model \mathcal{G} there exists a sub graph model \mathcal{P} of \mathcal{G} with a countable carrier set and such that $\text{Ord}(\mathcal{P}) = \text{Ord}(\mathcal{G})$, and hence $\text{Eq}(\mathcal{P}) = \text{Eq}(\mathcal{G})$.*

Proof. We will define an increasing sequence of countable subpairs \mathcal{A}_n of \mathcal{G} , and take for \mathcal{P} the sub graph model of \mathcal{G} generated by $\mathcal{A} \equiv \bigcup \mathcal{A}_n$.

First we define \mathcal{A}_0 . Let I be the countable set of inequations between closed λ -terms which fail in \mathcal{G} . Let $e \in I$. By Corollary 2 there exists a finite partial pair $\mathcal{A}_e \leq \mathcal{G}$ such that e fails in every partial pair \mathcal{B} satisfying $\mathcal{A}_e \leq \mathcal{B} \leq \mathcal{G}$. Then we define $\mathcal{A}_0 = \bigcup_{e \in I} \mathcal{A}_e \leq \mathcal{G}$. Assume now that \mathcal{A}_n has been defined. We define \mathcal{A}_{n+1} as follows. For each inequation $e \equiv M \sqsubseteq N$ which holds in \mathcal{G} and fails in the sub graph model $\mathcal{P}_n \leq \mathcal{G}$ generated by \mathcal{A}_n , we consider the set $L_e = \{\alpha \in P_n : \alpha \in M^{\mathcal{P}_n} - N^{\mathcal{P}_n}\}$. Let $\alpha \in L_e$. Since $\mathcal{P}_n \leq \mathcal{G}$ and $\alpha \in M^{\mathcal{P}_n}$, then by Lemma 4 we have that $\alpha \in M^{\mathcal{G}}$. By $\mathcal{G} \models M \sqsubseteq N$ we also obtain $\alpha \in N^{\mathcal{G}}$. By Lemma 5 there exists a partial pair $\mathcal{F}_{\alpha, e} \leq \mathcal{G}$ such that $\alpha \in N^{\mathcal{F}_{\alpha, e}}$. We define

\mathcal{A}_{n+1} as the union of the partial pair \mathcal{A}_n and the partial pairs $\mathcal{F}_{\alpha,e}$ for every $\alpha \in L_e$.

Finally take for \mathcal{P} the sub graph model of \mathcal{G} generated by $\mathcal{A} \equiv \cup \mathcal{A}_n$. By construction we have, for every inequation e which fails in \mathcal{G} : $\mathcal{A}_e \leq \mathcal{P}_n \leq \mathcal{P} \leq \mathcal{G}$. Now, $\text{Ord}(\mathcal{P}) \subseteq \text{Ord}(\mathcal{G})$ follows from Corollary 2 and from the choice of \mathcal{A}_e .

Let now $M \sqsubseteq N$ be an inequation which fails in \mathcal{P} but not in \mathcal{G} . Then there is an $\alpha \in M^{\mathcal{P}} - N^{\mathcal{P}}$. By Corollary 2 there is a finite partial pair $\mathcal{B} \leq \mathcal{P}$ satisfying the following condition: for every partial pair \mathcal{C} such that $\mathcal{B} \leq \mathcal{C} \leq \mathcal{P}$, we have $\alpha \in M^{\mathcal{C}} - N^{\mathcal{C}}$. Since B is finite, we have that $\mathcal{B} \leq \mathcal{P}_n$ for some n . This implies that $\alpha \in M^{\mathcal{P}_n} - N^{\mathcal{P}_n}$. By construction of \mathcal{P}_{n+1} we have that $\alpha \in N^{\mathcal{P}_{n+1}}$; this implies $\alpha \in N^{\mathcal{P}}$. Contradiction.

6 The minimum order graph theory

In this section we show one of the main theorems of the paper: the minimum order graph theory exists and it is the theory of an effective graph model. This result has the interesting consequence that no order graph theory can be r.e.

Lemma 6. *Suppose $\mathcal{A} \leq \mathcal{G}$ and let $f : E_{\mathcal{A}} \rightarrow G$ be defined by induction over the rank of $x \in E_{\mathcal{A}}$ as follows:*

$$f(x) = \begin{cases} x & \text{if } x \in A \\ c_{\mathcal{G}}(fa, f\alpha) & \text{if } x \notin A \text{ and } x \equiv (a, \alpha). \end{cases}$$

Then f is a morphism from $\mathcal{E}_{\mathcal{A}}$ into \mathcal{G} .

Lemma 7. *Suppose $\alpha \in M^{\mathcal{G}} - N^{\mathcal{G}}$ for some $M, N \in \Lambda^o$. Then there exists a finite $\mathcal{A} \leq \mathcal{G}$ such that: for all pairs \mathcal{B} satisfying $\mathcal{A} \leq \mathcal{B} \leq \mathcal{G}$, we have $\alpha \in M^{\mathcal{E}_{\mathcal{B}}} - N^{\mathcal{E}_{\mathcal{B}}}$.*

Proof. By Proposition 2 and Lemma 6.

Theorem 9. *There exists an effective graph model whose order/equational theory is the minimum order/equational graph theory.*

Proof. It is not difficult to define an effective bijective numeration \mathcal{N} of all the finite partial pairs whose carrier set is a subset of \mathbb{N} . We denote by \mathcal{N}_k the k -th finite partial pair with $N_k \subseteq \mathbb{N}$. We now make the carrier sets N_k ($k \in \mathbb{N}$) disjoint. Let p_k be the k -th prime natural number. Then we define another finite partial pair \mathcal{P}_k as follows: $P_k = \{p_k^{x+1} : x \in N_k\}$ and $c_{\mathcal{P}_k}(\{p_k^{\alpha_1+1}, \dots, p_k^{\alpha_n+1}\}, p_k^{\alpha+1}) = p_k^{c_{\mathcal{N}_k}(\{\alpha_1, \dots, \alpha_n\}, \alpha)+1}$ for all $(\{\alpha_1, \dots, \alpha_n\}, \alpha) \in \text{dom}(c_{\mathcal{N}_k})$. In this way we get an effective bijective numeration of all the finite partial pairs \mathcal{P}_k . Finally, we take $\mathcal{P} \equiv \cup_{k \in \mathbb{N}} \mathcal{P}_k$. It is an easy matter to prove that \mathcal{P} is a decidable subset of \mathbb{N} and that, after encoding, $c_{\mathcal{P}} = \cup_{k \in \mathbb{N}} c_{\mathcal{P}_k}$ is a computable map with a decidable domain and range. Then by Theorem 6(ii) $\mathcal{E}_{\mathcal{P}}$ is an effective graph model. Notice that $\mathcal{E}_{\mathcal{P}}$ is also isomorphic to the completion of the union $\cup_{k \in \mathbb{N}} \mathcal{E}_{\mathcal{P}_k}$, where $\mathcal{E}_{\mathcal{P}_k}$ is the completion of the partial pair \mathcal{P}_k .

We now prove that the order theory of $\mathcal{E}_{\mathcal{P}}$ is the minimum one. Let $e \equiv M \sqsubseteq N$ be an inequation which fails in some graph model \mathcal{G} . By Lemma 7 e fails in the completion of a finite partial pair \mathcal{A} . Without loss of generality, we may assume that the carrier set of \mathcal{A} is a subset of \mathbb{N} , and then that \mathcal{A} is one of the partial pairs \mathcal{P}_k . For such a \mathcal{P}_k , e fails in $\mathcal{E}_{\mathcal{P}_k}$. Now, it was shown by Bucciarelli and Salibra in [9, Proposition 2] that, if a graph model \mathcal{G} is the completion of the disjoint union of a family of graph models \mathcal{G}_i , then $Q^{\mathcal{G}_i} = Q^{\mathcal{G}} \cap G_i$ for any closed λ -term Q . Then we can conclude the proof as follows: if the inequation e holds in $\mathcal{E}_{\mathcal{P}}$, then by [9, Proposition 2] we get a contradiction: $M^{\mathcal{E}_{\mathcal{P}_k}} = M^{\mathcal{E}_{\mathcal{P}}} \cap E_{\mathcal{P}_k} \subseteq N^{\mathcal{E}_{\mathcal{P}}} \cap E_{\mathcal{P}_k} = N^{\mathcal{E}_{\mathcal{P}_k}}$.

Theorem 10. *Let \mathcal{T}_{min} and \mathcal{O}_{min} be, respectively, the minimum equational graph theory and the minimum order graph theory. We have:*

- (i) \mathcal{O}_{min} is not r.e.
- (ii) \mathcal{T}_{min} is an intersection of a countable set of non-r.e. equational graph theories.

Proof. (i) follows from Theorem 9 and from Theorem 4(iii), because \mathcal{O}_{min} is the theory of an effective λ -model.

(ii) By the proof of Theorem 9 we have that \mathcal{T}_{min} is an intersection of a countable set of graph theories, which are theories of completions of finite partial pairs. By Theorem 6(iii) these theories are not r.e.

Corollary 3. *For all graph models \mathcal{G} , $Ord(\mathcal{G})$ is not r.e.*

Proof. If $Ord(\mathcal{G})$ is r.e. and M is closed and β -normal, then $M^- = \{N \in \Lambda^o : N^{\mathcal{G}} \subseteq M^{\mathcal{G}}\}$ is a β -r.e. set, which contains the β -co-r.e. set $\{N \in \Lambda^o : \mathcal{O}_{min} \vdash N \sqsubseteq M\}$. By the FIP $M^- = \Lambda^o$. By the arbitrariness of M , it follows that $T^- = F^-$. Since $F \in T^-$ and conversely we get $F = T$ in \mathcal{G} , contradiction.

Corollary 4. *Let \mathfrak{G} be the class of all graph models. For any finite sequence M_1, \dots, M_n of closed β -normal forms, there exists a non-empty β -closed co-r.e. set \mathcal{U} of closed unsolvable terms such that*

$$(\forall \mathcal{G} \in \mathfrak{G})(\forall U \in \mathcal{U}) U^{\mathcal{G}} \subseteq M_1^{\mathcal{G}} \cap \dots \cap M_n^{\mathcal{G}}.$$

Proof. By Theorem 4(i) applied to any effective graph model with minimum theory, we have $(\forall U \in \mathcal{U}) \mathcal{O}_{min} \vdash U \sqsubseteq M_1 \wedge \dots \wedge \mathcal{O}_{min} \vdash U \sqsubseteq M_n$. The conclusion follows.

References

1. H.P. Barendregt. *The lambda calculus: Its syntax and semantics*. North-Holland Publishing Co., Amsterdam, 1984.
2. O. Bastonero, X. Gouy. Strong stability and the incompleteness of stable models of λ -calculus. *Annals of Pure and Applied Logic*, 100:247–277, 1999.
3. S. Berardi, C. Berline. $\beta\eta$ -complete models for system F , *Mathematical Structure in Computer Science*, 12:823–874, 2002.

4. C. Berline. From computation to foundations via functions and application: The λ -calculus and its webbed models. *Theoretical Computer Science*, 249:81–161, 2000.
5. C. Berline. Graph models of λ -calculus at work, and variations. *Math. Struct. in Comp. Science*, 16:185–221, 2006.
6. C. Berline, A. Salibra. Easiness in graph models, *Theoretical Computer Science* 354:4–23, 2006.
7. G. Berry. Stable models of typed lambda-calculi. In *Proc. 5th Int. Coll. on Automata, Languages and Programming*, LNCS 62, Springer-Verlag, Berline, 1978.
8. A. Bucciarelli, T. Ehrhard. Sequentiality and strong stability. *Sixth Annual IEEE Symposium on Logic in Computer Science (LICS 1991)*, IEEE Computer Society Press, 138–145, 1991.
9. A. Bucciarelli, A. Salibra. The minimal graph model of lambda calculus. MFCS'03, LNCS 2747, Springer-Verlag, 2003.
10. A. Bucciarelli, A. Salibra. The sensible graph theories of lambda calculus. *19th Annual IEEE Symposium on Logic in Computer Science (LICS 2004)*, IEEE Computer Society Press, 2004.
11. A. Bucciarelli, A. Salibra. Graph lambda theories. Preprint 2006 (submitted). <http://www.dsi.unive.it/~salibra/bs.pdf>
12. P. Di Gianantonio, F. Honsell, G.D. Plotkin. Uncountable limits and the lambda calculus. *Nordic J. Comput.*, 2:126–145, 1995.
13. P. Giannini, G. Longo. Effectively given domains and lambda-calculus models. *Information and Control*, 62:36–63, 1984.
14. A. Gruchalski. Computability on dI-Domains. *Information and Computation* 124:7–19, 1996.
15. F. Honsell, S. Ronchi della Rocca. An approximation theorem for topological lambda models and the topological incompleteness of lambda calculus. *Journal of Computer and System Sciences* 45:49–75, 1992.
16. R. Kerth. Isomorphism and equational equivalence of continuous lambda models. *Studia Logica* 61:403–415, 1998.
17. R. Kerth. On the construction of stable models of λ -calculus. *Theoretical Computer Science* 269:23–46, 2001.
18. G. Longo. Set-theoretical models of λ -calculus: theories, expansions and isomorphisms. *Ann. Pure Applied Logic* 24:153–188, 1983.
19. G.D. Plotkin. Set-theoretical and other elementary models of the λ -calculus. *Theoretical Computer Science* 121:351–409, 1993.
20. A. Salibra. A continuum of theories of lambda calculus without semantics. *16th Annual IEEE Symposium on Logic in Computer Science (LICS 2001)*, IEEE Computer Society Press, 334–343, 2001.
21. A. Salibra. Topological incompleteness and order incompleteness of the lambda calculus. (LICS'01 Special Issue). *ACM Transactions on Computational Logic* 4:379–401, 2003.
22. D.S. Scott. Continuous lattices. In *Toposes, Algebraic geometry and Logic*. LNM 274, Springer-Verlag, Berlin, 1972.
23. P. Selinger. Order-incompleteness and finite lambda reduction models. *Theoretical Computer Science* 309:43–63, 2003.
24. V. Stoltenberg-Hansen, I. Lindström, E. R. Griffor. *Mathematical theory of domains*. Cambridge Tracts in Theoretical Computer Science (No. 22), Cambridge University Press, 1994.
25. A. Visser. Numerations, λ -calculus and arithmetic. In: (J.R. Hindley and J.P. Seldin eds.) *To H.B. Curry: Essays on Combinatory Logic, Lambda-Calculus and Formalism*. Academic Press, New York, 259–284, 1980.