

# TD2 – Graphs, adjunctions, monads

Samuel Mimram

October 4, 2012

## 1 Graphs

1. Show that the category  $\mathbf{Cat}(\mathbf{Gr}, \mathbf{Set})$  of functors and natural transformations from the category with two objects 0, 1 and four morphisms

$$\text{id}_0 : 0 \rightarrow 0 \quad \text{id}_1 : 1 \rightarrow 1 \quad s, t : 1 \rightarrow 0$$

to the category  $\mathbf{Set}$  of sets and functions defines the category category of graphs, which is usually denoted  $\mathbf{Graph}$ .

2. Reformulate the definition of a category as a graph with some structure.
3. Explain that the category  $\mathbf{Cat}(\mathbf{Gr}_2, \mathbf{Set})$  of functors from the category  $\mathbf{Gr}_2$  with three objects 0, 1, 2 and height morphisms

$$\text{id}_0 : 0 \rightarrow 0 \quad \text{id}_1 : 1 \rightarrow 1 \quad \text{id}_2 : 2 \rightarrow 2 \quad s_1, t_1 : 2 \rightarrow 1 \quad s_0, t_0 : 1 \rightarrow 0 \quad s, t : 2 \rightarrow 0$$

with

$$s_0 \circ s_1 = s_0 \circ t_1 = s \quad \text{et} \quad t_0 \circ s_1 = t_0 \circ t_1 = t$$

defines a category category of 2-graphs and morphisms of 2-graphs.

4. Reformulate the definition of 2-categories using the notion of 2-graph.

## 2 Adjunctions between sets

We recall that a functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  is *left adjoint* to a functor  $G : \mathcal{D} \rightarrow \mathcal{C}$  if there is a natural bijection between  $\mathcal{D}(FA, B)$  and  $\mathcal{C}(A, GB)$ .

1. Suppose given two functions  $f : A \rightarrow B$  and  $g : B \rightarrow A$  between sets  $A$  and  $B$ . Show that the two following properties are equivalent:

- (i)  $f$  and  $g$  are bijections and  $f = g^{-1}$
- (ii)  $\forall a \in A, \forall b \in B, f(a) = b \text{ iff } a = f(b)$

2. Conclude that an adjunction between two discrete categories is a bijection.

## 3 The exception monad

We write  $\mathbf{pSet}$  for the category whose objects are *pointed sets*, i.e. pairs  $(A, a)$  where  $A$  is a set and  $a \in A$ , and morphisms  $f : (A, a) \rightarrow (B, b)$  are functions such that  $f(a) = b$ . Here the distinguished element of the pointed set will be seen as a particular value indicating an error or an exception.

1. Describe the *forgetful functor*  $U : \mathbf{pSet} \rightarrow \mathbf{Set}$  which to a pointed set associates the underlying set.

- Construct a functor  $F : \mathbf{Set} \rightarrow \mathbf{pSet}$  which is such that the sets  $\text{Hom}(FA, B)$  and  $\text{Hom}(A, UB)$  are isomorphic.
- [Facultative] Show that the families of isomorphisms

$$\varphi_{A,B} : \text{Hom}(FA, B) \rightarrow \text{Hom}(A, UB) \quad \text{and} \quad \psi_{A,B} : \text{Hom}(A, UB) \rightarrow \text{Hom}(FA, B)$$

described in previous question are natural. By “ $\varphi_{A,B}$  is *natural*”, we mean here that for every morphisms  $f : A \rightarrow A'$  in  $\mathbf{Set}$  and  $g : B \rightarrow B'$  in  $\mathbf{pSet}$  the diagram

$$\begin{array}{ccc} \text{Hom}(FA', B) & \xrightarrow{\phi_{A',B}} & \text{Hom}(A', UB) \\ g \circ - \circ Ff \downarrow & & \downarrow Ug \circ - \circ f \\ \text{Hom}(FA, B') & \xrightarrow{\phi_{A,B'}} & \text{Hom}(A, UB') \end{array}$$

commutes (in  $\mathbf{Set}$ ). Naturality of  $\psi$  is defined in a similar way.

- We recall that a *monad* consists of an endofunctor  $T : \mathcal{C} \rightarrow \mathcal{C}$  together with two natural transformations  $\mu : T \circ T \Rightarrow T$  and  $\eta : \text{id}_{\mathcal{C}} \Rightarrow T$  such that the following diagrams commute:

$$\begin{array}{ccc} T \circ T \circ T & \xrightarrow{T\mu} & T \circ T \\ \mu_T \downarrow & & \downarrow \mu \\ T \circ T & \xrightarrow{\mu} & T \end{array} \qquad \begin{array}{ccc} T & \xrightarrow{\eta_T} & T \circ T & \xleftarrow{T\eta} & T \\ \text{id}_T \searrow & & \downarrow \mu & & \swarrow \text{id}_T \\ & & T & & \end{array}$$

Represent those diagrams using pasting diagrams in the 2-category  $\mathbf{Cat}$ . Represent those diagrams using string diagrams.

- Describe a structure of monad on  $U \circ F$ .
- Given  $f : A \rightarrow B$  an OCaml function which might raise a unique exception  $e$  and  $g : B \rightarrow C$  a function which might raise a unique exception  $e'$ , construct a function corresponding to the composite of  $f$  and  $g$  which might raise a unique exception  $e''$ .
- We write  $\mathbf{Set}_T$  the category whose objects are the objects of  $\mathbf{Set}$  and morphisms  $f : A \rightarrow B$  in  $\mathbf{Set}_T$  are morphisms  $f : A \rightarrow TB$  in  $\mathbf{Set}$ . Compositions of two morphisms  $f : A \rightarrow B$  and  $g : B \rightarrow C$  in  $\mathbf{Set}_T$  is defined by  $g \circ f = \mu_C \circ Tg \circ f$  and identities are  $\text{id}_A = \eta_A$ . Show that the axioms of categories are satisfied.
- Give an explicit description of  $\mathbf{Set}_T$ .
- A *non-deterministic function* is a function that might return a set of values instead of a single value. How could we similarly define a category of non-deterministic functions by a Kleisli construction?

## 4 Free category on a graph

- Define the forgetful functor  $U : \mathbf{Cat} \rightarrow \mathbf{Graph}$ .
- Show that this functor  $F : \mathbf{Graph} \rightarrow \mathbf{Cat}$  admits a left adjoint.