

2-species exclusion processes and combinatorial algebras

Sylvie Corteel Arthur Nunge

IRIF, LIGM

March 2017

Non commutative symmetric functions

The algebra of noncommutative symmetric functions **Sym** is an algebra generalizing the symmetric functions. Its component of degree n has dimension 2^{n-1} . One can index its bases by compositions.

Non commutative symmetric functions

The algebra of noncommutative symmetric functions **Sym** is an algebra generalizing the symmetric functions. Its component of degree n has dimension 2^{n-1} . One can index its bases by compositions.

A composition of size n is a sequence of integers $I = (i_1, i_2, \dots, i_r)$ of sum n .

Non commutative symmetric functions

The algebra of noncommutative symmetric functions **Sym** is an algebra generalizing the symmetric functions. Its component of degree n has dimension 2^{n-1} . One can index its bases by compositions.

A composition of size n is a sequence of integers $I = (i_1, i_2, \dots, i_r)$ of sum n .

Complete basis (analog of h_λ)

For all n , define

$$S_n = \sum_{1 \leq j_1 \leq j_2 \leq \dots \leq j_n} a_{j_1} a_{j_2} \cdots a_{j_n}.$$

Non commutative symmetric functions

The algebra of noncommutative symmetric functions **Sym** is an algebra generalizing the symmetric functions. Its component of degree n has dimension 2^{n-1} . One can index its bases by compositions.

A composition of size n is a sequence of integers $I = (i_1, i_2, \dots, i_r)$ of sum n .

Complete basis (analog of h_λ)

For all n , define

$$S_n = \sum_{1 \leq j_1 \leq j_2 \leq \dots \leq j_n} a_{j_1} a_{j_2} \cdots a_{j_n}.$$

For any composition $I = (i_1, i_2, \dots, i_r)$,

$$S^I = S_{i_1} S_{i_2} \cdots S_{i_r}.$$

Non commutative symmetric functions

The algebra of noncommutative symmetric functions **Sym** is an algebra generalizing the symmetric functions. Its component of degree n has dimension 2^{n-1} . One can index its bases by compositions.

A composition of size n is a sequence of integers $I = (i_1, i_2, \dots, i_r)$ of sum n .

Complete basis (analog of h_λ)

For all n , define

$$S_n = \sum_{1 \leq j_1 \leq j_2 \leq \dots \leq j_n} a_{j_1} a_{j_2} \cdots a_{j_n}.$$

For any composition $I = (i_1, i_2, \dots, i_r)$,

$$S^I = S_{i_1} S_{i_2} \cdots S_{i_r}.$$

For example, $S_2(a_1, a_2, a_3) = a_1^2 + a_1 a_2 + a_1 a_3 + a_2^2 + a_2 a_3 + a_3^2$.

Ribbon basis

$$R_I = \sum_{J \preceq I} (-1)^{l(J) - l(I)} S^J.$$

For example, $R_{221} = S^{221} - S^{41} - S^{23} + S^5$.

Ribbon basis

$$R_I = \sum_{J \preceq I} (-1)^{l(J)-l(I)} S^J.$$

For example, $R_{221} = S^{221} - S^{41} - S^{23} + S^5$.

Polynomial realization

$$R_I = \sum_{\text{Des}(w)=I} w.$$

For example, $R_{221}(a_1, a_2) = a_1 a_2 a_1 a_2 a_1 + a_2 a_2 a_1 a_2 a_1$.

Tevlin's bases

In 2007 L. Tevlin defined the monomial (M_I) and fundamental (L_I) that are analog of the monomial basis and elementary basis of Sym. They both have binomial structure coefficients.

Tevlin's bases

In 2007 L. Tevlin defined the monomial (M_l) and fundamental (L_l) that are analog of the monomial basis and elementary basis of Sym. They both have binomial structure coefficients.

Transition matrices

The transition matrices between the ribbon basis and the fundamental basis of size 3 and 4 are:

$$\mathfrak{M}_3 = \begin{pmatrix} 1 & . & . & . \\ . & 2 & 1 & . \\ . & . & 1 & . \\ . & . & . & 1 \end{pmatrix}$$

$$\mathfrak{M}_4 = \begin{pmatrix} 1 & . & . & . & . & . & . & . \\ . & 3 & 2 & . & 1 & 1 & . & . \\ . & . & 2 & . & 1 & . & . & . \\ . & . & 1 & 3 & . & 2 & 1 & . \\ . & . & . & . & 1 & . & . & . \\ . & . & . & . & . & 2 & 1 & . \\ . & . & . & . & . & . & 1 & . \\ . & . & . & . & . & . & . & 1 \end{pmatrix}$$

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (*i.e.*, the values k such that $k + 1$ is on the left).

For $\sigma = 25783641$

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (i.e., the values k such that $k + 1$ is on the left).
For $\sigma = 2578364\mathbf{1}$, the recoils are $\{\mathbf{1}\}$

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (i.e., the values k such that $k + 1$ is on the left).

For $\sigma = \textcolor{red}{2}578\textcolor{blue}{3}641$, the recoils are $\{1\}$

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (i.e., the values k such that $k + 1$ is on the left).

For $\sigma = 2578\textcolor{red}{3}\textcolor{blue}{6}41$, the recoils are $\{1\}$

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (i.e., the values k such that $k + 1$ is on the left).

For $\sigma = 2\textcolor{blue}{5}7836\textcolor{red}{4}1$, the recoils are $\{1, \textcolor{red}{4}\}$

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (i.e., the values k such that $k + 1$ is on the left).

For $\sigma = 2\textcolor{red}{5}783\textcolor{blue}{6}41$, the recoils are $\{1, 4\}$

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (i.e., the values k such that $k + 1$ is on the left).

For $\sigma = 25783\textcolor{blue}{6}41$, the recoils are $\{1, 4, \textcolor{red}{6}\}$

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (i.e., the values k such that $k + 1$ is on the left).

For $\sigma = 25\textcolor{red}{7}\textcolor{blue}{8}3641$, the recoils are $\{1, 4, 6\}$

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (i.e., the values k such that $k + 1$ is on the left).

For $\sigma = 25783641$, the recoils are $\{\textcolor{red}{1}, 4, 6\}$ so $\text{Rec}(25783641) = \textcolor{red}{1}$.

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (i.e., the values k such that $k + 1$ is on the left).

For $\sigma = 25783641$, the recoils are $\{1, 4, 6\}$ so $\text{Rec}(25783641) = 13$.

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (i.e., the values k such that $k + 1$ is on the left).

For $\sigma = 25783641$, the recoils are $\{1, 4, 6\}$ so $\text{Rec}(25783641) = 13\textcolor{red}{2}$.

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (i.e., the values k such that $k + 1$ is on the left).

For $\sigma = 25783641$, the recoils are $\{1, 4, 6\}$ so $\text{Rec}(25783641) = 1322$.

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (*i.e.*, the values k such that $k + 1$ is on the left).
For $\sigma = 25783641$, the recoils are $\{1, 4, 6\}$ so $\text{Rec}(25783641) = 1322$.
- $\text{GC}(\sigma)$ is the composition associated with the values of descents (*i.e.*, the values $k = \sigma_i$ such that $\sigma_i > \sigma_{i+1}$) minus one.
For $\sigma = 25783641$, $\text{GC}(\sigma) = .$

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (*i.e.*, the values k such that $k + 1$ is on the left).
For $\sigma = 25783641$, the recoils are $\{1, 4, 6\}$ so $\text{Rec}(25783641) = 1322$.
- $\text{GC}(\sigma)$ is the composition associated with the values of descents (*i.e.*, the values $k = \sigma_i$ such that $\sigma_i > \sigma_{i+1}$) minus one.
For $\sigma = \textcolor{red}{2}\textcolor{blue}{5}783641$, $\text{GC}(\sigma) = .$

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (*i.e.*, the values k such that $k + 1$ is on the left).
For $\sigma = 25783641$, the recoils are $\{1, 4, 6\}$ so $\text{Rec}(25783641) = 1322$.
- $\text{GC}(\sigma)$ is the composition associated with the values of descents (*i.e.*, the values $k = \sigma_i$ such that $\sigma_i > \sigma_{i+1}$) minus one.
For $\sigma = 2578\textcolor{red}{3}\textcolor{blue}{6}41$, $\text{GC}(\sigma) = .$

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (*i.e.*, the values k such that $k + 1$ is on the left).
For $\sigma = 25783641$, the recoils are $\{1, 4, 6\}$ so $\text{Rec}(25783641) = 1322$.
- $\text{GC}(\sigma)$ is the composition associated with the values of descents (*i.e.*, the values $k = \sigma_i$ such that $\sigma_i > \sigma_{i+1}$) minus one.
For $\sigma = 257836\textcolor{blue}{4}1$, $\text{GC}(\sigma) = \textcolor{red}{3}$.

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (*i.e.*, the values k such that $k + 1$ is on the left).
For $\sigma = 25783641$, the recoils are $\{1, 4, 6\}$ so $\text{Rec}(25783641) = 1322$.
- $\text{GC}(\sigma)$ is the composition associated with the values of descents (*i.e.*, the values $k = \sigma_i$ such that $\sigma_i > \sigma_{i+1}$) minus one.
For $\sigma = 2\textcolor{red}{5}\textcolor{blue}{7}83641$, $\text{GC}(\sigma) = 3$.

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (i.e., the values k such that $k + 1$ is on the left).
For $\sigma = 25783641$, the recoils are $\{1, 4, 6\}$ so $\text{Rec}(25783641) = 1322$.
- $\text{GC}(\sigma)$ is the composition associated with the values of descents (i.e., the values $k = \sigma_i$ such that $\sigma_i > \sigma_{i+1}$) minus one.
For $\sigma = 25783\textcolor{red}{6}\textcolor{blue}{4}1$, $\text{GC}(\sigma) = 3\textcolor{red}{2}$.

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (i.e., the values k such that $k + 1$ is on the left).
For $\sigma = 25783641$, the recoils are $\{1, 4, 6\}$ so $\text{Rec}(25783641) = 1322$.
- $\text{GC}(\sigma)$ is the composition associated with the values of descents (i.e., the values $k = \sigma_i$ such that $\sigma_i > \sigma_{i+1}$) minus one.
For $\sigma = 25\textcolor{red}{7}\textcolor{blue}{8}3641$, $\text{GC}(\sigma) = 32$.

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (i.e., the values k such that $k + 1$ is on the left).
For $\sigma = 25783641$, the recoils are $\{1, 4, 6\}$ so $\text{Rec}(25783641) = 1322$.
- $\text{GC}(\sigma)$ is the composition associated with the values of descents (i.e., the values $k = \sigma_i$ such that $\sigma_i > \sigma_{i+1}$) minus one.
For $\sigma = 25783641$, $\text{GC}(\sigma) = 322$.

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (*i.e.*, the values k such that $k + 1$ is on the left).
For $\sigma = 25783641$, the recoils are $\{1, 4, 6\}$ so $\text{Rec}(25783641) = 1322$.
- $\text{GC}(\sigma)$ is the composition associated with the values of descents (*i.e.*, the values $k = \sigma_i$ such that $\sigma_i > \sigma_{i+1}$) minus one.
For $\sigma = 25783641$, $\text{GC}(\sigma) = 3221$.

Statistics on permutations

- $\text{Rec}(\sigma)$ is the composition associated with the values of recoils (*i.e.*, the values k such that $k + 1$ is on the left).
For $\sigma = 25783641$, the recoils are $\{1, 4, 6\}$ so $\text{Rec}(25783641) = 1322$.
- $\text{GC}(\sigma)$ is the composition associated with the values of descents (*i.e.*, the values $k = \sigma_i$ such that $\sigma_i > \sigma_{i+1}$) minus one.
For $\sigma = 25783641$, $\text{GC}(\sigma) = 3221$.

Combinatorial interpretation (F. Hivert, J.-C. Novelli, L. Tevlin, J.-Y. Thibon, 2009)

$$\begin{pmatrix} 1 & . & . & . & . & . & . & . \\ . & 3 & 2 & . & 1 & 1 & . & . \\ . & . & 2 & . & 1 & . & . & . \\ . & . & 1 & 3 & . & 2 & 1 & . \\ . & . & . & . & 1 & . & . & . \\ . & . & . & . & . & 2 & 1 & . \\ . & . & . & . & . & . & 1 & . \\ . & . & . & . & . & . & . & 1 \end{pmatrix}$$

GC \ Rec	4	31	22	211	13	121	112	1111
4	1234							
31		1243, 1423 4123	1342 3412		2341	2413		
22			1324 3124		2314			
211			3142	1432, 4132 4312		2431 4231	3241	
13					2134			
121						2143 4213	3421	
112							3214	
1111								4321

ASEP

The ASEP (Asymmetric Simple Exclusion Process) is a physical model in which particles hop back and forth (and in and out) of a one-dimensional lattice.



ASEP

The ASEP (Asymmetric Simple Exclusion Process) is a physical model in which particles hop back and forth (and in and out) of a one-dimensional lattice.



ASEP

The ASEP (Asymmetric Simple Exclusion Process) is a physical model in which particles hop back and forth (and in and out) of a one-dimensional lattice.



ASEP

The ASEP (Asymmetric Simple Exclusion Process) is a physical model in which particles hop back and forth (and in and out) of a one-dimensional lattice.



ASEP

The ASEP (Asymmetric Simple Exclusion Process) is a physical model in which particles hop back and forth (and in and out) of a one-dimensional lattice.



ASEP

The ASEP (Asymmetric Simple Exclusion Process) is a physical model in which particles hop back and forth (and in and out) of a one-dimensional lattice.



ASEP

The ASEP (Asymmetric Simple Exclusion Process) is a physical model in which particles hop back and forth (and in and out) of a one-dimensional lattice.



We associate the composition 1213 with the above state of the ASEP.

ASEP

The ASEP (Asymmetric Simple Exclusion Process) is a physical model in which particles hop back and forth (and in and out) of a one-dimensional lattice.



We associate the composition 1213 with the above state of the ASEP.

ASEP

The ASEP (Asymmetric Simple Exclusion Process) is a physical model in which particles hop back and forth (and in and out) of a one-dimensional lattice.



We associate the composition 1213 with the above state of the ASEP.

Combinatorial study of the ASEP

Let I be a composition associated with a state of the ASEP, the un-normalized steady-state probability of this state is given by:

Permutations

$$\sum_{\text{GC}(\sigma)=I} q^{\#_{31-2}(\sigma)}$$

Laguerre histories

$$\sum_{\text{type}(P)=I} q^{\text{weight}(P)}$$

Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.

Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.

The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.

The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

- \longrightarrow iff $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$;

Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.

The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

- \longrightarrow iff $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$;
- \nearrow iff $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$;

Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.

The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

- \longrightarrow iff $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$;
- \dashrightarrow iff $\sigma_{i-1} > \sigma_i > \sigma_{i+1}$;
- \nearrow iff $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$;

Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.

The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

- \longrightarrow iff $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$;
- \dashrightarrow iff $\sigma_{i-1} > \sigma_i > \sigma_{i+1}$;
- \nearrow iff $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$;
- \searrow iff $\sigma_{i-1} < \sigma_i > \sigma_{i+1}$.

Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.

The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

- \longrightarrow iff $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$;
- \dashrightarrow iff $\sigma_{i-1} > \sigma_i > \sigma_{i+1}$;
- \nearrow iff $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$;
- \searrow iff $\sigma_{i-1} < \sigma_i > \sigma_{i+1}$.

Its weight is the number of $31-2$ patterns where k is the 2.

Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \rightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.

The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

- \longrightarrow iff $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$;
- \nearrow iff $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$;
- \dashrightarrow iff $\sigma_{i-1} > \sigma_i > \sigma_{i+1}$;
- \searrow iff $\sigma_{i-1} < \sigma_i > \sigma_{i+1}$.

Its weight is the number of $31-2$ patterns where k is the 2.

 $25783641 \longrightarrow$

Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.


The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

- \longrightarrow iff $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$;
- \dashrightarrow iff $\sigma_{i-1} > \sigma_i > \sigma_{i+1}$;
- \nearrow iff $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$;
- \searrow iff $\sigma_{i-1} < \sigma_i > \sigma_{i+1}$.

Its weight is the number of $31-2$ patterns where k is the 2.

2578364**1** \longrightarrow



Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.

The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

- \longrightarrow iff $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$;
- \dashrightarrow iff $\sigma_{i-1} > \sigma_i > \sigma_{i+1}$;
- \nearrow iff $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$;
- \searrow iff $\sigma_{i-1} < \sigma_i > \sigma_{i+1}$.

Its weight is the number of $31-2$ patterns where k is the 2.

25783641 \longrightarrow



Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.


The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

- \longrightarrow iff $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$;
- \dashrightarrow iff $\sigma_{i-1} > \sigma_i > \sigma_{i+1}$;
- \nearrow iff $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$;
- \searrow iff $\sigma_{i-1} < \sigma_i > \sigma_{i+1}$.

Its weight is the number of $31-2$ patterns where k is the 2.

2578**3**641 \longrightarrow



Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.


The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

- \longrightarrow iff $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$;
- \dashrightarrow iff $\sigma_{i-1} > \sigma_i > \sigma_{i+1}$;
- \nearrow iff $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$;
- \searrow iff $\sigma_{i-1} < \sigma_i > \sigma_{i+1}$.

Its weight is the number of $31-2$ patterns where k is the 2.

25783641 \longrightarrow



Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.


The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

- \longrightarrow iff $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$;
- \dashrightarrow iff $\sigma_{i-1} > \sigma_i > \sigma_{i+1}$;
- \nearrow iff $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$;
- \searrow iff $\sigma_{i-1} < \sigma_i > \sigma_{i+1}$.

Its weight is the number of $31-2$ patterns where k is the 2.

25783641 \longrightarrow



Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.


The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

- \longrightarrow iff $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$;
- \dashrightarrow iff $\sigma_{i-1} > \sigma_i > \sigma_{i+1}$;
- \nearrow iff $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$;
- \searrow iff $\sigma_{i-1} < \sigma_i > \sigma_{i+1}$.

Its weight is the number of $31-2$ patterns where k is the 2.

25783**6**41 \longrightarrow



Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.


The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

- \longrightarrow iff $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$;
- \dashrightarrow iff $\sigma_{i-1} > \sigma_i > \sigma_{i+1}$;
- \nearrow iff $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$;
- \searrow iff $\sigma_{i-1} < \sigma_i > \sigma_{i+1}$.

Its weight is the number of $31-2$ patterns where k is the 2.

25783641 \longrightarrow



Laguerre histories

A Laguerre history is a weighted colored Motzkin path such that

- for a step \longrightarrow or \nearrow , $0 \leq w \leq h$;
- for a step \dashrightarrow or \searrow , $0 \leq w \leq h - 1$.


The Françon-Viennot bijection

Let $k = \sigma_i$ be a value of a permutation σ . The k -th step of $\phi_{FV}(\sigma)$ is

- \longrightarrow iff $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$;
- \dashrightarrow iff $\sigma_{i-1} > \sigma_i > \sigma_{i+1}$;
- \nearrow iff $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$;
- \searrow iff $\sigma_{i-1} < \sigma_i > \sigma_{i+1}$.


Its weight is the number of $31-2$ patterns where k is the 2.

25783641 \longrightarrow




2-ASEP

The 2-ASEP is a generalization of the ASEP with two kinds of particles.




2-ASEP

The 2-ASEP is a generalization of the ASEP with two kinds of particles.



2-ASEP


The 2-ASEP is a generalization of the ASEP with two kinds of particles.



A segmented composition is a sequence of integers separated by comas or bars.

2-ASEP


The 2-ASEP is a generalization of the ASEP with two kinds of particles.



A segmented composition is a sequence of integers separated by comas or bars. We associate the segmented composition $12|11|2$ with the above state of the 2-ASEP.

2-ASEP

The 2-ASEP is a generalization of the ASEP with two kinds of particles.



A segmented composition is a sequence of integers separated by comas or bars. We associate the segmented composition $12|11|2$ with the above state of the 2-ASEP.

Combinatorial study of the 2-ASEP

Let \mathbf{l} be a segmented composition associated with a state of the 2-ASEP, the un-normalized steady-state probability of this state is given by:

Partially signed permutations

$$\sum_{\text{GC}(\sigma)=\mathbf{l}} q^{\#_{31-2}(\sigma)+\#_{(31,2)}(\sigma)}$$

Marked Laguerre histories

$$\sum_{\text{type}(P)=\mathbf{l}} q^{\text{weight}(P)}$$

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = .$

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = \overline{1}|$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|\overline{1}$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .

For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|1$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|12$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .

For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|12$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|122$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|122$.
- $\text{GC}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{GC}(\overline{2}57836\overline{4}1) = .$

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|122$.
- $\text{GC}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{GC}(\overline{2}57836\overline{4}1) = \overline{1}|$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|122$.
- $\text{GC}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}578\textcolor{red}{3}\textcolor{blue}{6}\overline{4}1$, $\text{GC}(\overline{2}57836\overline{4}1) = 1|$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|122$.
- $\text{GC}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{GC}(\overline{2}57836\overline{4}1) = 1|2|$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|122$.
- $\text{GC}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{GC}(\overline{2}57836\overline{4}1) = 1|2|$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|122$.
- $\text{GC}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}5783\overline{6}\overline{4}1$, $\text{GC}(\overline{2}5783\overline{6}\overline{4}1) = 1|2|\overline{2}$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|122$.
- $\text{GC}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}5\overline{7}836\overline{4}1$, $\text{GC}(\overline{2}57836\overline{4}1) = 1|2|2$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|122$.
- $\text{GC}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57\textcolor{red}{8}\textcolor{blue}{3}6\overline{4}1$, $\text{GC}(\overline{2}57836\overline{4}1) = 1|2|2\textcolor{red}{2}$.

Partially signed permutations

A partially signed permutation is a permutation where all values except 1 can be overlined. For example, $\sigma = \overline{2}57836\overline{4}1$.

Statistics on partially signed permutations

- $\text{Rec}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{Rec}(\overline{2}57836\overline{4}1) = 1|2|122$.
- $\text{GC}(\sigma)$ is computed as previously, we add bars on the composition to retrieve the position of the overlined values in σ .
For $\sigma = \overline{2}57836\overline{4}1$, $\text{GC}(\overline{2}57836\overline{4}1) = 1|2|221$.

Marked Laguerre histories

A marked Laguerre history is a Laguerre history where all the steps but the first can be marked. Any marked step starting from height h increases its weight by h

Marked Laguerre histories

A marked Laguerre history is a Laguerre history where all the steps but the first can be marked. Any marked step starting from height h increases its weight by h

generalization of the Françon-Viennot bijection

Let σ be a partially signed permutation, apply ϕ_{FV} to σ . Then, mark the steps that correspond to the signed values.

Marked Laguerre histories

A marked Laguerre history is a Laguerre history where all the steps but the first can be marked. Any marked step starting from height h increases its weight by h


generalization of the Françon-Viennot bijection

Let σ be a partially signed permutation, apply ϕ_{FV} to σ . Then, mark the steps that correspond to the signed values.

$$\bar{2}57836\bar{4}1 \longrightarrow$$


A marked Laguerre history is a Laguerre history where all the steps but the first can be marked. Any marked step starting from height h increases its weight by h

Let σ be a partially signed permutation, apply ϕ_{FV} to σ . Then, mark the steps that correspond to the signed values.



A marked Laguerre history is a Laguerre history where all the steps but the first can be marked. Any marked step starting from height h increases its weight by h

Let σ be a partially signed permutation, apply ϕ_{FV} to σ . Then, mark the steps that correspond to the signed values.



The algebra of segmented compositions

In 2007, J.-C. Novelli and J.-Y. Thibon defined the algebra of segmented compositions (**SCQSym**) and its complete and ribbon bases.

The algebra of segmented compositions

In 2007, J.-C. Novelli and J.-Y. Thibon defined the algebra of segmented compositions (**SCQSym**) and its complete and ribbon bases.

Complete basis

$$S_I \cdot S_J = S_{I \cdot J} + S_{I|J}$$

For example, $S_{21|1} \cdot S_{32|21} = S_{21|132|21} + S_{21|1|32|21}$.

The algebra of segmented compositions

In 2007, J.-C. Novelli and J.-Y. Thibon defined the algebra of segmented compositions (**SCQSym**) and its complete and ribbon bases.

Complete basis

$$S_I \cdot S_J = S_{I \cdot J} + S_{I|J}$$

For example, $S_{21|1} \cdot S_{32|21} = S_{21|132|21} + S_{21|1|32|21}$.

Ribbon basis

Again we have

$$R_I = \sum_{J \preceq I} (-1)^{l(J) - l(I)} S_J.$$

For example, $R_{22|41} = S_{22|41} - S_{4|41} - S_{22|5} + S_{4|5}$.

Analogue of Tevlin's bases

We define a monomial basis (M_I) and a fundamental basis (L_I) in **SCQSym**.

Analogue of Tevlin's bases

We define a monomial basis (M_I) and a fundamental basis (L_I) in **SCQSym**.

Transition matrix

The coefficients in the transition matrices from the ribbon basis to the fundamental basis are

$$(\mathcal{M}_n)_{I,J} = \#\{\sigma \mid \text{GC}(\sigma) = I, \text{Rec}(\sigma) = J\}$$

$$\mathcal{M}_3 = \left(\begin{array}{cccc|cc|cc|c} 1 & . & . & . & . & . & . & . & . \\ . & 2 & 1 & . & . & . & . & . & . \\ . & . & 1 & . & . & . & . & . & . \\ . & . & . & 1 & . & . & . & . & . \\ \hline . & . & . & . & 3 & 1 & . & . & . \\ . & . & . & . & . & 2 & . & . & . \\ \hline . & . & . & . & . & . & 2 & . & . \\ . & . & . & . & . & . & 1 & 3 & . \\ \hline . & . & . & . & . & . & . & . & 6 \end{array} \right)$$

q -analogs

We define q -analogs of S_I and M_I and we obtain the four following transition matrices.

q -analogs

We define q -analogs of S_I and M_I and we obtain the four following transition matrices.

$$\begin{pmatrix}
 1 & 1 & 1 & 1 & . & . & . & . & . \\
 1 & q^2 + q + 1 & q + 1 & q^2 + 2q + 1 & . & . & . & . & . \\
 1 & 1 & q + 1 & q + 1 & . & . & . & . & . \\
 1 & q^2 + q + 1 & q^2 + q + 1 & q^3 + 2q^2 + 2q + 1 & . & . & . & . & . \\
 . & . & . & . & q^2 + q + 1 & q^2 + 2q + 1 & . & . & . \\
 . & . & . & . & q^2 + q + 1 & q^3 + 2q^2 + 2q + 1 & . & . & . \\
 . & . & . & . & . & . & q + 1 & q + 1 & . \\
 . & . & . & . & . & . & q^2 + q + 1 & q^3 + 2q^2 + 2q + 1 & . \\
 . & . & . & . & . & . & . & . & q^3 + 2q^2 + 2q + 1
 \end{pmatrix}$$

$S_I \rightarrow M_I$

q -analogs

We define q -analogs of S_I and M_I and we obtain the four following transition matrices.

$$\begin{pmatrix}
 1 & 1 & 1 & 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\
 \cdot & q+1 & 1 & q+2 & \cdot & \cdot & \cdot & \cdot & \cdot \\
 \cdot & \cdot & 1 & 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & \cdot & q^2+q+1 & q^2+2q+1 & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & \cdot & \cdot & q+1 & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & q+1 & q+1 & \cdot \\
 \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & q & q^2+2q+1 & \cdot \\
 \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & q^3+2q^2+2q+1
 \end{pmatrix}$$

$S_I \rightarrow L_I$

q -analogs

We define q -analogs of S_I and M_I and we obtain the four following transition matrices.

$$\begin{pmatrix}
 1 & 1 & 1 & 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\
 \cdot & q+1 & 1 & q+2 & \cdot & \cdot & \cdot & \cdot & \cdot \\
 \cdot & \cdot & 1 & 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & \cdot & q^2+q+1 & q^2+2q+1 & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & \cdot & \cdot & q+1 & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & q+1 & q+1 & \cdot \\
 \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & q & q^2+2q+1 & \cdot \\
 \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & q^3+2q^2+2q+1
 \end{pmatrix}$$

$S_I \rightarrow L_J$

$$\begin{pmatrix}
 1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
 \cdot & q+1 & 1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
 \cdot & \cdot & 1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & \cdot & q^2+q+1 & q & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & \cdot & \cdot & q+1 & \cdot & \cdot & \cdot \\
 \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & q+1 & \cdot & \cdot \\
 \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & q & q^2+q+1 & \cdot \\
 \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & q^3+2q^2+2q+1
 \end{pmatrix}$$

$R_I \rightarrow L_J$

Enumerative result

We obtain an enumerative formula for the probabilities of the 2-ASEP. Let I be a segmented composition, the steady-state probability of the state of the 2-ASEP encoded by I is given by the following formula.

$$[r+1]_q! \sum_{I \succeq J} \left(\frac{-1}{q} \right)^{I(I)-I(J)} q^{-\text{st}(I,J)} c_J(q)$$

Where $c_J(q) = [k]_q^{j_1} [k-1]_q^{j_2} \cdots [2]_q^{j_{k-1}} [1]_q^{j_k}$.

Perspectives

- Prove the combinatorial interpretation of the coefficients of the q -matrices.

Perspectives

- Prove the combinatorial interpretation of the coefficients of the q -matrices.
- find α and β statistics on partially signed permutations.

Perspectives

- Prove the combinatorial interpretation of the coefficients of the q -matrices.
- find α and β statistics on partially signed permutations.
- Understand the refinement (GC, Rec) on the 2-ASEP.