Signed bipartite circular cliques and a bipartite analogue of Grötzsch's theorem

Reza Naserasr¹ and Zhouningxin Wang^{*2}

¹Université de Paris, IRIF, CNRS, F-75013 Paris, France. ²School of Mathematical Sciences and LPMC, Nankai University, Tianjin 300071, China. Email addresses: reza@irif.fr, wangzhou@nankai.edu.cn.

Abstract

A circular r-coloring of a signed graph (G, σ) is to assign points of a circle of circumference r, $r \geq 2$, to the vertices of G such that vertices connected by a positive edge are at circular distance at least 1 and vertices connected by a negative edge are at circular distance at most $\frac{r}{2} - 1$. The restriction of the notion of circular colorings to the class of signed bipartite graphs is already of high interest because the circular chromatic number of an (unsigned) graph can be obtained by bounding the circular chromatic number of an associated signed bipartite graph.

In this paper, we define signed bipartite circular cliques $B_{p;q}^s$ and $\hat{B}_{p;q}^s$ having the property that a signed bipartite graph admits a circular $\frac{p}{q}$ -coloring if and only if it admits an edge-sign preserving homomorphism to $B_{p;q}^s$ and a switching homomorphism to $\hat{B}_{p;q}^s$, respectively. Then as a bipartite analogue of Grötzsch's theorem, we prove that every signed bipartite planar graph of negative girth at least 6 admits a circular 3-coloring.

1 Introduction

A signed graph is a graph G (allowing loops and multi-edges) together with an assignment $\sigma : E(G) \rightarrow \{+, -\}$, denoted (G, σ) . We note that signed graphs are allowed to have multi-edges, but we consider multi-edges only if they are of different signs. The signed graph on two vertices connected by two parallel edges of different signs is called *digon*. Furthermore, unless specified, graphs are considered to have no loop. The sign of a closed walk of (G, σ) is the product of signs of all its edges (allowing repetition). Given a signed graph (G, σ) and a vertex v of (G, σ) , a switching at v is to switch the signs of all the edges incident to v. We say a signed graph (G, σ') is switching equivalent to (G, σ) if it is obtained from (G, σ) by a series of switchings at vertices. In this case, we say the signature σ' is equivalent to σ . It has been proved in [25] that two signed graphs (G, σ_1) and (G, σ_2) are switching equivalent if and only if they have the same set of negative cycles.

A switching homomorphism, or simply a homomorphism, of a signed graph (G, σ) to (H, π) is a mapping of V(G) and E(G) to V(H) and E(H) (respectively) such that the adjacencies, the incidences and the signs of the closed walks are preserved. When there exists such a homomorphism, we write $(G, \sigma) \to (H, \pi)$. A homomorphism of (G, σ) to (H, π) is said to be *edge-sign preserving* if it, furthermore, preserves the signs of the edges. When there exists an edge-sign preserving homomorphism of (G, σ) to (H, π) , we write $(G, \sigma) \xrightarrow{s.p.} (H, \pi)$. The connection between these two kinds of homomorphisms is established as follows: Given two signed graphs (G, σ) and (H, π) , $(G, \sigma) \to (H, \pi)$ if and only if there exists a signature σ'

^{*}Corresponding author.

which is equivalent to σ , such that $(G, \sigma') \xrightarrow{s.p.} (H, \pi)$. An equivalent reformulation of this is through the following definition.

Definition 1.1. Given a signed graph (G, σ) , the Double Switch Graph of it, denoted $DSG(G, \sigma)$, is the signed graph built from two disjoint copies (G_1, σ_1) and (G_2, σ_2) of (G, σ) by adding the following set of edges in between. If xy is a positive (resp. negative) edge of (G, σ) , then x_1y_2 and x_2y_1 are negative (resp. positive) edges of $DSG(G, \sigma)$. Here x_1, x_2, y_1, y_2 are representing copies of x and y in G_1 and G_2 in the most natural way.

The connection between the two notions of homomorphisms is as follows.

Theorem 1.2. [1] A signed graph (G, σ) admits a switching homomorphism to (H, π) if and only if it admits an edge-sign preserving homomorphism to $DSG(H, \pi)$.

We note that in building $DSG(H, \pi)$ for the purpose of this theorem, if for a vertex x there already exists a vertex x' which is like obtained from x by a switching, then we do not need to add a copy of x. When (H, π) has the property that for each vertex x of it, there is a such vertex x' (which is like a switched copy of x), then a signed graph (G, σ) admits a homomorphism to (H, π) if and only if it also admits an edge-sign preserving homomorphism to (H, π) . That is because if a vertex u of (G, σ) is mapped to the vertex x of (H, π) after a switching, then one may instead map u to x' without a switching. We note that some of the circular cliques that we will discuss here are of this form.

Observe that the parity of the lengths and the signs of closed walks are preserved by a homomorphism. Given a signed graph (G, σ) and an element $ij \in \mathbb{Z}_2^2$, we define $g_{ij}(G, \sigma)$ to be the length of a shortest closed walk whose number of negative edges modulo 2 is i and whose length modulo 2 is j. When there exists no such closed walk, we say $g_{ij}(G, \sigma) = \infty$. For any signed graph (G, σ) containing at least one edge, $g_{00}(G, \sigma) = 2$. In general, the length of a shortest negative closed walk of a signed graph is said to be its *negative girth*. By the definition of the homomorphism of signed graphs, we have the following no-homomorphism lemma.

Lemma 1.3. [17][No-homomorphism lemma] If $(G, \sigma) \to (H, \pi)$, then $g_{ij}(G, \sigma) \geq g_{ij}(H, \pi)$ for each $ij \in \mathbb{Z}_2^2$.

1.1 Homomorphisms of signed bipartite graphs

Given a graph G, we construct a signed bipartite graph S(G) as follows: The vertex set of S(G) consists of V(G) and $\{x_{uv}, y_{uv} : uv \in E(G)\}$; For the edge set, we join each of x_{uv} and y_{uv} to each of u and v; For the signature, we assign signs such that each 4-cycle $ux_{uv}vy_{uv}$ is negative. Intuitively, in constructing S(G), we replace each edge uv of G with a negative 4-cycle. We note there is more than one choice of signature here, and moreover, having vertices already labeled, not every two such signatures are switching equivalent. That which of the two sides of a 4-cycle is chosen to be negative makes a difference here. However, up to a switching isomorphism, any two signature choices of S(G) are the same.

The first easy observation on S(G) is that it is a signed bipartite graph where in one part all vertices are of degree 2. This construction was introduced in [16] where the next two theorems are proved in order to show the importance of the study of homomorphisms of signed bipartite graphs.

Theorem 1.4. [16] Given graphs G and H, $G \to H$ if and only if $S(G) \to S(H)$.

Theorem 1.5. [16] Given a graph G, we have the following.

- $\chi(G) \leq 2$ if and only if $S(G) \rightarrow (K_{2,2}, e)$;
- $\chi(G) \leq k$ if and only if $S(G) \rightarrow (K_{k,k}, M)$ for $k \geq 3$.

Here $(K_{k,k}, M)$ is a signed graph on the complete bipartite graph $K_{k,k}$ where the edges in a perfect matching each is assigned with a negative sign. As the problem of mapping signed graphs to $(K_{k,k}, M)$ could capture the problem of the coloring of ordinary graphs, signed graphs $(K_{k,k}, M)$ are of special interests in the study of the homomorphism of signed graphs. In particular, when k = 4, we have a restatement of the 4-Color Theorem as follows.

Theorem 1.6. [4-Color Theorem restated] For any planar graph $G, S(G) \rightarrow (K_{4,4}, M)$.

Notice that the planarity is preserved when we construct S(G) from a planar graph G. Moreover, based on an edge-coloring result of B. Guenin [6] which in turn is based on the 4-Color Theorem, the following strengthening of the 4-Color Theorem has been proved in [16]: Every signed planar graph (G, σ) satisfying that $g_{ij}(G, \sigma) \ge g_{ij}(K_{4,4}, M)$ for $ij \in \mathbb{Z}_2^2$ maps to $(K_{4,4}, M)$. Since [6] is not available, for a direct proof of the statement we refer to [19]. A signed graph (G, σ) satisfies the conditions $g_{ij}(G, \sigma) \ge g_{ij}(K_{4,4}, M)$ if and only if (G, σ) is bipartite and has no digon. The next theorem is a reformulation of this strengthening of the 4-Color Theorem.

Theorem 1.7. [19] Every signed bipartite planar simple graph admits a homomorphism to $(K_{4,4}, M)$.

Motivated by these results, in this work we are interested in the study of circular chromatic numbers of signed bipartite graphs on their own. In particular, after noting that all signed bipartite (simple) planar graphs map to $(K_{4,4}, M)$, and that $(K_{4,4}, M)$ is the smallest signed bipartite simple graph having that property, we would like to ask under what girth condition we may use a subgraph of $(K_{4,4}, M)$ as a homomorphism target. To this end, we prove the following bipartite analogue of the Grötzsch theorem in this work, noting that our proof is based on the 4-Color Theorem.

Theorem 1.8. Every signed bipartite planar graph of negative girth at least 6 admits a homomorphism to $(K_{3,3}, M)$. Moreover, the girth condition is best possible.

2 Circular chromatic number of signed graphs

Given a signed graph (G, σ) and a positive integer k, a 0-free 2k-coloring of (G, σ) (introduced in [24]) is a mapping $f: V(G) \to \{\pm 1, \pm 2, \ldots, \pm k\}$ such that for any edge e = uv, $f(u) \neq \sigma(e)f(v)$. The notion of the circular coloring of signed graphs defined in [18] is a common extension of circular colorings of graphs and 0-free 2k-colorings of signed graphs.

For a real number $r \ge 1$, let C^r be a circle of circumference r. For two points x, y on C^r , the distance between x and y on C^r , denoted $d_{C^r}(x, y)$, is the length of the shorter arc of C^r connecting x and y. For each point x on C^r , the *antipodal* of x, denoted \bar{x} , is the unique point at distance $\frac{r}{2}$ from x.

Given a real number r, a *circular r-coloring* of a signed graph (G, σ) is a mapping $\varphi : V(G) \to C^r$ such that

- for each positive edge uv of (G, σ) , $d_{C^r}(\varphi(u), \varphi(v)) \ge 1$;
- for each negative edge uv of $(G, \sigma), d_{C^r}(\varphi(u), \overline{\varphi(v)}) \ge 1$.

The *circular chromatic number* of a signed graph (G, σ) is defined as

 $\chi_c(G,\sigma) = \inf\{r \ge 1 : (G,\sigma) \text{ admits a circular } r\text{-coloring}\}.$

We note that the infimum is attained and can then be replaced with the minimum. In particular, the circular chromatic number is always a rational. For integers $p \ge 2q > 0$ such that p is even, the signed circular clique $K_{p;q}^s$ has the vertex set $[p] = \{0, 1, \ldots, p-1\}$, in which ij is a positive edge if and only if $q \le |i-j| \le p-q$ and ij is a negative edge if and only if either $|i-j| \le \frac{p}{2} - q$ or $|i-j| \ge \frac{p}{2} + q$. Moreover, let $\widehat{K}_{p;q}^s$ be the signed subgraph of $K_{p;q}^s$ induced by vertices $\{0, 1, \ldots, \frac{p}{2} - 1\}$. In this definition loops are allowed, and indeed, by the definition, there will be a negative loop on each vertex but there will be no positive loop. As shown in [18], the following statements are equivalent:

- (G, σ) admits a circular $\frac{p}{q}$ -coloring;
- (G, σ) admits an edge-sign preserving homomorphism to $K^s_{p;a}$;
- (G, σ) admits a switching homomorphism to $\widehat{K}^{s}_{p;q}$.

In other words, in the order induced by edge-sign preserving homomorphism on the class of all signed graphs, the circular chromatic number of a signed graph (G, σ) is the smallest value of a rational number $\frac{p}{q}$ such that $(G, \sigma) \xrightarrow{s.p.} K_{p;q}^s$. Normally we choose the minimal element (the *core*) of each homomorphically equivalent class to represent the class. In such cases we will choose $K_{p;q}^s$ where p is an even integer and, with respect to this condition, $\frac{p}{q}$ is in its simplest form, e.g. $K_{16;5}^s$ or $K_{6;2}^s$. Observe that in $K_{p;q}^s$ if we apply a switching at a vertex $i, i \leq \frac{p}{2} - 1$, then we get a copy of the vertex $i + \frac{p}{2}$. Furthermore, $K_{p;q}^s = \text{DSG}(\hat{K}_{p;q}^s)$. Thus $\hat{K}_{p;q}^s$ is a homomorphic image of $K_{p;q}^s$ with respect to the switching homomorphism. Moreover, with the same assumption on p and q, $\hat{K}_{p;q}^s$ is a core.

The next lemma is a straightforward consequence of the transitivity of the homomorphism relation.

Lemma 2.1. If $(G, \sigma) \to (H, \pi)$, then $\chi_c(G, \sigma) \leq \chi_c(H, \pi)$.

Let D be a digon. It follows immediately that every signed bipartite graph admits an edge-sign preserving homomorphism to D and, as $\chi_c(D) = 4$, we have an upper bound of 4 for the circular chromatic number of signed bipartite graphs. However, the restriction of the problem to this subclass of signed graphs is still of high interest as shown by the following result of [18].

Theorem 2.2. [18] Given a graph G, we have

$$\chi_c(S(G)) = 4 - \frac{4}{\chi_c(G) + 1}.$$

That is equivalent to: $\chi_c(G) = \frac{\chi_c(S(G))}{4 - \chi_c(S(G))}$. In particular, we have that

- $\chi_c(G) \leq 4$ if and only if $\chi_c(S(G)) \leq \frac{16}{5}$,
- $\chi_c(G) \leq 3$ if and only if $\chi_c(S(G)) \leq 3$.

In the next section, we study the restriction of the circular chromatic number to the class of signed bipartite graphs and especially we introduce the signed bipartite circular clique.

2.1 Signed bipartite circular clique

One may view the class of signed circular cliques $K_{p;q}^s$ or $\widehat{K}_{p;q}^s$ as a representation of rational numbers in the homomorphism order of the class of all signed graphs. Then the circular chromatic number of a signed graph (G, σ) is determined by the first element of this chain (representing rational numbers) which is larger than (G, σ) with respect to the homomorphism order.

In Theorems 1.4, 1.5 and 2.2, we have seen the importance of the restriction of the study into the subclass SB of signed bipartite graphs. A natural question to ask is if the homomorphism order restricted to this subclass behaves similarly? More precisely, we would like to know if there is a chain of signed bipartite graphs in the homomorphism order on SB which plays the role of circular clique?

We note that no signed circular clique $\widehat{K}_{p;q}^s$ or $K_{p;q}^s$ is bipartite. Indeed each vertex in any of these cliques has a negative loop on it. In this section, for $\frac{p}{q} \leq 4$, we introduce a bipartite subgraph of these circular cliques that plays the role of circular clique in the restricted class SB.

Definition 2.3. Given a rational number $\frac{p}{q}$ where p is an even number, $2 \leq \frac{p}{q} \leq 4$ and subject to these conditions $\frac{p}{q}$ is in its simplest form, we define the signed graph $B_{p;q}$ to be the following subgraph of $K_{p;q}^s$: The vertex set $[p] = \{0, 1, \dots, p-1\}$ is partitioned to two parts X and Y where $X = \{0, 2, \dots, p-2\}$ and $Y = \{1, 3, \dots, p-1\}$. The edge set is formed by the edges of $K_{p;q}^s$ which have exactly one endpoint in X and the other endpoint in Y. The signs of edges are also induced by $K_{p:a}^s$.

We will show that $B_{p;q}$, which itself is a signed bipartite graph, plays the role of circular clique in the subclass of signed bipartite graphs. However, this class of signed graphs is partitioned into two subclasses depending on whether p is a multiple of 4 or it is 2 (mod 4).

When p is a multiple of 4, then we will show that $B_{p;q}$ is a circular clique with respect to the edge-sign preserving homomorphism. It means that any signed bipartite graph of circular chromatic number at most $\frac{p}{q}$ admits an edge-sign preserving homomorphism to $B_{p;q}$. In this case, as we will show, the subgraph induced on the vertices $\left[\frac{p}{2}\right] = \{0, 1, \dots, \frac{p}{2} - 1\}$ forms the core of $B_{p;q}$ and will play the role of signed bipartite clique with respect to the switching homomorphism. For example, $B_{16;5}$ is depicted in Figure 1 and its switching core, which is a signed graph on $K_{4,4}$, is depicted in Figure 9.



Figure 2: $B_{10:3}$

When $p \equiv 2 \pmod{4}$, and again noting our assumption that $\frac{p}{q}$ is in its simplest form subject to p being even, the signed graph $B_{p;q}$ is already a core with respect to the switching homomorphism. For example, $B_{10:3}$ is depicted in Figure 2. In this case then to have a signed circular clique with respect to the edge-sign preserving homomorphism, we must consider $DSG(B_{p;q})$. To be more precise, when $p \equiv 2 \pmod{4}$, for a signed bipartite graph (G, σ) to satisfy that $\chi_c(G, \sigma) \leq \frac{p}{q}$, it is necessary and sufficient that (G, σ) admits a switching homomorphism to $B_{p;q}$. However, for some choices of σ , this homomorphism might not be an edge-sign preserving homomorphism and a switching might be necessary. To be sure to have an edge-sign preserving homomorphism then we must consider $DSG(B_{p;q})$. For the example of p = 6 and q = 2, which corresponds to circular chromatic number at most 3, see Figures 3, 4, 5 and 6. The first one, the signed graph of Figure 3 on three vertices, is the signed circular 3-clique with respect to the switching homomorphism. The second one, the signed graph of Figure 4, which is the Double Switch Graph of the first one, is the signed circular 3-clique with respect to the edge-sign preserving homomorphism. The third one, the signed graph of Figure 5, also on 6 vertices, is the signed bipartite circular 3-clique with respect to the switching homomorphism. Finally the last one, the signed graph of Figure 6, on 12 vertices, is the Double Switch Graph of the previous one and is the signed bipartite circular 3-clique with respect to the edge-sign preserving homomorphism.

To distinguish which of the two notions of homomorphisms we are working with, we may define $B_{v;a}^s$ and $\widehat{B}^s_{p;q}$ as follows.

Given a positive even integer p and a positive integer q such that subject to p being even, $\frac{p}{q}$ is in its simplest form and $\frac{p}{q} \geq 2$, we define $B_{p;q}^s$ to be $B_{p;q}$ when $4 \mid p$ and to be $DSG(B_{p;q})$ when $4 \nmid p$. As mentioned before, these signed graphs $B_{p;q}^s$ play the role of signed bipartite circular clique with respect



to the edge-sign preserving homomorphism. For the switching homomorphism, we define $\widehat{B}_{p;q}^s$ to be $B_{p;q}$ when $4 \nmid p$ and to be the subgraph of $B_{p;q}$ induced on the vertices $\{0, \ldots, \frac{p}{2} - 1\}$ when $4 \mid p$.

We should also note that in defining $K_{p;q}^s$ and $\widehat{K}_{p;q}^s$, we did not need to assume $\frac{p}{q}$ is in the simplest form, however, we note that $K_{ap;aq}^s$ and $\widehat{K}_{ap;aq}^s$ map, respectively, to $K_{p;q}^s$ and $\widehat{K}_{p;q}^s$. By taking such a homomorphism and then taking the pre-image of $B_{p;q}^s$ and $\widehat{B}_{p;q}^s$, one may define $B_{ap;aq}^s$ and $\widehat{B}_{ap;aq}^s$.

That $B_{p;q}^s$ and $\widehat{B}_{p;q}^s$ play the roles of circular cliques in the subclass of signed bipartite graphs is the subject of the next theorem. For simplicity, it is stated using $B_{p;q}$ and switching homomorphism but one can easily restate it by using $B_{p;q}^s$ or $\widehat{B}_{p;q}^s$ and the associated notion of homomorphisms.

Theorem 2.4. Given a signed bipartite graph (G, σ) and a rational number $\frac{p}{q}$ in [2,4] where p is a positive even integer and subject to this, $\frac{p}{q}$ is in its simplest form, we have

$$\chi_c(G,\sigma) \leq \frac{p}{q}$$
 if and only if $(G,\sigma) \to B_{p;q}$.

Proof. Let (G, σ) be a signed bipartite graph. One direction is quite trivial. As $B_{p;q}$ is a subgraph of $K_{p;q}^s$, $\chi_c(B_{p;q}) \leq \frac{p}{q}$. If $(G, \sigma) \to B_{p;q}$, then, by Lemma 2.1, we have $\chi_c(G, \sigma) \leq \frac{p}{q}$. It remains to show that if $\chi_c(G, \sigma) \leq \frac{p}{q}$, then $(G, \sigma) \to B_{p;q}$. Since $B_{p;q}$ behaves differently depending

It remains to show that if $\chi_c(G, \sigma) \leq \frac{p}{q}$, then $(G, \sigma) \to B_{p;q}$. Since $B_{p;q}$ behaves differently depending on whether p divides 4 or not, we divide the proof into two cases based on this criteria: (1) p = 4k. (2) p = 4k + 2. We note that in the first case, q must be an odd number.

Case 1 p = 4k.

As $\frac{p}{q} \geq 2$, we know q is an odd number smaller than 2k - 1. Let (X, Y) be the bipartition of $B_{4k;q}$ and let (A, B) be the bipartition of (G, σ) . Since $\chi_c(G, \sigma) \leq \frac{4k}{q}$, there is an edge-sign preserving homomorphism of (G, σ) to $K^s_{4k;q}$. Let φ be such a homomorphism. Our goal is to modify φ , if needed, so that we obtain a mapping of (G, σ) to $B_{4k;q}$. This would of course be based on the bipartition of G. One such modification is given as follows:

$$\phi(u) = \begin{cases} \varphi(u) + 1 & \text{either } u \in A \text{ and } \varphi(u) \in Y \text{ or } u \in B \text{ and } \varphi(u) \in X, \\ \varphi(u) & \text{otherwise.} \end{cases}$$

Intuitively, we aim at modifying the mapping such that the vertices in the part A of G are mapped to the vertices in the part X of $B_{4k;q}$ and the vertices in the part B are mapped to the vertices in the part Y. In defining ϕ , for vertices of G satisfying these conditions under the mapping φ , we give the same image under ϕ . If this condition is not met, then we shift the image by 1 in the clockwise direction of the circle.

What remains is to show that ϕ is also an edge-sign preserving homomorphism of (G, σ) to $K^s_{4k;q}$. Then it would naturally be a homomorphism of (G, σ) to $B_{4k;q}$ as well.

Given an edge e = uv of G, if both $\phi(u) = \varphi(u)$ and $\phi(v) = \varphi(v)$ hold, then e is already mapped to an edge of the same sign under φ and nothing is left to show. If $\phi(u) = \varphi(u) + 1$ and $\phi(v) = \varphi(v) + 1$, then the claim follows from the circular structure of $K_{p;q}^s$, that is, if there is an edge ij of sign η in $B_{4k;q}$, then there is also an (i + 1)(j + 1) (additions done modulo 4k) edge of sign η . It remains to consider the case that only one endpoint of e = uv has been shifted. By the symmetry, we may assume $\phi(u) = \varphi(u)$ and $\phi(v) = \varphi(v) + 1$. Moreover, noting that u and v must be in different parts of the bipartite graph G, and again by the symmetry, we assume $u \in A$ and $v \in B$ with $\varphi(u), \varphi(v) \in X$. Hence, by our assumption, $\phi(u) = \varphi(u)$ and $\phi(v) = \varphi(v) + 1 \in Y$. Depending on the signature of e, we consider two cases.

If e is a positive edge, then $\varphi(u)\varphi(v)$ is a positive edge of $K_{4k;q}^s$. Thus $q \leq |\varphi(u) - \varphi(v)| \leq p - q = 4k - q$. Observe that, as $\varphi(u)$ and $\varphi(v)$ are both in X, they have the same parity, and thus $|\varphi(u) - \varphi(v)|$ is an even number. However, since q is an odd number, both sides of the inequality (i.e., q and 4k - q) are odd numbers and, therefore, equality cannot hold there. It is implied that if we change (only) one of $\varphi(u)$ and $\varphi(v)$ by a value of at most 1, then the inequality would still hold. Thus $\phi(u)\phi(v)$ is a positive edge of $K_{4k:q}^s$.

If e is a negative edges, then (only) one of the following must hold: either $|\varphi(u) - \varphi(v)| \leq \frac{p}{2} - q = 2k - q$ or $|\varphi(u) - \varphi(v)| \geq \frac{p}{2} + q = 2k + q$. As in the previous case, we conclude that $|\varphi(u) - \varphi(v)|$ is an even number. However, $\frac{p}{2} = 2k$ is an even number while q must be an odd number. Thus both of 2k - q and 2k + q are odd numbers and once again the equality cannot hold. Therefore, after shifting only one of the values of $\varphi(u), \varphi(v)$ by 1, the corresponding inequality holds with respect to the new function which is ϕ , that is to say, either $|\phi(u) - \phi(v)| \leq 2k - q$ or $|\phi(u) - \phi(v)| \geq 2k + q$. Hence e is mapped to a negative edge $\phi(u)\phi(v)$ of $K_{4k;a}^s$.

Case 2 p = 4k + 2.

Notice that in this case, $\frac{p}{2} = 2k + 1$ is an odd number. Let (X, Y) be the bipartition of $B_{4k+2;q}$ and let (A, B) be a bipartition of (G, σ) .

Since $\chi_c(G, \sigma) \leq \frac{4k+2}{q}$, there exists an edge-sign preserving homomorphism of (G, σ) to $K^s_{4k+2;q}$, say φ . Our goal is to modify φ to obtain a switching homomorphism of (G, σ) to $B_{4k+2;q}$. This would be based on the bipartition of G. Intuitively, we want a mapping that maps vertices in A to X and those in B to Y. We observe that for each pair of antipodal vertices of $K^s_{4k+2;q}$, one is in X and the other is in Y. Thus in the mapping φ , if one vertex is not mapped to the correct part, then we first apply a switching at that vertex and then map it to the antipodal of the original image. This is formalized as follows.

$$\phi(u) = \begin{cases} \overline{\varphi(u)} \text{ (switching at } u) & \text{ either } u \in A \text{ and } \varphi(u) \in Y \text{ or } u \in B \text{ and } \varphi(u) \in X, \\ \varphi(u) & \text{ otherwise.} \end{cases}$$

What remains is to show that ϕ is a switching homomorphism of (G, σ) to $K^s_{4k+2;q}$. Then it would naturally be a homomorphism of (G, σ) to $B_{4k+2;q}$ as well.

Given an edge e = uv of G, if both $\phi(u) = \varphi(u)$ and $\phi(v) = \varphi(v)$ hold, then it follows easily that $\phi(u)\phi(v)$ is the required edge. If $\phi(u) = \overline{\varphi(u)}$ and $\phi(v) = \overline{\varphi(u)}$, then we switch at both of vertices u and v. Thus the sign of uv does not change. Moreover, vertices i and j are connected by an edge of sign η in $K^s_{4k+2;q}$, then their antipodals are also connected by an edge of the same sign. Therefore, $\phi(u)\phi(v)$ is an edge of $K^s_{4k+2;q}$ with the same sign as $\varphi(u)\varphi(v)$ and thus as uv. The final case is that only one endpoint of uv has been switched and mapped to the antipodal. By the symmetry, we may assume that we switch at v and $\phi(v) = \overline{\varphi(v)}$. Moreover, noting that u and v must be in different parts of the bipartite graph G, and again by the symmetry, we assume $u \in A$ and $v \in B$ with $\varphi(u), \varphi(v) \in X$. Hence, by our assumption, $\phi(u) = \varphi(u)$ and $\phi(v) = \overline{\varphi(v)} \in Y$. Depending on the sign of e, we consider two cases.

If e is a positive edge, then $\varphi(u)\varphi(v)$ is a positive edge of $K_{4k+2;q}^s$. Thus $q \leq |\varphi(u) - \varphi(v)| \leq p - q = 4k + 2 - q$. As $|\varphi(v) - \overline{\varphi(v)}| = \frac{p}{2} = 2k + 1$, we have $|\varphi(u) - \overline{\varphi(v)}| \leq 2k + 1 - q$ or $|\varphi(u) - \overline{\varphi(v)}| \geq 2k + 1 + q$. Note that now uv is a negative edge of (G, σ') where σ' is obtained from σ by switching at v. Since

 $\varphi(u)\overline{\varphi(v)}$ satisfies the condition for being a negative edge of $K^s_{4k+2;q}$, $\phi(u)\phi(v)$ is a negative edge that we required.

If e is a negative edge, then (only) one of the following must hold: either $|\varphi(u) - \varphi(v)| \leq \frac{p}{2} - q = 2k + 1 - q$ or $|\varphi(u) - \varphi(v)| \geq \frac{p}{2} + q = 2k + 1 + q$. As in the previous case, we have that $|\varphi(v) - \overline{\varphi(v)}| = \frac{p}{2} = 2k + 1$. Thus $q \leq |\varphi(u) - \overline{\varphi(v)}| \leq 4k + 2 - q$. Switching at v makes uv become a positive edge. Now $\varphi(u)\overline{\varphi(v)}$ satisfies the condition for being a positive edge of $K_{4k+2;q}^s$, in other words, $\phi(u)\phi(v)$ is a positive edge. Therefore, we verify that ϕ is a switching homomorphism of (G, σ) to $K_{4k+2;q}^s$ and thus also to its signed bipartite subgraph $B_{4k+2;q}$.

We note that the assumption $\frac{p}{q} \leq 4$ is not used explicitly in the proof. If $\frac{p}{q} > 4$, then the signed bipartite graph induced by odd versus even vertices will contain a digon which admits a homomorphism from any signed bipartite graph and provides the upper bound of 4 for the circular chromatic number of this class of signed graphs. That leaves us with circular 3-coloring as a special case. In this case, by switching at all vertices of one part of $B_{6;2}$, we get an isomorphic copy of $(K_{3,3}, M)$. Hence, as a special case we have:

Corollary 2.5. Given a signed bipartite graph (G, σ) ,

$$\chi_c(G,\sigma) \leq 3$$
 if and only if $(G,\sigma) \to (K_{3,3},M)$.

Another special case is when p = 4k and q = 2k - 1. In this case, one may observe that the (switching) core of $B_{4k;2k-1}$ (on 2k vertices) is switching equivalent to the negative cycle C_{-2k} . Hence, we have the following corollary.

Corollary 2.6. Given a signed bipartite graph (G, σ) ,

$$\chi_c(G,\sigma) \leq \frac{4k}{2k-1}$$
 if and only if $(G,\sigma) \to C_{-2k}$.

This is analogous to circular $\frac{2k+1}{k}$ -colorings of graphs because $\chi_c(G) \leq \frac{2k+1}{k}$ if and only if $G \to C_{2k+1}$. Viewing a graph as a signed graph where all edges are positive, the two can be combined to the following.

Theorem 2.7. Given a positive integer ℓ , $\ell \geq 2$, and a signed graph (G, σ) satisfying $g_{ij}(G, \sigma) \geq g_{ij}(C_{-\ell})$ for each $ij \in \mathbb{Z}_2^2$, we have

$$\chi_c(G, -\sigma) \leq \frac{2\ell}{\ell - 1}$$
 if and only if $(G, \sigma) \to C_{-\ell}$.

One of the most important conjectures in the study of the circular chromatic numbers of planar graphs is the well-known Jaeger-Zhang conjecture which claims that every planar graph of odd-girth at least 4k + 1 admits a homomorphism to C_{2k+1} , i.e., a circular $\frac{2k+1}{k}$ -coloring. Attempting to fill the gaps in the statement of the conjecture and potentially provide a better room for induction on k, we propose the following strengthening.

Conjecture 2.8. Given a positive integer $k, k \geq 2$, any signed planar graph (G, σ) satisfying $g_{ij}(G, -\sigma) \geq g_{ij}(C_{-(2k-r)})$ where $r \in \{0, 1\}$ has circular chromatic number at most $\frac{2k}{k-1}$.

The case (k, r) = (2, 0) is trivial as every signed bipartite graph admits a circular 4-coloring. The case (k, r) = (2, 1) is the 4-Color Theorem. The case (3, 1) is the well-known Grötzsch theorem. The case (k, r) = (2p + 1, 1) is the Jaeger-Zhang conjecture. Theorem 1.8 proves the case (k, r) = (3, 0) of this conjecture. More details are given in Section 3.2. For some supporting evidence we refer to [8].

2.2 Circular coloring and subdivision

A classic relation between the chromatic number of a graph and homomorphism from a certain subdivisions of it to the odd cycle is extended, in [14], to a relation between the circular chromatic number of signed graphs and homomorphism of its subdivision to negative cycles. Here we present a slightly stronger version and then use it to build examples in the next sections.

Definition 2.9. Given a signed graph (G, σ) and a positive integer ℓ , we define $T^*_{\ell}(G, \sigma)$ to be the signed graph obtained from (G, σ) by replacing each edge e with a path P_{ℓ} of length ℓ where internal vertices of the path are disjoint and assigning a signature satisfying that P_{ℓ} contains an odd number of positive edges if e is a positive edge and P_{ℓ} contains an even number of positive edges if e is a negative edge.

We note that there are many choices for the signature in defining $T_{\ell}^*(G, \sigma)$, but, as all such choices are switching equivalent, one may take any. The relation between the circular chromatic number of (G, σ) and $T_{\ell}^*(G, \sigma)$ follows from two lemmas based on the following notation of indicator.

Given a signed graph I with two specific vertices u and v, we refer to $\mathcal{I} = (I, u, v)$ as an indicator. Given an indicator \mathcal{I} and a real number $r, r \geq 2$, with [0, r) viewed as the circle of circumference r, we define $Z(\mathcal{I})$ to be the set of possible choices for u in a circular r-coloring of \mathcal{I} where v is colored by 0.

Given two indicators $\mathcal{I}_+ = (I_1, u_1, v_2)$ and $\mathcal{I}_- = (I_2, u_2, v_2)$, for each signed graph Ω , we define $\Omega(\mathcal{I}_+, \mathcal{I}_-)$ to be the signed graph obtained from Ω by replacing each positive edge xy with a distinct copy of \mathcal{I}_+ where x is identified with u_1 and y with u_2 and similarly replacing each negative edge with \mathcal{I}_- . For some indicators, the circular chromatic number of $\Omega(\mathcal{I}_+, \mathcal{I}_-)$ could be determined by $\chi_c(\Omega)$.

Lemma 2.10. [18] Assume \mathcal{I}_+ and \mathcal{I}_- are two signed indicators, $r \geq 2$ is a real number such that $Z(\mathcal{I}_+) = [t, \frac{r}{2}]$ and $Z(\mathcal{I}_-) = [0, \frac{r}{2} - t]$ for some $0 < t < \frac{r}{2}$. Then for any signed graph Ω , we have

$$\chi_c(\Omega(\mathcal{I}_+, \mathcal{I}_-)) = t\chi_c(\Omega).$$

We denote a path of length ℓ which contains an odd number of positive edges by P_{ℓ}^{o} and a path of length ℓ which contains an even number of positive edges by P_{ℓ}^{e} . The special choice for the indicators are P_{ℓ}^{o} and P_{ℓ}^{e} , with the two endpoints as special vertices. The range of possible choices for the ends in the circular colorings of these paths is computed in [22].

Lemma 2.11. [22] Given an integer $\ell \geq 1$ and a real number $r < \frac{2\ell}{\ell-1}$,

$$Z(P_{\ell}^{e}) = [0, \ell \frac{r}{2} - \ell] \text{ and } Z(P_{\ell}^{o}) = [\ell - (\ell - 1)\frac{r}{2}, \frac{r}{2}].$$

Combining these two lemmas, where we take $\mathcal{I}_+ = P_{\ell}^o, \mathcal{I}_- = P_{\ell}^e$ and $t = \ell - (\ell - 1)\frac{r}{2}$, we have the following.

Lemma 2.12. For any signed graph Ω ,

$$\chi_c(T^*_{\ell}(\Omega)) = \frac{2\ell\chi_c(\Omega)}{(\ell-1)\chi_c(\Omega)+2}.$$

A few comments are to be mentioned here.

The first is to note that for each positive integer ℓ , and by considering a graph G as a signed graph where all edges are regarded positive, the 4-colorability of G is equivalent to proving that $\chi_c(T_\ell^*(G)) \leq \frac{8\ell}{4\ell-2}$. Furthermore, noting that subdivision preserves the planarity, for each choices of ℓ , we have a reformulation of the 4-Color Theorem. For each such ℓ then one line of study is to introduce an interesting classes of signed graphs that includes $T_\ell^*(G)$ for all planar graphs G and admits the same upper bound for the circular chromatic number.

A second note here is that for even values of ℓ , $T^*_{\ell}(\Omega)$ is a signed bipartite graph and thus the subject of the main study in this work.

And the last note is that the S(G) construction mentioned before is a special case of this indicator construction. Given a graph G, one may first build a signed graph \tilde{G} by replacing each edge with a digon. Then $T_2^*(\tilde{G})$ is the same as S(G).

3 Coloring planar signed graphs

We have already noted that, via constructions such as S(G), most homomorphisms and coloring questions can be restated in the language of homomorphisms of singed bipartite graphs. Here we have a look at what this means to the coloring of planar graphs and restate some famous theorems such as the 4-Color Theorem and Grötzsch's theorem.

As a direct corollary of Theorem 1.4, we have the following reformulation of the 4-Color Theorem.

Theorem 3.1. [4-Color Theorem restated] For any planar graph G, we have $S(G) \to S(K_4)$.

Noting that the 4-Color Theorem is equivalent to bounding the circular chromatic number of all planar graphs by 4, and applying Theorem 2.2, another restatement of 4-Color Theorem is as follows.

Theorem 3.2. [4-Color Theorem restated] For any planar graph G, we have $\chi_c(S(G)) \leq \frac{16}{5}$.

Since every S(G) is a signed bipartite graph, the claim of this theorem is equivalent to the existence of an edge-sign preserving mapping from S(G) to $B_{16;5}^s$. Note that $\widehat{B}_{16;5}^s$, the switching core of $B_{16;5}^s$, is a signed graph on $K_{4,4}$. With one random choice of a signature, (among all equivalent signatures), this core is presented in Figure 9. We recall that the 4-Color Theorem is also restated in Theorem 1.6 in the form of mapping S(G), for every planar G, to $(K_{4,4}, M)$.

Let \mathcal{P}' be the class of all simple planar graphs and let $S(\mathcal{P}') = \{S(G) : G \in \mathcal{P}'\}$. Then, by the discussion above, the 4-Color Theorem is equivalent to bounding the class $S(\mathcal{P}')$ by either of the signed bipartite graphs of Figures 7, 8, or 9. One may observe that $S(K_4)$ admits a homomorphism to both $(K_{4,4}, M)$ and $\widehat{B}^s_{16;5}$ but $(K_{4,4}, M)$ and $\widehat{B}^s_{16;5}$ are homomorphically incomparable. The latter is a consequence of the following two facts: 1. Any pair of nonadjacent vertices in $(K_{4,4}, M)$ or in $\widehat{B}^s_{16;5}$ belongs to a negative 4-cycle which means identifying them would result in a digon; 2. the two signed graphs are not switching isomorphic, for example, $\chi_c(K_{4,4}, M) = 4$ and $\chi_c(\widehat{B}^s_{16;5}) = \frac{16}{5}$.



Observe that $S(K_4)$ itself is in the family $S(\mathcal{P}')$ and bounds the family. By taking a larger class of 4-colorable graphs in place of \mathcal{P}' , such as the class of K_5 -minor-free graphs we may strengthen the result, but we do not expect further extension. In contrast, since $S(K_4)$ admits a homomorphism to each of $(K_{4,4}, M)$ and $\hat{B}^s_{16;5}$, it would not be a surprise if a stronger statement can be proved regarding these two targets. Indeed that is the case for $(K_{4,4}, M)$: it bounds the class of all signed bipartite planar simple graphs [15]. As the limit of the circular chromatic numbers of signed bipartite planar simple graphs is 4 (see [18] and [7]), this cannot be the case for $\hat{B}^s_{16;5}$. Thus it remains an open question to bound a larger class of signed bipartite planar graphs with $\hat{B}^s_{16;5}$.

It is shown in [14] that every signed bipartite planar graph of negative girth at least 8 maps to C_{-4} and that this girth condition cannot be improved to 6. We observe that C_{-4} is a subgraph of each of the three homomorphism targets of this discussion.

Another common subgraph of $(K_{4,4}, M)$ and $\widehat{B}_{16;5}^s$ which is of high interest for this discussion is $(K_{3,3}, M)$. A restatement of the Grötzsch theorem is the following.

Theorem 3.3. [Grötzsch's theorem restated] For any triangle-free planar graph G (with no loop), we have $\chi_c(S(G)) \leq 3$.

In the next subsection, we prove Theorem 1.8 which may be viewed as a parallel theorem to Grötzsch's theorem. In Section 4 then we propose a question as potentially common strengthening of the two theorems.

3.1 Bounding the circular chromatic number by 3

For a class \mathcal{C} of signed graphs, we define $\chi_c(\mathcal{C}) = \sup\{\chi_c(G,\sigma) : (G,\sigma) \in \mathcal{C}\}.$

For a given integer k, let \mathcal{P}_k^* be the class of singed planar graphs (G, σ) such that the signed graph $(G, -\sigma)$ satisfies the following conditions: for each $ij \in \mathbb{Z}_2^2$, we have $g_{ij}(G, -\sigma) \geq g_{ij}(C_{-k})$. Thus for an odd integer k, and after suitable switchings, \mathcal{P}_k^* consists of all planar graphs of odd girth at least k with all edges being assigned positive signs. For an even value of k, the class \mathcal{P}_k^* consists of all signed bipartite planar graphs of negative girth at least k.

A main question then is to find $\chi_c(\mathcal{P}_k^*)$ for each k. For k = 3 and 4 both answers are 4, the first by the 4-Color Theorem, the second by the observation that 4 is the upper bound for the class of signed bipartite simple graphs and a construction given in [18] showing that 4 cannot be improved (see also [7]). For k = 5, we have the Grötzsch theorem, that gives upper bound of 3 which is also shown to be the optimal value. For k = 4p + 1, this question is the subject of widely studied Jaeger-Zhang conjecture. And for other values of k, similar conjectures are proposed. Here, addressing the case k = 6 we prove the following result.

Theorem 3.4. We have $\frac{14}{5} \le \chi_c(\mathcal{P}_6^*) \le 3$.

The proof of the upper bound is based on the following theorem which is implied by combining several results from the literature, noting that the 4-Color Theorem is used in proving this claim.

Theorem 3.5. Given a signed bipartite planar graph (G, σ) of negative girth at least 6, one can find six disjoint subsets of edges, E_1, E_2, \ldots, E_6 , such that each of the signed graphs (G, σ_i) , $i \in \{1, 2, \ldots, 6\}$, where E_i is the set of negative edges of (G, σ_i) , is switching equivalent to (G, σ) .

To follow the literature for a proof of this claim, one first should note that in [15] it is shown that the claim of Theorem 3.5 is equivalent to the following: Given a 6-regular planar multigraph G, if for every set X of odd number of vertices there are at least six edges in the edge-cut $(X, V(G) \setminus X)$, then G is 6-edge-colorable. Replacing 6 with a general integer k is the subject of a conjecture of P. Seymour while the case k = 3 is a classic restatement of the 4-Color Theorem. The proof of the case k = 6 is given in [3]. However, this proof is based on induction on k, thus not only relies on the 4-Color Theorem, but also on the proof of the cases k = 4 and k = 5 of Seymour's conjecture. Proofs of these two cases were claimed by B. Guenin in 2003, but there has been no publication of it since then. However, these cases are verified independently through the notion of packing signatures in signed graphs in [19] and [20].

Proof of Theorem 3.4. Let (G, σ) be a signed bipartite planar graph of negative girth at least 6 with a bipartition (A, B). By Theorem 3.5, there are disjoint subsets E_1, E_2, \ldots, E_6 of edges of G such that for each $i \in [6]$, the signature σ_i , whose negative edges are E_i , is equivalent to σ .

We consider the signed graph (G, σ_1) where the set of negative edges is E_1 . Let G' be the graph obtained from G by contracting all the edges in E_1 . In this notion of contracting, we delete the contracted edge (those in E_1) but all other edges remain. Thus in theory we may have loops and parallel edge in the resulting graph. However, we show next that not only G' has no loop, it has no triangle either. In other words, we claim that every odd cycle of G' is of length at least 5.

To see this, let C' be an odd cycle of G'. This cycle is obtained from a cycle C of G by contracting some edges (of E_1). As G is bipartite, C must be of even length. Thus the number of the contracted edges is odd. Therefore, C is a negative cycle in the signed graph (G, σ_1) . As all the (G, σ_i) , $i = 1, 2, \ldots, 6$, are equivalent, C is negative in all of them which means it has an odd number of edges from each of E_i 's. As these sets are disjoint, and as for i = 2, 3, ... 6, they still present in G', the cycle C' has an odd number of edges from each E_i , i = 2, 3, ... 6. In particular, that is at least one edge from each, and noting again that they are disjoint sets, we conclude that C' is of length at least 5.

Having shown that G' is a triangle-free planar graph with no loop (might have parallel edges), we may apply the Grötzsch theorem to obtain a 3-coloring $\varphi : V(G') \to \{1, 2, 3\}$ of G'. Let (X, Y) be the bipartition of $(K_{3,3}, M)$. Label the vertices $X = \{x_1, x_2, x_3\}$ and $Y = \{y_1, y_2, y_3\}$ such that $\{x_1y_1, x_2y_2, x_3y_3\}$ is the set of negative edges.

We define the mapping ψ of (G, σ_1) to $(K_{3,3}, M)$ as follows:

$$\psi(u) = \begin{cases} x_i, & \text{if } u \in A \text{ and } \varphi(u) = i \\ y_i, & \text{if } u \in B \text{ and } \varphi(u) = i. \end{cases}$$

It remains to show that ψ is an edge-sign preserving mapping of (G, σ_1) to $(K_{3,3}, M)$. For any positive edge uv of (G, σ_1) , without loss of generality, we may assume that $u \in A, v \in B$ and that $\varphi(u) = i$, $\varphi(v) = j$. Noting that uv is also an edge of G', as φ is a proper 3-coloring, we have that $i \neq j$. Thus $\psi(u)\psi(v) = x_ix_j$ is a positive edge in $(K_{3,3}, M)$. For any negative edge uv of (G, σ_1) , without loss of generality, assume $u \in A, v \in B$. As $uv \in E_1$ is contracted to a vertex to obtain G', $\varphi(u) = \varphi(v) = i$. So $\psi(u)\psi(v) = x_iy_i$ is a negative edge. Hence, ψ is an edge-sign preserving homomorphism of (G, σ_1) to $(K_{3,3,M})$. This completes the proof of the upper bound.

For the lower bound we recall that an example of a signed simple planar graph (G, σ) satisfying $\chi_c(G, \sigma) = \frac{14}{3}$ is given in [18]. Then it follows from Lemma 2.12 that $\chi_c(T_2^*(G, \sigma)) = \frac{14}{5}$. It is easily observed that, since (G, σ) is a signed simple planar graph, $T_2^*(G, \sigma)$ has (negative) girth at least 6 and obviously it is a signed bipartite graph.

3.2 Bounds based on girth

We will denote the class of all signed planar graphs with \mathcal{P} , where we will allow loops and multi-edges. The subclass of \mathcal{P} where the shortest cycle of each member is at least k will be denoted by \mathcal{P}_k . Thus, in particular, \mathcal{P}_2 is the class of all loop-free signed planar graphs and \mathcal{P}_3 is the class of all signed planar simple graphs.

We note that \mathcal{P}_k^* is not a subclass of \mathcal{P}_k as signed graphs in \mathcal{P}_k^* may have positive even cycles of any length. However, we expect that the circular chromatic number of \mathcal{P}_k^* is determined by the subclass $\mathcal{P}_k^* \cap \mathcal{P}_k$.

The questions of determining $\chi_c(\mathcal{P}_k^*)$ is closely related to some of the most well known theorems and conjectures in the theory of graph coloring, such the 4-Color theorem, Grötzsch's theorem and Jaeger-Zhang conjecture. This also leads to the importance of the question of determining $\chi_c(\mathcal{P}_k)$. In the table below we summarize the best known results for these questions for various values of k.

Circular chromatic number of \mathcal{P}_k^* and \mathcal{P}_k

k	Bounds on $\chi_c(\mathcal{P}_k^*)$	Reference	Bounds on $\chi_c(\mathcal{P}_k)$	Reference
2	$\chi_c(\mathcal{P}_2^*) = 4$	[Bipartite]	$\chi_c(\mathcal{P}_2) = 8$	[4CT]
3	$\chi_c(\mathcal{P}_3^*) = 4$	[4CT]	$\chi_c(\mathcal{P}_3) \le 6$	[18]
4	$\chi_c(\mathcal{P}_4^*) \cong 4$	[7]	$\chi_c(\mathcal{P}_4) \le 4$	[12]
5	$\chi_c(\mathcal{P}_5^*) = 3$	[5], [23]	*	
6	$\chi_c(\mathcal{P}_6^*) \leq 3$	[this paper]	*	
7	*		$\chi_c(\mathcal{P}_7) \le 3$	[13]
8	$\chi_c(\mathcal{P}_8^*) \cong \frac{8}{3}$	[14]	*	
11	$\chi_c(\mathcal{P}_{11}^*) \le \frac{5}{2}$	[4], [2]	*	
14	$\chi_c(\mathcal{P}_{14}^*) \le \frac{12}{5}$	[10]	*	
17	$\chi_c(\mathcal{P}_{17}^*) \le \frac{7}{3}$	[2], [21]	*	
6p - 2	$\chi_c(\mathcal{P}^*_{6p-2}) \le \frac{4p}{2p-1}$	[8]	$\chi_c(\mathcal{P}_{6p-2}) \le \frac{8p-2}{4p-3}$	[8]
6p - 1	$\chi_c(\mathcal{P}^*_{6p-1}) \le \frac{4p}{2p-1}$	[11]	$\chi_c(\mathcal{P}_{6p-1}) \le \frac{4p}{2p-1}$	[8]
6 <i>p</i>	*		$\chi_c(\mathcal{P}_{6p}) < \frac{4p}{2p-1}$	[8]
6p + 1	$\chi_c(\mathcal{P}^*_{6p+1}) \le \frac{2p+1}{p}$	[9]	$\chi_c(\mathcal{P}_{6p+1}) \le \frac{8p+2}{4p-1}$	[8]
6p + 2	*		$\chi_c(\mathcal{P}_{6p+2}) \le \frac{2p+1}{p}$	[8]
6p + 3	$\chi_c(\mathcal{P}^*_{6p+3}) < \frac{2p+1}{p}$	[11]	$\chi_c(\mathcal{P}_{6p+3}) < \frac{2p+1}{p}$	[8]

In this table, when we write $\chi_c(\mathcal{C}) = r$, it means that $\chi_c(\widehat{G}) \leq r$ for each member \widehat{G} of the class \mathcal{C} and that the equality is known to hold for at least one member of the class. When we write $\chi_c(\mathcal{C}) \cong r$, we mean that there is a sequence of signed graphs of \mathcal{C} whose limit of the circular chromatic number is r. In such cases, sometimes it is verified that the r is never reached by a single member of \mathcal{C} . For example, this is indeed the case for \mathcal{P}_4^* as shown in [7]. In other cases, it is not known if the equality holds for some members or the inequality is always strict. In particular, for \mathcal{P}_8^* the sequence that gives the limit of $\frac{8}{3}$ is $\{T_2^*(\Gamma_i)\}$ where Γ_i is the sequence reaching the limit of 4 for \mathcal{P}_4^* . It remains an open problem whether the equality can be reached in this case.

There are some trivial inclusion among the classes considered here: $\mathcal{P}_{k+2}^* \subseteq \mathcal{P}_k^*$ and $\mathcal{P}_{k+1} \subseteq \mathcal{P}_k$. In such cases, any upper bound for the larger class works also on the smaller one and any lower bound for the smaller one works on the larger one as well. In the entries of the table where we write * the best known bounds come from the other entries of the table based on these inclusion.

To tight the gap in the bounds or, more ambitiously, to determine the exact values, is the subject of some of main work in the theory of coloring planar graphs. A notable conjecture is that of Jaeger-Zhang which can be restated as:

Conjecture 3.6. [Jaeger-Zhang Conjecture] Given a positive integer p, we have $\chi_c(\mathcal{P}^*_{4p+1}) \leq \frac{2p+1}{p}$.

A bipartite analogue of this conjecture was first proposed in [16], but considering the result of [14], it is modified to the following. Recall that the class \mathcal{P}_{2k}^* consists of all signed planar bipartite graphs of negative girth at least 2k.

Conjecture 3.7. [Bipartite analogue of Jaeger-Zhang Conjecture] Given a positive integer p, we have $\chi_c(\mathcal{P}_{4p}^*) \leq \frac{4p}{2p-1}$.

As a common generalization of the two, we have proposed Conjecture 2.8.

4 Conclusions and further questions

In this paper, verifying the importance of the study of circular chromatic number of signed bipartite graphs we have presented the signed bipartite circular cliques. Then, using the 4-Color Theorem, we have shown an upper bound of 3 for the circular chromatic number of signed bipartite planar graphs of negative girth at least 6. We have provided a table summarizing the best known results on the circular chromatic number of signed planar graphs with a girth condition. Beside all the open questions that are summarized in the table, there are two questions of interest to mention.

The first is about the use the 4-Color Theorem in our proof of the upper bound of 3 for the circular chromatic number of the subclass $\chi_c(\mathcal{P}_2^*)$. Could one find a relatively short proof of this without using the 4-Color Theorem? Or can one show that, on the contrary, this result implies the 4-Color Theorem? We recall that, in Section 2, reformulations of the 4-Color Theorem using special classes of planar graphs of high girth are given. So this would not be a surprise.

The second question is to potentially strengthen our result to include the Grötzsch theorem as a special case. One possibility is observed by reformulating the Grötzsch theorem itself as follows.

Theorem 4.1. [Grötzsch's theorem restated] If G is a planar graph satisfying that $K_3 \not\rightarrow G$, then $G \rightarrow K_3$.

We recall that if $K_3 \not\rightarrow G$, then $S(K_3) \not\rightarrow S(G)$ and that if $G \rightarrow K_3$ then $S(G) \rightarrow S(K_3)$. Thus a potential strengthening of our result, which would include the Grötzsch theorem, is as follows.

Conjecture 4.2. If (G, σ) is a signed bipartite planar graph with the property that $S(K_3) \not\rightarrow (G, \sigma)$, then $\chi_c(G, \sigma) \leq 3$, i.e., $(G, \sigma) \rightarrow (K_{3,3}, M)$.

Acknowledgement. This work is supported by the French ANR project HOSIGRA (ANR-17-CE40-0022). The second author has also received funding from the European Union's Horizon 2020 research and innovation program under the Marie Sklodowska-Curie grant agreement No 754362 and the Fundamental Research Funds for the Central Universities, Nankai University.

References

- Brewster, R. C. and Graves, T. (2009). Edge-switching homomorphisms of edge-coloured graphs. Discrete Mathematics, 18(309), 5540–5546.
- [2] Cranston, D. W. and Li, J. (2020). Circular flows in planar graphs. SIAM Journal on Discrete Mathematics 34(1), 497-519.
- [3] Dvořák, Z., Kawarabayashi, K. I. and Kral. D. (2016). Packing six T-joins in plane graphs. Journal of Combinatorial Theory, Series B, 116, 287-305.
- [4] Dvořák, Z., and Postle, L. (2017). Density of 5/2-critical graphs. Combinatorica, 37(5), 863-886.
- [5] Grötzsch, H. (1959). Ein Dreifarbensatz für dreikreisfreie Netze auf der Kugel. Wiss. Z. Martin-Luther-Univ. Halle-Wittenberg Math.-Natur. Reihe, 8, pp. 109-120
- [6] Guenin, B. (2003). Packing T-joins and edge colouring in planar graphs. Manuscript.
- [7] Kardoš, F., Narboni, J., Naserasr, R., and Wang, Z. (2021). Circular (4ϵ) -coloring of some classes of signed graphs. arXiv preprint arXiv:2107.12126.
- [8] Li, J., Naserasr, R., Wang, Z., and Zhu, X. (2022). Circular flows in mono-directed signed graphs. arXiv preprint arXiv:2212.10757.
- [9] Lovász, L. M., Thomassen, C., Wu, Y., and Zhang, C. Q. (2013). Nowhere-zero 3-flows and modulo k-orientations. Journal of Combinatorial Theory, Series B, 103(5), 587-598.
- [10] Li, J., Shi, Y., Wang, Z., and Wei, C. (2023). Homomorphisms to small negative even cycles. Submitted.

- [11] Li, J., Wu, Y., and Zhang, C.-Q. (2020). Circular flows via extended Tutte orientations. Journal of Combinatorial Theory, Series B, 145, 307-322.
- [12] Máčajová, E., Raspaud, A. and Skoviera, M. (2016). The Chromatic Number of a Signed Graph. Electronic Journal of Combinatorics 23(1), 1-14.
- [13] Naserasr, R., Skrekovski, R., Wang, Z., and Xu, R. (2023). Mapping sparse signed graphs to (K_{2k}, M) . Journal of Graph Theory, (2023), 1-34, https://doi.org/10.1002/jgt.22992
- [14] Naserasr, R., Pham, L. A., and Wang, Z. (2022). Density of C₋₄-critical signed graphs. Journal of Combinatorial Theory, Series B 153: 81-104.
- [15] Naserasr, R., Rollová, E., and Sopena, E. (2013). On homomorphisms of planar signed graphs to signed projective cubes. In The Seventh European Conference on Combinatorics, Graph Theory and Applications (pp. 271-276). Edizioni della Normale, Pisa.
- [16] Naserasr, R., Rollová, E., and Sopena, É. (2015). Homomorphisms of signed graphs. Journal of Graph Theory, 79(3), 178-212.
- [17] Naserasr, R., Sopena, E., and Zaslavsky, T. (2021). Homomorphisms of signed graphs: An update. European Journal of Combinatorics, 91, 103222.
- [18] Naserasr, R., Wang, Z., and Zhu, X. (2021). Circular chromatic number of signed graphs. The Electronic Journal of Combinatorics, 28(2), P2.44.
- [19] Naserasr, R. and Yu, W. (2023). Packing signatures in signed graphs. To appear in SIAM Disc. Math. Available at: https://hal.science/hal-03960731
- [20] Naserasr, R. and Yu, W. (2023). On the packing number of antibalanced signed simple planar graphs of negative girth at least 5. https://hal.science/hal-04080407
- [21] Postle, L. and Smith-Roberge, E. (2022). On the Density of C₇-Critical Graphs. Combinatorica, 48pp. DOI:10.1007/s00493-020-4177-y
- [22] Pan, Z., and Zhu, X. (2022). The circular chromatic numbers of signed series-parallel graphs. Discrete Mathematics, 345(3), 112733.
- [23] Steinberg, R. and Younger, D.H. (1989). Grötzsch's theorem for the projective plane. Ars Combin., 28:15–31.
- [24] Zaslavsky, T. (1982). Signed graph coloring. Discrete Mathematics, 39, 215–228.
- [25] Zaslavsky, T. (1982). Signed graphs. Discrete Applied Mathematics, 4(1), 47-74.