

Numbers with integer expansion in the numeration system with negative base

Petr Ambrož

joint work with D. Dombek, Z. Masáková, and E. Pelantová

Doppler Institute and Department of Mathematics
FNSPE, Czech Technical University in Prague

Digital expansions, dynamics and tilings
Aussois, April 2010

- 1 Rényi expansions
- 2 Ito-Sadahiro expansions
- 3 $(-\beta)$ -integers
- 4 Examples
- 5 Open questions

β -expansions of real numbers

Consider $\beta > 1$ and $T_\beta : [0, 1) \mapsto [0, 1)$ given by

$$T_\beta(x) := \beta x - \lfloor \beta x \rfloor.$$

Representation of $x \in [0, 1)$ of the form

$$x = \frac{x_1}{\beta} + \frac{x_2}{\beta^2} + \frac{x_3}{\beta^3} + \dots,$$

where

$$x_i = \lfloor \beta T_\beta^{i-1}(x) \rfloor$$

is called the β -**expansion** of x .

We write

$$d_\beta(x) := x_1 x_2 x_3 \dots.$$

β -expansions of real numbers

Consider $\beta > 1$ and $T_\beta : [0, 1) \mapsto [0, 1)$ given by

$$T_\beta(x) := \beta x - \lfloor \beta x \rfloor.$$

Representation of $x \in [0, 1)$ of the form

$$x = \frac{x_1}{\beta} + \frac{x_2}{\beta^2} + \frac{x_3}{\beta^3} + \dots,$$

where

$$x_i = \lfloor \beta T_\beta^{i-1}(x) \rfloor$$

is called the **β -expansion** of x .

We write

$$d_\beta(x) := x_1 x_2 x_3 \dots.$$

β -expansions of real numbers

Consider $\beta > 1$ and $T_\beta : [0, 1) \mapsto [0, 1)$ given by

$$T_\beta(x) := \beta x - \lfloor \beta x \rfloor.$$

Representation of $x \in [0, 1)$ of the form

$$x = \frac{x_1}{\beta} + \frac{x_2}{\beta^2} + \frac{x_3}{\beta^3} + \dots,$$

where

$$x_i = \lfloor \beta T_\beta^{i-1}(x) \rfloor$$

is called the **β -expansion** of x .

We write

$$d_\beta(x) := x_1 x_2 x_3 \dots.$$

β -expansions of real numbers

The β -expansion of $x \geq 1$ can be naturally defined:

- find an exponent $k \in \mathbb{N}$ such that $\frac{x}{\beta^k} \in [0, 1)$
- using the transformation T_β derive the β -expansion of $\frac{x}{\beta^k}$

$$\frac{x}{\beta^k} = \frac{x_1}{\beta} + \frac{x_2}{\beta^2} + \frac{x_3}{\beta^3} + \dots,$$

- then

$$x = x_1\beta^{k-1} + x_2\beta^{k-2} + \dots + x_{k-1}\beta + x_k + \frac{x_{k+1}}{\beta} + \dots.$$

- does not depend on k

β -expansions of real numbers

The β -expansion of $x \geq 1$ can be naturally defined:

- find an exponent $k \in \mathbb{N}$ such that $\frac{x}{\beta^k} \in [0, 1)$
- using the transformation T_β derive the β -expansion of $\frac{x}{\beta^k}$

$$\frac{x}{\beta^k} = \frac{x_1}{\beta} + \frac{x_2}{\beta^2} + \frac{x_3}{\beta^3} + \dots,$$

- then

$$x = x_1\beta^{k-1} + x_2\beta^{k-2} + \dots + x_{k-1}\beta + x_k + \frac{x_{k+1}}{\beta} + \dots.$$

- does not depend on k

β -expansions of real numbers

The β -expansion of $x \geq 1$ can be naturally defined:

- find an exponent $k \in \mathbb{N}$ such that $\frac{x}{\beta^k} \in [0, 1)$
- using the transformation T_β derive the β -expansion of $\frac{x}{\beta^k}$

$$\frac{x}{\beta^k} = \frac{x_1}{\beta} + \frac{x_2}{\beta^2} + \frac{x_3}{\beta^3} + \dots,$$

- then

$$x = x_1\beta^{k-1} + x_2\beta^{k-2} + \dots + x_{k-1}\beta + x_k + \frac{x_{k+1}}{\beta} + \dots.$$

- does not depend on k

β -expansions of real numbers

The β -expansion of $x \geq 1$ can be naturally defined:

- find an exponent $k \in \mathbb{N}$ such that $\frac{x}{\beta^k} \in [0, 1)$
- using the transformation T_β derive the β -expansion of $\frac{x}{\beta^k}$

$$\frac{x}{\beta^k} = \frac{x_1}{\beta} + \frac{x_2}{\beta^2} + \frac{x_3}{\beta^3} + \dots,$$

- then

$$x = x_1\beta^{k-1} + x_2\beta^{k-2} + \dots + x_{k-1}\beta + x_k + \frac{x_{k+1}}{\beta} + \dots.$$

- does not depend on k

β -integers

An integer sequence

$$x_1 x_2 x_3 \cdots$$

is said to be **β -admissible** if there exists $x \in [0, 1)$

$$d_\beta(x) = x_1 x_2 x_3 \cdots .$$

Set of non-negative β -integers is

$$\mathbb{Z}_\beta^+ := \{x_k \beta^k + \cdots + x_1 \beta + x_0 \mid x_k \cdots x_0 0^\omega \text{ is a } \beta\text{-admissible sequence}\} .$$

[Thurston]: The distances between consecutive β -integers take values in $\{\Delta_i \mid i = 0, 1, 2, \dots\}$, where

$$\Delta_i = \sum_{j=1}^{\infty} \frac{t_{i+j}}{\beta^j} \quad \text{and} \quad d_\beta(1) = t_1 t_2 t_3 \cdots .$$

β -integers

An integer sequence

$$x_1 x_2 x_3 \cdots$$

is said to be **β -admissible** if there exists $x \in [0, 1)$

$$d_\beta(x) = x_1 x_2 x_3 \cdots .$$

Set of non-negative **β -integers** is

$$\mathbb{Z}_\beta^+ := \{x_k \beta^k + \cdots + x_1 \beta + x_0 \mid x_k \cdots x_0 0^\omega \text{ is a } \beta\text{-admissible sequence}\} .$$

[Thurston]: The distances between consecutive β -integers take values in $\{\Delta_i \mid i = 0, 1, 2, \dots\}$, where

$$\Delta_i = \sum_{j=1}^{\infty} \frac{t_{i+j}}{\beta^j} \quad \text{and} \quad d_\beta(1) = t_1 t_2 t_3 \cdots .$$

β -integers

An integer sequence

$$x_1 x_2 x_3 \cdots$$

is said to be **β -admissible** if there exists $x \in [0, 1)$

$$d_\beta(x) = x_1 x_2 x_3 \cdots .$$

Set of non-negative **β -integers** is

$$\mathbb{Z}_\beta^+ := \{x_k \beta^k + \cdots + x_1 \beta + x_0 \mid x_k \cdots x_0 0^\omega \text{ is a } \beta\text{-admissible sequence}\} .$$

[Thurston]: The distances between consecutive β -integers take values in $\{\Delta_i \mid i = 0, 1, 2, \dots\}$, where

$$\Delta_i = \sum_{j=1}^{\infty} \frac{t_{i+j}}{\beta^j} \quad \text{and} \quad d_\beta(1) = t_1 t_2 t_3 \cdots .$$

- 1 Rényi expansions
- 2 Ito-Sadahiro expansions**
- 3 $(-\beta)$ -integers
- 4 Examples
- 5 Open questions

$(-\beta)$ -expansions of real numbers

Consider $-\beta < -1$ and $T_{-\beta} : \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1}\right) \mapsto \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1}\right)$ given by

$$T_{-\beta}(x) := -\beta x - \left\lfloor -\beta x + \frac{\beta}{\beta+1} \right\rfloor.$$

Representation of $x \in I_\beta \equiv [l_\beta, r_\beta) \equiv \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1}\right)$ of the form

$$x = \frac{x_1}{-\beta} + \frac{x_2}{(-\beta)^2} + \frac{x_3}{(-\beta)^3} + \dots,$$

where

$$x_i = \left\lfloor -\beta T_{-\beta}^{i-1}(x) + \frac{\beta}{\beta+1} \right\rfloor,$$

is called the **$(-\beta)$ -expansion** of x , denoted $d_{-\beta}(x) = x_1 x_2 x_3 \dots$.

We denote

$$d_{-\beta}(I_\beta) = d_{-\beta}\left(\frac{-\beta}{\beta+1}\right) = l_1 l_2 l_3 \dots$$

$(-\beta)$ -expansions of real numbers

Consider $-\beta < -1$ and $T_{-\beta} : \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1}\right) \mapsto \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1}\right)$ given by

$$T_{-\beta}(x) := -\beta x - \left\lfloor -\beta x + \frac{\beta}{\beta+1} \right\rfloor.$$

Representation of $x \in I_\beta \equiv [l_\beta, r_\beta) \equiv \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1}\right)$ of the form

$$x = \frac{x_1}{-\beta} + \frac{x_2}{(-\beta)^2} + \frac{x_3}{(-\beta)^3} + \dots,$$

where

$$x_i = \left\lfloor -\beta T_{-\beta}^{i-1}(x) + \frac{\beta}{\beta+1} \right\rfloor,$$

is called the **$(-\beta)$ -expansion** of x , denoted $d_{-\beta}(x) = x_1 x_2 x_3 \dots$.

We denote

$$d_{-\beta}(I_\beta) = d_{-\beta}\left(\frac{-\beta}{\beta+1}\right) = l_1 l_2 l_3 \dots$$

$(-\beta)$ -expansions of real numbers

Consider $-\beta < -1$ and $T_{-\beta} : \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1}\right) \mapsto \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1}\right)$ given by

$$T_{-\beta}(x) := -\beta x - \left\lfloor -\beta x + \frac{\beta}{\beta+1} \right\rfloor.$$

Representation of $x \in I_\beta \equiv [l_\beta, r_\beta) \equiv \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1}\right)$ of the form

$$x = \frac{x_1}{-\beta} + \frac{x_2}{(-\beta)^2} + \frac{x_3}{(-\beta)^3} + \dots,$$

where

$$x_i = \left\lfloor -\beta T_{-\beta}^{i-1}(x) + \frac{\beta}{\beta+1} \right\rfloor,$$

is called the **$(-\beta)$ -expansion** of x , denoted $d_{-\beta}(x) = x_1 x_2 x_3 \dots$.

We denote

$$d_{-\beta}(l_\beta) = d_{-\beta}\left(\frac{-\beta}{\beta+1}\right) = l_1 l_2 l_3 \dots$$

Admissibility condition

An integer sequence $x_1x_2x_3 \cdots$ is said to be **$(-\beta)$ -admissible** if there exists $x \in I_\beta$ such that $d_{-\beta}(x) = x_1x_2x_3 \cdots$.

Theorem (Ito-Sadahiro)

The string $x_1x_2x_3 \cdots$ is $(-\beta)$ -admissible, if and only if for all $i = 1, 2, 3, \dots$,

$$d_{-\beta}(I_\beta) \preceq_{alt} x_i x_{i+1} x_{i+2} \prec_{alt} d_{-\beta}^*(r_\beta),$$

where $d_{-\beta}^*(r_\beta) = \lim_{\varepsilon \rightarrow 0^+} d_{-\beta}(r_\beta - \varepsilon)$.

Admissibility condition

An integer sequence $x_1x_2x_3 \cdots$ is said to be **$(-\beta)$ -admissible** if there exists $x \in I_\beta$ such that $d_{-\beta}(x) = x_1x_2x_3 \cdots$.

Theorem (Ito-Sadahiro)

The string $x_1x_2x_3 \cdots$ is $(-\beta)$ -admissible, if and only if for all $i = 1, 2, 3, \dots$,

$$d_{-\beta}(l_\beta) \preceq_{alt} x_i x_{i+1} x_{i+2} \prec_{alt} d_{-\beta}^*(r_\beta),$$

where $d_{-\beta}^*(r_\beta) = \lim_{\varepsilon \rightarrow 0^+} d_{-\beta}(r_\beta - \varepsilon)$.

Admissibility condition

An integer sequence $x_1x_2x_3 \cdots$ is said to be **$(-\beta)$ -admissible** if there exists $x \in I_\beta$ such that $d_{-\beta}(x) = x_1x_2x_3 \cdots$.

Theorem (Ito-Sadahiro)

The string $x_1x_2x_3 \cdots$ is $(-\beta)$ -admissible, if and only if for all $i = 1, 2, 3, \dots$,

$$d_{-\beta}(l_\beta) \preceq_{alt} x_i x_{i+1} x_{i+2} \prec_{alt} d_{-\beta}^*(r_\beta),$$

where $d_{-\beta}^*(r_\beta) = \lim_{\varepsilon \rightarrow 0^+} d_{-\beta}(r_\beta - \varepsilon)$.

Alternate order.

$$x_1x_2x_3 \cdots \prec_{alt} y_1y_2y_3 \cdots$$

if $(-1)^i(x_i - y_i) > 0$ for the smallest i such that $x_i \neq y_i$.

Admissibility condition

An integer sequence $x_1x_2x_3 \cdots$ is said to be **$(-\beta)$ -admissible** if there exists $x \in I_\beta$ such that $d_{-\beta}(x) = x_1x_2x_3 \cdots$.

Theorem (Ito-Sadahiro)

The string $x_1x_2x_3 \cdots$ is $(-\beta)$ -admissible, if and only if for all $i = 1, 2, 3, \dots$,

$$d_{-\beta}(l_\beta) \preceq_{alt} x_i x_{i+1} x_{i+2} \prec_{alt} d_{-\beta}^*(r_\beta),$$

where $d_{-\beta}^*(r_\beta) = \lim_{\varepsilon \rightarrow 0^+} d_{-\beta}(r_\beta - \varepsilon)$.

Relation between $d_{-\beta}^*(r_\beta)$ and $d_{-\beta}(l_\beta)$.

$$d_{-\beta}^*(r_\beta) = \begin{cases} (0l_1 \cdots l_{2l}(l_{2l+1} - 1))^\omega & \text{for } d_{-\beta}(l_\beta) = (l_1 \cdots l_{2l+1})^\omega \\ 0d_{-\beta}(l_\beta) & \text{otherwise.} \end{cases}$$

Uniqueness problem

Consider $x = \frac{\beta^2}{\beta+1} \notin I_\beta = \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1} \right)$.

- $\frac{x}{-\beta} = \frac{-\beta}{\beta+1}$. Thus

$$d_{-\beta}\left(\frac{x}{-\beta}\right) = l_1 l_2 l_3 \dots$$

- $\frac{x}{(-\beta)^3} = \frac{1}{-\beta(\beta+1)} \in I_\beta$. We compute

$$x_1 = \left\lfloor -\beta \frac{x}{(-\beta)^3} + \frac{\beta}{\beta+1} \right\rfloor = \left\lfloor \frac{1}{\beta+1} + \frac{\beta}{\beta+1} \right\rfloor = 1$$

and

$$T_{-\beta}\left(\frac{x}{(-\beta)^3}\right) = -\beta \frac{x}{(-\beta)^3} - 1 = \frac{1}{\beta+1} - 1 = \frac{-\beta}{\beta+1}.$$

Uniqueness problem

Consider $x = \frac{\beta^2}{\beta+1} \notin I_\beta = \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1} \right)$.

- $\frac{x}{-\beta} = \frac{-\beta}{\beta+1}$. Thus

$$d_{-\beta}\left(\frac{x}{-\beta}\right) = l_1 l_2 l_3 \dots$$

- $\frac{x}{(-\beta)^3} = \frac{1}{-\beta(\beta+1)} \in I_\beta$. We compute

$$x_1 = \left\lfloor -\beta \frac{x}{(-\beta)^3} + \frac{\beta}{\beta+1} \right\rfloor = \left\lfloor \frac{1}{\beta+1} + \frac{\beta}{\beta+1} \right\rfloor = 1$$

and

$$T_{-\beta}\left(\frac{x}{(-\beta)^3}\right) = -\beta \frac{x}{(-\beta)^3} - 1 = \frac{1}{\beta+1} - 1 = \frac{-\beta}{\beta+1}.$$

Uniqueness problem

Consider $x = \frac{\beta^2}{\beta+1} \notin I_\beta = \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1} \right)$.

- $\frac{x}{-\beta} = \frac{-\beta}{\beta+1}$. Thus

$$d_{-\beta}\left(\frac{x}{-\beta}\right) = l_1 l_2 l_3 \dots$$

- $\frac{x}{(-\beta)^3} = \frac{1}{-\beta(\beta+1)} \in I_\beta$. We compute

$$x_1 = \left\lfloor -\beta \frac{x}{(-\beta)^3} + \frac{\beta}{\beta+1} \right\rfloor = \left\lfloor \frac{1}{\beta+1} + \frac{\beta}{\beta+1} \right\rfloor = 1$$

and

$$T_{-\beta}\left(\frac{x}{(-\beta)^3}\right) = -\beta \frac{x}{(-\beta)^3} - 1 = \frac{1}{\beta+1} - 1 = \frac{-\beta}{\beta+1}.$$

Uniqueness problem

Consider $x = \frac{\beta^2}{\beta+1} \notin I_\beta = \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1} \right)$.

- $\frac{x}{-\beta} = \frac{-\beta}{\beta+1}$. Thus

$$d_{-\beta}\left(\frac{x}{-\beta}\right) = l_1 l_2 l_3 \dots$$

- $\frac{x}{(-\beta)^3} = \frac{1}{-\beta(\beta+1)} \in I_\beta$. We compute

$$x_1 = \left\lfloor -\beta \frac{x}{(-\beta)^3} + \frac{\beta}{\beta+1} \right\rfloor = \left\lfloor \frac{1}{\beta+1} + \frac{\beta}{\beta+1} \right\rfloor = 1$$

and

$$T_{-\beta}\left(\frac{x}{(-\beta)^3}\right) = -\beta \frac{x}{(-\beta)^3} - 1 = \frac{1}{\beta+1} - 1 = \frac{-\beta}{\beta+1}.$$

Uniqueness problem

Consider $x = \frac{\beta^2}{\beta+1} \notin I_\beta = \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1}\right)$.

- $\frac{x}{-\beta} = \frac{-\beta}{\beta+1}$. Thus

$$d_{-\beta}\left(\frac{x}{-\beta}\right) = l_1 l_2 l_3 \dots$$

- $\frac{x}{(-\beta)^3} = \frac{1}{-\beta(\beta+1)} \in I_\beta$. Thus

$$d_{-\beta}\left(\frac{x}{(-\beta)^3}\right) = 1 l_1 l_2 l_3 \dots$$

Therefore

$$\frac{1}{-\beta} + \frac{l_1}{(-\beta)^2} + \frac{l_2}{(-\beta)^3} + \dots = \frac{l_1}{(-\beta)^3} + \frac{l_2}{(-\beta)^4} + \frac{l_3}{(-\beta)^5} + \dots$$

Uniqueness problem

Consider $x = \frac{\beta^2}{\beta+1} \notin I_\beta = \left[\frac{-\beta}{\beta+1}, \frac{1}{\beta+1}\right)$.

- $\frac{x}{-\beta} = \frac{-\beta}{\beta+1}$. Thus

$$d_{-\beta}\left(\frac{x}{-\beta}\right) = l_1 l_2 l_3 \dots$$

- $\frac{x}{(-\beta)^3} = \frac{1}{-\beta(\beta+1)} \in I_\beta$. Thus

$$d_{-\beta}\left(\frac{x}{(-\beta)^3}\right) = 1 l_1 l_2 l_3 \dots$$

Therefore

$$\frac{1}{-\beta} + \frac{l_1}{(-\beta)^2} + \frac{l_2}{(-\beta)^3} + \dots = \frac{l_1}{(-\beta)^3} + \frac{l_2}{(-\beta)^4} + \frac{l_3}{(-\beta)^5} + \dots$$

Lemma

Let $x_1x_2x_3\cdots$ be a $(-\beta)$ -admissible sequence with $x_1 \neq 0$. For fixed $k \in \mathbb{Z}$, denote

$$z = \sum_{i=1}^{\infty} x_i(-\beta)^{k-i}.$$

Then

$$z \in \begin{cases} \left[\frac{\beta^{k-1}}{\beta+1}, \frac{\beta^{k+1}}{\beta+1} \right] & \text{for } k \text{ odd,} \\ \left[-\frac{\beta^{k+1}}{\beta+1}, -\frac{\beta^{k-1}}{\beta+1} \right] & \text{for } k \text{ even.} \end{cases}$$

Remark. Numbers with two different $(-\beta)$ -admissible expansions

$$z = \frac{(-\beta)^k}{\beta+1} \quad \text{for } k \in \mathbb{Z}.$$

Lemma

Let $x_1x_2x_3\cdots$ be a $(-\beta)$ -admissible sequence with $x_1 \neq 0$. For fixed $k \in \mathbb{Z}$, denote

$$z = \sum_{i=1}^{\infty} x_i (-\beta)^{k-i}.$$

Then

$$z \in \begin{cases} \left[\frac{\beta^{k-1}}{\beta+1}, \frac{\beta^{k+1}}{\beta+1} \right] & \text{for } k \text{ odd,} \\ \left[-\frac{\beta^{k+1}}{\beta+1}, -\frac{\beta^{k-1}}{\beta+1} \right] & \text{for } k \text{ even.} \end{cases}$$

Remark. Numbers with two different $(-\beta)$ -admissible expansions

$$z = \frac{(-\beta)^k}{\beta+1} \quad \text{for } k \in \mathbb{Z}.$$

Greedy algorithm

Greedy algorithm for computing the $(-\beta)$ -expansion of $x \in \mathbb{R}$

Require: $x \in \mathbb{R}$

while $x \neq 0$ **do**

if $x > 0$ **then**

 find the maximal even $k \in \mathbb{Z}$ such that $x \geq \frac{\beta^k}{\beta+1}$.

end if

if $x < 0$ **then**

 find the maximal odd $k \in \mathbb{Z}$ such that $x \leq \frac{-\beta^k}{\beta+1}$.

end if

$$x_k \leftarrow \lfloor \frac{x}{(-\beta)^k} + \frac{\beta}{\beta+1} \rfloor$$

$$x \leftarrow x - x_k (-\beta)^k$$

end while

- 1 Rényi expansions
- 2 Ito-Sadahiro expansions
- 3 $(-\beta)$ -integers**
- 4 Examples
- 5 Open questions

$(-\beta)$ -integers

Set of $(-\beta)$ -integers

$$\mathbb{Z}_{-\beta} = \{x_k(-\beta)^k + \dots + x_1(-\beta) + x_0 \mid x_k \dots x_1 x_0 0^\omega \text{ is } (-\beta)\text{-admissible}\}.$$

Remark.

- $0 \in I_\beta$ and $T_{-\beta}(0) = 0 \Rightarrow d_{-\beta}(0) = 0^\omega$ and thus $0 \in \mathbb{Z}_{-\beta}$
- β minimal Pisot number, then $d_{-\beta}(I_\beta) = 1001^\omega$
 $x_k \dots x_1 x_0 0^\omega \neq 0^\omega$ is $(-\beta)$ -admissible \Rightarrow so is 10^ω
But $1001^\omega \not\leq_{\text{alt}} 10^\omega \Rightarrow \mathbb{Z}_{-\beta} = \{0\}$.

Lemma

$\mathbb{Z}_{-\beta} = \{0\}$ iff $10^{2k}1$ is a prefix of $d_{-\beta}(I_\beta)$ for some $k \geq 0$.

$(-\beta)$ -integers

Set of $(-\beta)$ -integers

$$\mathbb{Z}_{-\beta} = \{x_k(-\beta)^k + \dots + x_1(-\beta) + x_0 \mid x_k \dots x_1 x_0 0^\omega \text{ is } (-\beta)\text{-admissible}\}.$$

Remark.

- $0 \in I_\beta$ and $T_{-\beta}(0) = 0 \Rightarrow d_{-\beta}(0) = 0^\omega$ and thus $0 \in \mathbb{Z}_{-\beta}$
- β minimal Pisot number, then $d_{-\beta}(I_\beta) = 1001^\omega$
 $x_k \dots x_1 x_0 0^\omega \neq 0^\omega$ is $(-\beta)$ -admissible \Rightarrow so is 10^ω
But $1001^\omega \not\leq_{\text{alt}} 10^\omega \Rightarrow \mathbb{Z}_{-\beta} = \{0\}$.

Lemma

$\mathbb{Z}_{-\beta} = \{0\}$ iff $10^{2k}1$ is a prefix of $d_{-\beta}(I_\beta)$ for some $k \geq 0$.

$(-\beta)$ -integers

Set of $(-\beta)$ -integers

$$\mathbb{Z}_{-\beta} = \{x_k(-\beta)^k + \dots + x_1(-\beta) + x_0 \mid x_k \dots x_1 x_0 0^\omega \text{ is } (-\beta)\text{-admissible}\}.$$

Remark.

- $0 \in I_\beta$ and $T_{-\beta}(0) = 0 \Rightarrow d_{-\beta}(0) = 0^\omega$ and thus $0 \in \mathbb{Z}_{-\beta}$
- β minimal Pisot number, then $d_{-\beta}(l_\beta) = 1001^\omega$

$x_k \dots x_1 x_0 0^\omega \neq 0^\omega$ is $(-\beta)$ -admissible \Rightarrow so is 10^ω

But $1001^\omega \not\prec_{\text{alt}} 10^\omega \Rightarrow \mathbb{Z}_{-\beta} = \{0\}$.

Lemma

$\mathbb{Z}_{-\beta} = \{0\}$ iff $10^{2k}1$ is a prefix of $d_{-\beta}(l_\beta)$ for some $k \geq 0$.

$(-\beta)$ -integers

Set of $(-\beta)$ -integers

$$\mathbb{Z}_{-\beta} = \{x_k(-\beta)^k + \dots + x_1(-\beta) + x_0 \mid x_k \dots x_1 x_0 0^\omega \text{ is } (-\beta)\text{-admissible}\}.$$

Remark.

- $0 \in I_\beta$ and $T_{-\beta}(0) = 0 \Rightarrow d_{-\beta}(0) = 0^\omega$ and thus $0 \in \mathbb{Z}_{-\beta}$
- β minimal Pisot number, then $d_{-\beta}(I_\beta) = 1001^\omega$
 $x_k \dots x_1 x_0 0^\omega \neq 0^\omega$ is $(-\beta)$ -admissible \Rightarrow so is 10^ω
But $1001^\omega \not\leq_{\text{alt}} 10^\omega \Rightarrow \mathbb{Z}_{-\beta} = \{0\}$.

Lemma

$\mathbb{Z}_{-\beta} = \{0\}$ iff $10^{2k}1$ is a prefix of $d_{-\beta}(I_\beta)$ for some $k \geq 0$.

$(-\beta)$ -integers

Set of $(-\beta)$ -integers

$$\mathbb{Z}_{-\beta} = \{x_k(-\beta)^k + \dots + x_1(-\beta) + x_0 \mid x_k \dots x_1 x_0 0^\omega \text{ is } (-\beta)\text{-admissible}\}.$$

Remark.

- $0 \in I_\beta$ and $T_{-\beta}(0) = 0 \Rightarrow d_{-\beta}(0) = 0^\omega$ and thus $0 \in \mathbb{Z}_{-\beta}$
- β minimal Pisot number, then $d_{-\beta}(I_\beta) = 1001^\omega$
 $x_k \dots x_1 x_0 0^\omega \neq 0^\omega$ is $(-\beta)$ -admissible \Rightarrow so is 10^ω
But $1001^\omega \not\stackrel{L}{\sim}_{\text{alt}} 10^\omega \Rightarrow \mathbb{Z}_{-\beta} = \{0\}$.

Lemma

$\mathbb{Z}_{-\beta} = \{0\}$ iff $10^{2k}1$ is a prefix of $d_{-\beta}(I_\beta)$ for some $k \geq 0$.

$(-\beta)$ -integers

Set of $(-\beta)$ -integers

$$\mathbb{Z}_{-\beta} = \{x_k(-\beta)^k + \dots + x_1(-\beta) + x_0 \mid x_k \dots x_1 x_0 0^\omega \text{ is } (-\beta)\text{-admissible}\}.$$

Remark.

- $0 \in I_\beta$ and $T_{-\beta}(0) = 0 \Rightarrow d_{-\beta}(0) = 0^\omega$ and thus $0 \in \mathbb{Z}_{-\beta}$
- β minimal Pisot number, then $d_{-\beta}(l_\beta) = 1001^\omega$
 $x_k \dots x_1 x_0 0^\omega \neq 0^\omega$ is $(-\beta)$ -admissible \Rightarrow so is 10^ω
But $1001^\omega \not\stackrel{L}{\sim}_{\text{alt}} 10^\omega \Rightarrow \mathbb{Z}_{-\beta} = \{0\}$.

Lemma

$\mathbb{Z}_{-\beta} = \{0\}$ iff $10^{2k}1$ is a prefix of $d_{-\beta}(l_\beta)$ for some $k \geq 0$.

Gaps in $\mathbb{Z}_{-\beta}$ — general proposition

$\mathcal{S}(k) = \{x_{k-1}x_{k-2} \cdots x_0 0^\omega \mid x_{k-1}x_{k-2} \cdots x_0 0^\omega \text{ is } (-\beta)\text{-admissible}\}.$

$\text{Max}(k) =$ maximal in $\mathcal{S}(k)$ with respect to the alternate order,

$\text{Min}(k) =$ minimal in $\mathcal{S}(k)$ with respect to the alternate order.

Proposition

Let Δ be the distance of two consecutive $(-\beta)$ -integers.

Then there exists a $k \in \{0, 1, 2, \dots\}$ such that

$$\Delta = \beta^{2k} + \gamma(\text{Min}(2k)) - \gamma(\text{Max}(2k))$$

or

$$\Delta = \beta^{2k+1} + \gamma(\text{Max}(2k+1)) - \gamma(\text{Min}(2k+1)).$$

Gaps in $\mathbb{Z}_{-\beta}$ — general proposition

$\mathcal{S}(k) = \{x_{k-1}x_{k-2} \cdots x_0 0^\omega \mid x_{k-1}x_{k-2} \cdots x_0 0^\omega \text{ is } (-\beta)\text{-admissible}\}.$

$\text{Max}(k) =$ maximal in $\mathcal{S}(k)$ with respect to the alternate order,

$\text{Min}(k) =$ minimal in $\mathcal{S}(k)$ with respect to the alternate order.

Proposition

Let Δ be the distance of two consecutive $(-\beta)$ -integers.

Then there exists a $k \in \{0, 1, 2, \dots\}$ such that

$$\Delta = \beta^{2k} + \gamma(\text{Min}(2k)) - \gamma(\text{Max}(2k))$$

or

$$\Delta = \beta^{2k+1} + \gamma(\text{Max}(2k+1)) - \gamma(\text{Min}(2k+1)).$$

Gaps in $\mathbb{Z}_{-\beta}$ — general proposition

$\mathcal{S}(k) = \{x_{k-1}x_{k-2} \cdots x_0 0^\omega \mid x_{k-1}x_{k-2} \cdots x_0 0^\omega \text{ is } (-\beta)\text{-admissible}\}.$

$\text{Max}(k) =$ maximal in $\mathcal{S}(k)$ with respect to the alternate order,

$\text{Min}(k) =$ minimal in $\mathcal{S}(k)$ with respect to the alternate order.

Proposition

Let Δ be the distance of two consecutive $(-\beta)$ -integers.

Then there exists a $k \in \{0, 1, 2, \dots\}$ such that

$$\Delta = \beta^{2k} + \gamma(\text{Min}(2k)) - \gamma(\text{Max}(2k))$$

or

$$\Delta = \beta^{2k+1} + \gamma(\text{Max}(2k+1)) - \gamma(\text{Min}(2k+1)).$$

Theorem

Let $d_{-\beta}(l_\beta) = l_1 l_2 l_3 \cdots$ where $0 < l_i < l_1$ for all $i = 2, 3, 4, \dots$.
Then the distances between adjacent $(-\beta)$ -integers take values

$$\Delta_0 = 1$$

$$\Delta_k = \left| (-1)^k + \sum_{i=1}^{\infty} \frac{l_{k-1+i} - l_{k+i}}{(-\beta)^i} \right|, \quad k = 1, 2, 3, \dots$$

Moreover, all the distances are less than 2.

Gaps in $\mathbb{Z}_{-\beta}$ — second case

Theorem

Let $d_{-\beta}(l_\beta) = l_1 l_2 \cdots l_m 0^\omega$, where $l_m \neq 0$.

If $0 < l_i < l_1$ for all $i = 2, 3, 4, \dots, m$, then

$$\left. \begin{array}{l} m \text{ even} \\ \\ \\ \end{array} \right\} \begin{cases} \Delta_0 = 1, \\ \Delta_k = \left| (-1)^k + \sum_{i=1}^{\infty} \frac{l_{k-1+i} - l_{k+i}}{(-\beta)^i} \right|, & k = 1, \dots, m \end{cases}$$
$$\left. \begin{array}{l} m \text{ odd} \\ \\ \\ \end{array} \right\} \begin{cases} \Delta_0 = 1, \\ \Delta_k = \left| (-1)^k + \sum_{i=1}^{\infty} \frac{l_{k-1+i} - l_{k+i}}{(-\beta)^i} \right|, & k = 1, \dots, m-1, \\ \Delta_m = \frac{l_m}{\beta} \end{cases}$$

Moreover, all the distances are less than 2.

- 1 Rényi expansions
- 2 Ito-Sadahiro expansions
- 3 $(-\beta)$ -integers
- 4 Examples**
- 5 Open questions

Tribonacci number β , root of $x^3 - x^2 - x - 1$

β -expansions

- $d_\beta(1) = 1110^\omega$
- $\Delta_0 = 1$, $\Delta_1 = \beta - 1$ and $\Delta_2 = \frac{1}{\beta}$.

$(-\beta)$ -expansions

- $d_{-\beta}(l_\beta) = 101^\omega$,
- $d_{-\beta}^*(r_\beta) = 0101^\omega$,

$$\text{Min}(2k) = 10(11)^{k-1}, \quad \text{Min}(2k+1) = 10(11)^{k-1}0,$$

$$\text{Max}(2k) = 010(11)^{k-2}0, \quad \text{Max}(2k+1) = 010(11)^{k-1},$$

$$\text{Max}(2) = 01.$$

Distances are

$$\Delta_0 = 1, \quad \Delta_1 = \beta - 1, \quad \text{and} \quad \Delta_2 = \frac{1}{\beta}.$$

Tribonacci number β , root of $x^3 - x^2 - x - 1$

β -expansions

- $d_\beta(1) = 1110^\omega$
- $\Delta_0 = 1$, $\Delta_1 = \beta - 1$ and $\Delta_2 = \frac{1}{\beta}$.

$(-\beta)$ -expansions

- $d_{-\beta}(l_\beta) = 101^\omega$,
- $d_{-\beta}^*(r_\beta) = 0101^\omega$,

$$\text{Min}(2k) = 10(11)^{k-1}, \quad \text{Min}(2k+1) = 10(11)^{k-1}0,$$

$$\text{Max}(2k) = 010(11)^{k-2}0, \quad \text{Max}(2k+1) = 010(11)^{k-1},$$

$$\text{Max}(2) = 01.$$

Distances are

$$\Delta_0 = 1, \quad \Delta_1 = \beta - 1, \quad \text{and} \quad \Delta_2 = \frac{1}{\beta}.$$

Tribonacci number β , root of $x^3 - x^2 - x - 1$

β -expansions

- $d_\beta(1) = 1110^\omega$
- $\Delta_0 = 1$, $\Delta_1 = \beta - 1$ and $\Delta_2 = \frac{1}{\beta}$.

$(-\beta)$ -expansions

- $d_{-\beta}(l_\beta) = 101^\omega$,
- $d_{-\beta}^*(r_\beta) = 0101^\omega$,

$$\text{Min}(2k) = 10(11)^{k-1}, \quad \text{Min}(2k+1) = 10(11)^{k-1}0,$$

$$\text{Max}(2k) = 010(11)^{k-2}0, \quad \text{Max}(2k+1) = 010(11)^{k-1},$$

$$\text{Max}(2) = 01.$$

Distances are

$$\Delta_0 = 1, \quad \Delta_1 = \beta - 1, \quad \text{and} \quad \Delta_2 = \frac{1}{\beta}.$$

Tribonacci number β , root of $x^3 - x^2 - x - 1$

β -expansions

- $d_\beta(1) = 1110^\omega$
- $\Delta_0 = 1$, $\Delta_1 = \beta - 1$ and $\Delta_2 = \frac{1}{\beta}$.

$(-\beta)$ -expansions

- $d_{-\beta}(l_\beta) = 101^\omega$,
- $d_{-\beta}^*(r_\beta) = 0101^\omega$,

$$\text{Min}(2k) = 10(11)^{k-1}, \quad \text{Min}(2k+1) = 10(11)^{k-1}0,$$

$$\text{Max}(2k) = 010(11)^{k-2}0, \quad \text{Max}(2k+1) = 010(11)^{k-1},$$

$$\text{Max}(2) = 01.$$

Distances are

$$\Delta_0 = 1, \quad \Delta_1 = \beta - 1, \quad \text{and} \quad \Delta_2 = \frac{1}{\beta}.$$

Tribonacci number β , root of $x^3 - x^2 - x - 1$

β -expansions

- $d_\beta(1) = 1110^\omega$
- $\Delta_0 = 1$, $\Delta_1 = \beta - 1$ and $\Delta_2 = \frac{1}{\beta}$.

$(-\beta)$ -expansions

- $d_{-\beta}(l_\beta) = 101^\omega$,
- $d_{-\beta}^*(r_\beta) = 0101^\omega$,

$$\text{Min}(2k) = 10(11)^{k-1}, \quad \text{Min}(2k+1) = 10(11)^{k-1}0,$$

$$\text{Max}(2k) = 010(11)^{k-2}0, \quad \text{Max}(2k+1) = 010(11)^{k-1},$$

$$\text{Max}(2) = 01.$$

Distances are

$$\Delta_0 = 1, \quad \Delta_1 = \beta - 1, \quad \text{and} \quad \Delta_2 = \frac{1}{\beta}.$$

β root of $x^3 - 2x^2 - x + 1$

β -expansions

- $d_\beta(1) = 2(01)^\omega$
- $\Delta_0 = 1$, $\Delta_1 = \beta - 2$, and $\Delta_3 = \beta^2 - 2\beta$.

$(-\beta)$ -integers

- $d_{-\beta}(1_\beta) = 210^\omega$
- By Theorem

$$\tilde{\Delta}_0 = 1,$$

$$\tilde{\Delta}_1 = \beta - 1,$$

$$\tilde{\Delta}_2 = 1 - \frac{1}{\beta}.$$

Distances in \mathbb{Z}_β and $\mathbb{Z}_{-\beta}$ are different.

β root of $x^3 - 2x^2 - x + 1$

β -expansions

- $d_\beta(1) = 2(01)^\omega$
- $\Delta_0 = 1$, $\Delta_1 = \beta - 2$, and $\Delta_3 = \beta^2 - 2\beta$.

$(-\beta)$ -integers

- $d_{-\beta}(1_\beta) = 210^\omega$
- By Theorem

$$\tilde{\Delta}_0 = 1,$$

$$\tilde{\Delta}_1 = \beta - 1,$$

$$\tilde{\Delta}_2 = 1 - \frac{1}{\beta}.$$

Distances in \mathbb{Z}_β and $\mathbb{Z}_{-\beta}$ are different.

β root of $x^3 - 2x^2 - x + 1$

β -expansions

- $d_\beta(1) = 2(01)^\omega$
- $\Delta_0 = 1$, $\Delta_1 = \beta - 2$, and $\Delta_3 = \beta^2 - 2\beta$.

$(-\beta)$ -integers

- $d_{-\beta}(1_\beta) = 210^\omega$
- By Theorem

$$\tilde{\Delta}_0 = 1,$$

$$\tilde{\Delta}_1 = \beta - 1 > 1,$$

$$\tilde{\Delta}_2 = 1 - \frac{1}{\beta}.$$

Distances in \mathbb{Z}_β and $\mathbb{Z}_{-\beta}$ are **different**.


- 1 Rényi expansions
- 2 Ito-Sadahiro expansions
- 3 $(-\beta)$ -integers
- 4 Examples
- 5 Open questions**

Open questions


- Gaps in $\mathbb{Z}_{-\beta}$ in general
- What does the projection of $\mathbb{Z}_{-\beta}$ into the contracting plane give?

Open questions

- Gaps in $\mathbb{Z}_{-\beta}$ in general
- What does the projection of $\mathbb{Z}_{-\beta}$ into the contracting plane give?




Projection of \mathbb{Z}_{β} ,
 β Tribonacci number.



Projection of $\mathbb{Z}_{-\beta}$,
 β Tribonacci number.

Open questions

- Gaps in $\mathbb{Z}_{-\beta}$ in general
- What does the projection of $\mathbb{Z}_{-\beta}$ into the contracting plane give?



Projection of \mathbb{Z}_β ,
 β root of $x^3 = 2x^2 + x - 1$.



Projection of $\mathbb{Z}_{-\beta}$,
 β root of $x^3 = 2x^2 + x - 1$.

Open questions

- Gaps in $\mathbb{Z}_{-\beta}$ in general
- What does the projection of $\mathbb{Z}_{-\beta}$ into the contracting plane give?

If we code gaps in $\mathbb{Z}_{-\beta}$ by an infinite word $\mathbf{u}_{-\beta}$

- Is $\mathbf{u}_{-\beta}$ fixed point of some substitution?
- Is there any relation with the canonical substitution φ_β associated to β -numeration system?

S. Ito and T. Sadahiro,
 $(-\beta)$ -expansions of real numbers,
Integers 9 (2009), 239–259.

W.P. Thurston,
Groups, tilings, and finite state automata,
AMS Colloquium Lecture Notes, American Mathematical Society,
Boulder, 1989.

Petr Ambrož, Daniel Dombek, Zuzana Masáková, Edita Pelantová,
Numbers with integer expansion in the system with negative base,
submitted (2010).