

Circular Coloring of Signed Graphs

Zhouningxin Wang

Université de Paris

wangzhou4@irif.fr

(A joint work with Reza Naserasr, Xuding Zhu)

March 20, 2021

- 1 Introduction
 - Circular coloring of graphs
 - Homomorphism of signed graphs
- 2 Circular coloring of signed graphs
 - Circular chromatic number
 - Signed indicators
 - Tight cycle argument
- 3 Results on some classes of signed graphs
 - Signed bipartite planar graphs
 - Signed d-degenerate graphs
 - Signed planar graphs
- 4 Discussion

Circular coloring of graphs

Given a real number r , a **circular r -coloring** of a graph G is a mapping $f : V(G) \rightarrow C^r$ such that for any edge $uv \in E(G)$,

$$d_{(\text{mod } r)}(f(u), f(v)) \geq 1.$$

The **circular chromatic number** of G is defined as

$$\chi_c(G) = \inf\{r : G \text{ admits a circular } r\text{-coloring}\}.$$

Circular coloring of graphs

- A 3-chromatic graph is not 2-colorable, but if its circular chromatic number is near 2, then it is somehow “just barely” not 2-colorable.
- By Grotzsch’s theorem, every triangle-free planar graph is 3-colorable. In generalizing this to circular chromatic number, we may ask what threshold on girth is needed to force the circular chromatic number to be at most $2 + \frac{1}{t}$.

Jaeger-Zhang conjecture [C.-Q. Zhang 2002]

Every planar graph of odd-girth $4k + 1$ admits a circular $(2 + \frac{1}{k})$ -coloring.

Homomorphism of signed graphs

- A **signed graph** is a graph $G = (V, E)$ together with an assignment $\{+, -\}$ on its edges, denoted by (G, σ) .
- A **switching** at vertex v is to switch the signs of all the edges incident to this vertex.
- The **sign** of a closed walk is the product of signs of all the edges of this walk.
- A **homomorphism** of signed graph (G, σ) to a signed graph (H, π) is a mapping φ from $V(G)$ and $E(G)$ correspondingly to $V(H)$ and $E(H)$ such that the adjacency, the incidence and the signs of the closed walks are preserved.
- If there exists one, we write $(G, \sigma) \rightarrow (H, \pi)$.

Homomorphism of signed graphs

- An **edge-sign preserving homomorphism** of a signed graph (G, σ) to (H, π) is a mapping $f : V(G) \rightarrow V(H)$ such that for every positive (respectively, negative) edge uv of (G, σ) , $f(u)f(v)$ is a positive (respectively, negative) edge of (H, π) .
- If there exists one, we write $(G, \sigma) \xrightarrow{s.p.} (H, \pi)$.

Proposition

Given two signed graphs (G, σ) and (H, π) ,

$$(G, \sigma) \rightarrow (H, \pi) \Leftrightarrow \exists \sigma' \equiv \sigma, (G, \sigma') \xrightarrow{s.p.} (H, \pi).$$

Double Switching Graphs

Given a signed graph (G, σ) on the vertex set $V = \{x_1, \dots, x_n\}$, the **Double Switching Graph** of (G, σ) , denoted $\text{DSG}(G, \sigma)$, is a signed graph built as follows:

- We have two disjoint copies of V , $V^+ = \{x_1^+, x_2^+, \dots, x_n^+\}$ and $V^- = \{x_1^-, x_2^-, \dots, x_n^-\}$ in $\text{DSG}(G, \sigma)$.
- Each set of vertices V^+, V^- then induces a copy of (G, σ) .
- Furthermore, a vertex x_i^- connects to vertices in V^+ as it is obtained from a switching on x_i .

Double Switching Graphs




Figure: Signed graphs (C_4, e) and $DSG(C_4, e)$

Theorem [R.C. Brewster and T. Graves 2009]

Given signed graphs (G, σ) and (H, π) ,

$$(G, \sigma) \rightarrow (H, \pi) \Leftrightarrow (G, \sigma) \xrightarrow{s.p.} DSG(H, \pi).$$

- 1 Introduction
 - Circular coloring of graphs
 - Homomorphism of signed graphs
- 2 Circular coloring of signed graphs
 - Circular chromatic number
 - Signed indicators
 - Tight cycle argument
- 3 Results on some classes of signed graphs
 - Signed bipartite planar graphs
 - Signed d-degenerate graphs
 - Signed planar graphs
- 4 Discussion

Circular coloring of signed graphs

Given a signed graph (G, σ) with no positive loop and a real number r , a **circular r -coloring** of (G, σ) is a mapping $f : V(G) \rightarrow C^r$ such that for each positive edge uv of (G, σ) ,

$$d_{(\text{mod } r)}(f(u), f(v)) \geq 1,$$

and for each negative edge uv of (G, σ) ,

$$d_{(\text{mod } r)}(f(u), \overline{f(v)}) \geq 1.$$

The **circular chromatic number** of (G, σ) is defined as

$$\chi_c(G, \sigma) = \inf\{r \geq 1 : (G, \sigma) \text{ admits a circular } r\text{-coloring}\}.$$

Refinement of 0-free $2k$ -coloring of signed graphs

Definition [T. Zaslavsky 1982]

Given a signed graph (G, σ) and a positive integer k , a **0-free $2k$ -coloring** of (G, σ) is a mapping $f : V(G) \rightarrow \{\pm 1, \pm 2, \dots, \pm k\}$ such that for any edge uv of (G, σ) , $f(u) \neq \sigma(uv)f(v)$.

Proposition

Assume (G, σ) is a signed graph and k is a positive integer. Then (G, σ) is 0-free $2k$ -colorable if and only if (G, σ) is circular $2k$ -colorable.

Equivalent definition

Note that for $s, t \in [0, r)$, $d_{(\text{mod } r)}(s, t) = \min\{|s - t|, r - |s - t|\}$.

- A **circular r -coloring** of a signed graph (G, σ) is a mapping $f : V(G) \rightarrow [0, r)$ such that for each positive edge uv ,

$$1 \leq |f(u) - f(v)| \leq r - 1$$

and for each negative edge uv ,

$$\text{either } |f(u) - f(v)| \leq \frac{r}{2} - 1 \text{ or } |f(u) - f(v)| \geq \frac{r}{2} + 1.$$

Equivalent definition: (p, q) -coloring of signed graphs

For $i, j, x \in \{0, 1, \dots, p-1\}$, we define

$d_{(\text{mod } p)}(i, j) = \min\{|i - j|, p - |i - j|\}$ and $\bar{x} = x + \frac{p}{2} \pmod{p}$.

- Assume p is an even integer and $q \leq \frac{p}{2}$ is a positive integer.

A (p, q) -coloring of a signed graph (G, σ) is a mapping

$f : V(G) \rightarrow \{0, 1, \dots, p-1\}$ such that for any positive edge

uv ,

$$d_{(\text{mod } p)}(f(u), f(v)) \geq q,$$

and for any negative edge uv ,

$$d_{(\text{mod } p)}(f(u), \overline{f(v)}) \geq q.$$

The **circular chromatic number** of (G, σ) is

$$\chi_c(G, \sigma) = \inf\left\{\frac{p}{q} : (G, \sigma) \text{ has a } (p, q)\text{-coloring}\right\}.$$

Signed circular clique

Circular chromatic number of signed graphs are also defined through graph homomorphism.

For integers $p \geq 2q > 0$ such that p is even, the **signed circular clique** $K_{p,q}^s$ has vertex set $[p] = \{0, 1, \dots, p-1\}$, in which

- ij is a positive edge if $q \leq |i-j| \leq p-q$;
- ij is a negative edge if $|i-j| \leq \frac{p}{2} - q$ or $|i-j| \geq \frac{p}{2} + q$.

Signed circular clique

Lemma

Given a signed graph (G, σ) and a positive even integer p , a positive integer q with $p \geq 2q$, (G, σ) has a (p, q) -coloring if and only if $(G, \sigma) \xrightarrow{s.p.} K_{p;q}^s$.

Hence the circular chromatic number of (G, σ) is

$$\chi_c(G, \sigma) = \inf \left\{ \frac{p}{q} : p \text{ is even and } (G, \sigma) \xrightarrow{s.p.} K_{p;q}^s \right\}.$$

Lemma

If $(G, \sigma) \xrightarrow{s.p.} (H, \pi)$, then $\chi_c(G, \sigma) \leq \chi_c(H, \pi)$.



Lemma

Given even positive integers p, p' , if $\frac{p}{q} \leq \frac{p'}{q'}$, then $K_{p;q}^s \xrightarrow{s.p.} K_{p';q'}^s$.

Signed circular clique

Let $\hat{K}_{p;q}^s$ be the signed subgraph of $K_{p;q}^s$ induced by vertices $\{0, 1, \dots, \frac{p}{2} - 1\}$. Notice that $K_{p;q}^s = \text{DSG}(\hat{K}_{p;q}^s)$.
The circular chromatic number of (G, σ) is also

$$\chi_c(G, \sigma) = \inf \left\{ \frac{p}{q} : p \text{ is even and } (G, \sigma) \rightarrow \hat{K}_{p;q}^s \right\}.$$

Figure: $K_{8;3}^s$ Figure: $\hat{K}_{8;3}^s$

Circular chromatic number of cycles

For a non-zero integer ℓ , we denote by C_ℓ the cycle of length $|\ell|$ whose sign agrees with the sign of ℓ .

Proposition

$$\chi_c(C_{2k}) = \chi_c(C_{-(2k+1)}) = 2; \quad \chi_c(C_{2k+1}) = \frac{2k+1}{k};$$

$$\chi_c(C_{-2k}) = \frac{4k}{2k-1}.$$

Observe that the signed graph $\hat{K}_{4k;2k-1}^s$ is obtained from C_{-2k} by adding a negative loop at each vertex.

C_{2k+1} -coloring and C_{-2k} -coloring

Proposition

- Given a graph G , $G \rightarrow C_{2k+1}$ if and only if $\chi_c(G) \leq \frac{2k+1}{k}$;
- Given a signed bipartite graph (G, σ) ,

$$(G, \sigma) \rightarrow C_{-2k} \text{ if and only if } \chi_c(G, \sigma) \leq \frac{4k}{2k-1}.$$

Signed indicator

Let G be a graph and let Ω be a signed graph.

- A **signed indicator** \mathcal{I} is a triple $\mathcal{I} = (\Gamma, u, v)$ such that Γ is a signed graph and u, v are two distinct vertices of Γ .
- **Replacing e of G with a copy of \mathcal{I}** is the following operation: Take the disjoint union of Ω and \mathcal{I} , delete the edge e from Ω , identify x with u and identify y with v .
- Given a signed indicator \mathcal{I} , we denote by $G(\mathcal{I})$ the signed graph obtained from G by replacing each edge with a copy of \mathcal{I} .
- Given two signed indicators \mathcal{I}_+ and \mathcal{I}_- , we denote by $\Omega(\mathcal{I}_+, \mathcal{I}_-)$ the signed graph obtained from Ω by replacing each positive edge with a copy of \mathcal{I}_+ and replacing each negative edge with a copy of \mathcal{I}_- .

Signed indicator

Assume $\mathcal{I} = (\Gamma, u, v)$ is a signed indicator and $r \geq 2$ is a real number.

- For $a, b \in [0, r)$, we say the color pair (a, b) is **feasible for \mathcal{I}** (with respect to r) if there is a circular r -coloring ϕ of Γ such that $\phi(u) = a$ and $\phi(v) = b$.
- Define

$$Z(\mathcal{I}, r) = \{b \in [0, \frac{r}{2}] : (0, b) \text{ is feasible for } \mathcal{I} \text{ with respect to } r\}.$$

Lemma

Assume that $\mathcal{I} = (\Gamma, u, v)$ is a signed indicator, $r \geq 2$ is a real number and $Z(\mathcal{I}, r) = [t, \frac{r}{2} - t]$ for some $0 < t < \frac{r}{4}$. Then for any graph G ,

$$\chi_c(G) = \frac{\chi_c(G(\mathcal{I}))}{2t}.$$

Examples

- If Γ is a positive 2-path connecting u and v , and $\mathcal{I} = (\Gamma, u, v)$, then for any ϵ , $0 < \epsilon < 1$, and $r = 4 - 2\epsilon$,

$$Z(\mathcal{I}, r) = [0, 2 - 2\epsilon] = [0, \frac{r}{2} - \epsilon].$$

- If Γ' is a negative 2-path connecting u and v , and $\mathcal{I}' = (\Gamma', u, v)$, then for any ϵ , $0 < \epsilon < 1$, and $r = 4 - 2\epsilon$,

$$Z(\mathcal{I}', r) = [\epsilon, \frac{r}{2}].$$

- If Γ'' consists of a negative 2-path and a positive 2-path connecting u and v , and $\mathcal{I}'' = (\Gamma'', u, v)$, then for any ϵ , $0 < \epsilon < 1$, and $r = 4 - 2\epsilon$,

$$Z(\mathcal{I}'', r) = [\epsilon, \frac{r}{2} - \epsilon].$$

Indicator construction $S(G)$

Given a graph G , a signed graph $S(G)$ is built as follows.




Figure: $S(K_3)$




Figure: $S(C_5)$

Corollary

For any graph G ,

$$\chi_c(S(G)) = 4 - \frac{4}{\chi_c(G) + 1}.$$

Signed indicator

Lemma

Assume that \mathcal{I}_+ and \mathcal{I}_- are indicators, $r \geq 2$ is a real number and

$$Z(\mathcal{I}_+, r) = [t, \frac{r}{2}], Z(\mathcal{I}_-, r) = [0, \frac{r}{2} - t]$$

for some $0 < t < \frac{r}{2}$. Then for any signed graph Ω ,

$$\chi_c(\Omega) = \frac{\chi_c(\Omega(\mathcal{I}_+, \mathcal{I}_-))}{t}.$$

Tight cycle argument

Assume (G, σ) is a signed graph and $\phi : V(G) \rightarrow [0, r)$ is a circular r -coloring of (G, σ) . The **partial orientation** $D = D_\phi(G, \sigma)$ of G with respect to a circular r -coloring ϕ is defined as follows: (u, v) is an arc of D if and only if one of the following holds:

- uv is a positive edge and $(\phi(v) - \phi(u))(\bmod r) = 1$.
- uv is a negative edge and $(\overline{\phi(v)} - \phi(u))(\bmod r) = 1$.

Arcs in $D_\phi(G, \sigma)$ are called **tight arcs** of (G, σ) with respect to ϕ . A directed cycle in $D_\phi(G, \sigma)$ is called a **tight cycle** with respect to ϕ .

Tight cycle argument

Lemma

Let (G, σ) be a signed graph and let ϕ be a circular r -coloring of (G, σ) . If $D_\phi(G, \sigma)$ is acyclic, then there exists an $r_0 \not\leq r$ such that (G, σ) admits an r_0 -circular coloring.

Notice that assume $D_\phi(G, \sigma)$ is acyclic and among all such ϕ , $D_\phi(G, \sigma)$ has minimum number of arcs, then $D_\phi(G, \sigma)$ has no arc.

Lemma

Given a signed graph (G, σ) , $\chi_c(G, \sigma) = r$ if and only if (G, σ) is circular r -colorable and every circular r -coloring ϕ of (G, σ) has a tight cycle.

Tight cycle argument

Proposition

Any signed graph (G, σ) , which is not a forest, has a cycle with s positive edges and t negative edges such that

$$\chi_c(G, \sigma) = \frac{2(s + t)}{2a + t}$$

for some non-negative integer a .

Corollary

Given a signed graph (G, σ) on n vertices, $\chi_c(G, \sigma) = \frac{p}{q}$ for some $p \leq 2n$ and q .

- 1 Introduction
 - Circular coloring of graphs
 - Homomorphism of signed graphs
- 2 Circular coloring of signed graphs
 - Circular chromatic number
 - Signed indicators
 - Tight cycle argument
- 3 Results on some classes of signed graphs**
 - Signed bipartite planar graphs
 - Signed d -degenerate graphs
 - Signed planar graphs
- 4 Discussion

Classes of signed graphs

Given a class \mathcal{C} of signed graphs,

$$\chi_c(\mathcal{C}) = \sup\{\chi_c(G, \sigma) \mid (G, \sigma) \in \mathcal{C}\}.$$



- \mathcal{SBP} the class of signed bipartite planar simple graphs,
- \mathcal{SD}_d the class of signed d -degenerate simple graphs,
- \mathcal{SP} the class of signed planar simple graphs.

Signed bipartite planar graphs


Proposition

$$\chi_c(SBP) = 4.$$

Let Γ_1 be a positive 2-path connecting u_1 and v_1 . For $i \geq 2$,





Even step



Odd step

Signed bipartite planar graphs

Figure: Γ_4 Figure: Γ_5

Lemma

$$\chi_c(\Gamma_n) = \frac{4n}{n+1}.$$

Results on signed bipartite planar graphs with girth condition

- $\chi_c(\mathcal{SBP}_6) \leq 3$. (Corollary of a result that every signed bipartite planar graph of negative girth 6 admits a homomorphism to $(K_{3,3}, M)$ [R. Naserasr and Z. Wang 2021+])
- $\chi_c(\mathcal{SBP}_8) \leq \frac{8}{3}$. (Corollary of a result that C_{-4} -critical signed graph has density $|E(G)| \geq \frac{3|V(G)|-2}{4}$ [R. Naserasr, L-A. Pham and Z. Wang 2020+])

Signed d -degenerate graphs

Proposition

For any positive integer d , $\chi_c(\mathcal{SD}_d) = 2\lfloor \frac{d}{2} \rfloor + 2$.

Sketch of the proof:

- First we show that every $(G, \sigma) \in \mathcal{SD}_d$ admits a circular $(2\lfloor \frac{d}{2} \rfloor + 2)$ -coloring.


For the tightness,

- For odd integer d , we consider the signed complete graphs $(K_{d+1}, +)$.
- For $d = 2$, we consider the signed graph Γ_n built before.
- For even integer $d \geq 4$, we construct a signed d -degenerate graph (G, σ) such that $\chi_c(G, \sigma) = d + 2$.

Signed d -degenerate graphs

Proof for even $d \geq 4$

- Define a signed graph Ω_d as follows.



- Let φ be a circular r -coloring of Ω_d where $r < d + 2$. Without loss of generality, $\varphi(x_1), \dots, \varphi(x_d)$ are cyclically ordered on C^r and assume that $d_{(\text{mod } r)}(\varphi(x_1), \varphi(x_2))$ is maximized. We prove that there is no place for $y_{1,1+\frac{d}{2}}$.

Signed planar graphs

Proposition

$$4 + \frac{2}{3} \leq \chi_c(\mathcal{SP}) \leq 6.$$




Figure: Mini-gadget (T, π)




Figure: A signed Wenger Graph \mathbb{W}

Signed planar graphs

$\ell_{\phi;u,v}$: the minimum length of an interval which contains $\phi(u) \cup \phi(v)$.

Lemma

Let $r = \frac{14}{3} - \epsilon$ with $0 < \epsilon \leq \frac{2}{3}$. For any circular r -coloring ϕ of \tilde{W} , $\ell_{\phi;u,v} \geq \frac{4}{9}$.

Let Γ be obtained from \tilde{W} by adding a negative edge uv . Let $\mathcal{I} = (\Gamma, u, v)$.

Theorem

Let $\Omega = K_4(\mathcal{I})$. Then Ω is a signed planar simple graph with $\chi_c(\Omega) = \frac{14}{3}$.

Sketch of the proof of the theorem

- First we show that Ω admits a circular $\frac{14}{3}$ -coloring. We find a circular $\frac{14}{3}$ -coloring ϕ of Γ such that $\phi(u) = \phi(v) = 0$ and then extend it to each of inner triangles.
- Let ϕ be a circular r -coloring of Ω for $r < \frac{14}{3}$. For any $1 \leq i < j \leq 4$, $\frac{4}{9} \leq d_{(\text{mod } r)}(\phi(v_i), \phi(v_j)) \leq \frac{r}{2} - 1$. Assume that $\phi(x_1), \phi(x_2), \phi(x_3), \phi(x_4)$ are on C^r in this cyclic order.
 - $l([\phi(v_1), \phi(v_4)]) = l([\phi(v_1), \phi(v_2)]) + l([\phi(v_2), \phi(v_3)]) + l([\phi(v_3), \phi(v_4)]) \geq 3 \times \frac{4}{9} = \frac{4}{3} > \frac{r}{2} - 1$,
 - $l([\phi(v_4), \phi(v_1)]) \geq r - (l([\phi(v_1), \phi(v_3)]) + l([\phi(v_2), \phi(v_4)])) \geq 2 > \frac{r}{2} - 1$.

Contradiction.

Results on signed planar graphs with girth condition

- $\chi_c(\mathcal{SP}_4) = 4$. (By the 3-degeneracy of triangle-free planar graph)
- $\chi_c(\mathcal{SP}_7) \leq 3$. (Corollary of a result that every signed graph of $mad < \frac{14}{5}$ admits a homomorphism to (K_6, M) [R. Naserasr, R. Škrekovski, Z. Wang and R. Xu 2020+])

Signed circular chromatic number

For a simple graph G , the **signed circular chromatic number** $\chi_c^s(G)$ of G is defined as

$$\chi_c^s(G) = \max\{\chi_c(G, \sigma) : \sigma \text{ is a signature of } G\}.$$

Proposition

For every graph G , $\chi_c^s(G) \leq 2\chi_c(G)$.

Signed chromatic number of k -chromatic graph

Theorem

For any integers $k, g \geq 2$ and any $\epsilon > 0$, there is a graph G of girth at least g satisfying that $\chi(G) = k$ and $\chi_c^s(G) > 2k - \epsilon$.

We will prove that for any integer p , there is a graph G for which the followings hold:

- G is of girth at least g and has chromatic number at most k .
- There is a signature σ such that (G, σ) is not $(2kp, (p + 1))$ -colorable.

Augmented tree


- A **complete k -ary tree** is a rooted tree in which each non-leaf vertex has k children and all the leaves are of the same level.
- A **q -augmented k -ary tree** is obtained from a complete k -ary tree by adding, for each leaf v , q edges connecting v to q of its ancestors. These q edges are called the **augmenting edges** from v .
- For positive integers k, q, g , a **(k, q, g) -graph** is a q -augmented k -ary tree which is bipartite and has girth at least g .

Lemma [Alon, N., Kostochka, A., Reiniger, B., West, D., and Zhu, X 2016]

For any positive integers $k, q, g \geq 2$, there exists a (k, q, g) -graph.

Augmented tree

- A standard labeling of a complete k -ary tree T ;
- A f -path P_f of T with respect to a given k -coloring f .




Construction of k -chromatic graph G

- H : $(2kp, k, 2kg)$ -graph with underline tree T .
- ϕ : a standard $2kp$ -labeling of the edges of T .
- $\ell(v)$: the level of v , i.e., the distance from v to the root vertex in T . Let $\theta(v) = \ell(v) \pmod k$.

For each leaf v of T , let $u_{v,1}, u_{v,2}, \dots, u_{v,k}$ be the vertices on P_v that are connected to v by augmenting edges. Let $u'_{v,i} \in P_v$ be the closest descendant of $u_{v,i}$ with $\theta(u'_{v,i}) = i$ and let $e_{v,i}$ be the edge connecting $u'_{v,i}$ to its child on P_v .

Let $s_{v,i} = \phi(e_{v,i})$ and let

- $A_{v,i} = \{s_{v,i}, s_{v,i} + 1, \dots, s_{v,i} + p\}$,
- $B_{v,i} = \{a + kp : a \in A_{v,i}\}$,
- $C_{v,i} = A_{v,i} \cup B_{v,i}$.



Construction of the signature σ on G

Note that $B_{v,i}$ is a kp -shift of $A_{v,i}$. Two possibilities:

- $A_{v,i} \cap A_{v,j} \neq \emptyset$ (then $B_{v,i} \cap B_{v,j} \neq \emptyset$)

$$d_{(\text{mod } 2kp)}(\phi(e_{v,i}), \phi(e_{v,j})) \leq p.$$

- $A_{v,i} \cap B_{v,j} \neq \emptyset$ (then $B_{v,i} \cap A_{v,j} \neq \emptyset$)

$$d_{(\text{mod } 2kp)}(\phi(e_{v,i}), \overline{\phi(e_{v,j})}) \leq p.$$

Let L be the set of leaves of T . For each $v \in L$, we define one edge e_v on $V(T)$ as follows:

- If $d_{(\text{mod } 2kp)}(\phi(e_{v,i}), \phi(e_{v,j})) \leq p$, then let e_v be a positive edge connecting $u'_{v,i}$ and $u'_{v,j}$.
- If $d_{(\text{mod } 2kp)}(\phi(e_{v,i}), \overline{\phi(e_{v,j})}) \leq p$, then let e_v be a negative edge connecting $u'_{v,i}$ and $u'_{v,j}$.

Proof for “ (G, σ) is not circular $\frac{2kp}{p+1}$ -colorable”

Let (G, σ) be the signed graph with vertex set $V(T)$ and with edge set $\{e_v : v \in L\}$, where the signs of the edges are defined as above.

- Assume f is a $(2kp, p+1)$ -coloring of (G, σ) .
- As f is also a $2kp$ -coloring of the vertices of T , there is a unique f -path P_v . Assume that $e_v = u'_{v,i}u'_{v,j}$. By definition,

$$f(u'_{v,i}) = \phi(e_{v,i}) \text{ and } f(u'_{v,j}) = \phi(e_{v,j}).$$

- If e_v is a positive edge, then $d_{(\text{mod } 2kp)}(\phi(e_{v,i}), \phi(e_{v,j})) \leq p$.
If e_v is a negative edge, then $d_{(\text{mod } 2kp)}(\phi(e_{v,i}), \overline{\phi(e_{v,j})}) \leq p$.
Contradiction.

Mapping signed graphs to signed cycles

Let C_ℓ^{o+} be signed cycle of length ℓ where the number of positive edges is odd. Then $\chi_c(C_\ell^{o+}) = \frac{2\ell}{\ell-1}$.

Theorem

Given a positive integer ℓ and a signed graph (G, σ) satisfying $g_{ij}(G, \sigma) \geq g_{ij}(C_\ell^{o+})$ for $ij \in \mathbb{Z}_\ell^2$, we have $\chi_c(G, \sigma) \leq \frac{2\ell}{\ell-1}$ if and only if $(G, \sigma) \rightarrow C_\ell^{o+}$.

Circular chromatic number of signed planar graphs

Question

Given a positive integer ℓ , what is the smallest value $f(\ell)$ (with $f(\infty) = \infty$) such that for every signed planar graph (G, σ) satisfying $g_{ij}(G, \sigma) \geq g_{ij}(C_\ell^{o+})$ and $g_{ij}(G, \sigma) \geq f(\ell)$ for all $ij \in \mathbb{Z}_2^2$, we have $\chi_c(G, \sigma) \leq \frac{2\ell}{\ell - 1}$.

Jaeger-Zhang conjecture

When $\ell = 2k + 1$,

Jaeger-Zhang conjecture [C.-Q. Zhang 2002]

Every planar graph of odd-girth $f(2k + 1) = 4k + 1$ admits a circular $\frac{2k+1}{k}$ -coloring, i.e., C_{2k+1} -coloring.

- $f(3) = 5$ [Grötzsch's theorem];
- $f(5) \leq 11$ [Z. Dvořák and L. Postle 2017][D. W. Cranston and J. Li 2020];
- $4k + 1 \leq f(2k + 1) \leq 6k + 1$ [C. Q. Zhang 2002; L. M. Lovász, C. Thomassen, Y. Wu and C. Q. Zhang 2013];

Bipartite analogue of Jaeger-Zhang conjecture

When $\ell = 2k$,

Bipartite analogue of Jaeger-Zhang conjecture

Every signed bipartite planar graph of negative-girth $f(2k)$ admits a circular $\frac{4k}{2k-1}$ -coloring, i.e., C_{-2k} -coloring.

- $f(4) = 8$ [R. Naserasr, L. A. Pham and Z. Wang 2020+];
($f(2k) > 4k - 2$ when $k = 2$.)
- $f(2k) \leq 8k - 2$ [C. Charpentier, R. Naserasr and E. Sopena 2020].

Odd-Hadwiger Conjecture

Theorem [P.A. Catlin 1979]

If $(G, -)$ has no $(K_4, -)$ -minor, then $\chi_c(G, +) \leq 3$.

The Odd-Hadwiger conjecture was proposed independently by B. Gerard and P. Seymour.

Odd-Hadwiger conjecture

If a signed graph $(G, -)$ has no $(K_{k+1}, -)$ -minor, then $\chi_c(G, +) \leq k$.

Question

Assuming (G, σ) has no $(K_{k+1}, -)$ -minor, what is the best upper bound on $\chi_c(G, -\sigma)$?

The end. Thank you!